



# Effect of Sodium Nitrate ( $\text{NaNO}_3$ ) on Sperm Motility and Abnormality: An *In vitro* Approach

Rajini SV<sup>1</sup>, Chaithra B<sup>2</sup> and Shivabasavaiah<sup>1\*</sup>

<sup>1</sup>Department of Studies in Zoology, University of Mysore, India

<sup>2</sup>Department of Zoology, Vijaya College, Bangalore, India

\*Corresponding author: Shivabasavaiah, Professor, Department of Studies in Zoology, University of Mysore, Manasagangotri, Mysore 570006, Karnataka, India; Email: shivabasavaiahuom@gmail.com

## Research Article

Volume 6 Issue 2

Received Date: February 10, 2021

Published Date: April 28, 2021

DOI: 10.23880/act-16000211

## Abstract

Nitrates ( $\text{NO}_3$ ) are the naturally occurring inorganic ions with beneficial effects in both plants and animals. However,  $\text{NO}_3$  in excess has adverse effect on male and female reproductive systems. The study was undertaken to investigate the effect of 10mg/ml and 100mg/ml of sodium nitrate ( $\text{NaNO}_3$ ) on sperm motility and abnormality at different time intervals (5,10,15,20,25 and 30 min). Treatment of cauda epididymal sperm suspension with 10mg/ml of  $\text{NaNO}_3$  significantly reduced the sperm motility only at 25 and 30 min time intervals compared to controls. However, 100mg/ml of  $\text{NaNO}_3$  resulted in significant reduction of sperm motility from 5 min to 30 min time interval in time dependent manner and complete loss of sperm motility was observed at 15 min of incubation. In addition, the number of abnormal spermatozoa was significantly high in 100mg/ml of  $\text{NaNO}_3$  treated groups compared to controls and 10 mg/ml of  $\text{NaNO}_3$ . These results clearly reflected the toxic effect of  $\text{NO}_3$  on spermatozoa and its capability in fertilization.

**Keywords:** Abnormality; *In Vitro*; Motility; Sodium nitrate; Spermatozoa

## Introduction

Nitrate ( $\text{NO}_3$ ) and nitrite ( $\text{NO}_2$ ) are the naturally occurring inorganic ions involved in nitrogen cycle. There are numerous sources of nitrates which can be classified into exogenous sources and endogenous sources. Leafy vegetables are the exogenous sources of nitrates which account for more than 70% of nitrates ingested in human diet. Nitrites are produced endogenously through the oxidation of nitric oxide (NO) and through a reduction of  $\text{NO}_3$  by commensal bacteria in mouth and gastrointestinal tract [1]. In biological system,  $\text{NO}_3$  gets converted to  $\text{NO}_2$  and NO and are interchangeable [2].

Besides having beneficial effect,  $\text{NO}_3$  has adverse effects on the body. The connection between human health and harmful effect of  $\text{NO}_3$  was first reported by Comly in 1945

after observing cyanosis in residents of Iowa (USA). The primary effect of  $\text{NO}_3$  is the formation of methemoglobinemia in infants which reduces capacity to release oxygen to tissues due to oxidation of iron group of haem group [3,4]. Further,  $\text{NO}_3$  causes improper thyroid functioning via inhibition of sodium iodide symporter [5]. Increase in  $\text{NO}_3$  concentration causes myocardial infraction and hypotension [6]. And also affects liver by increasing bilirubin and transaminase levels [7].

Numerous studies have reported the toxic effect of  $\text{NO}_3$  on reproductive system [8-12]. It is evident that  $\text{NO}_3$  is a potential endocrine disruptor of reproductive endocrinology [13]. The adverse effects of  $\text{NO}_3$  on reproductive system viz, mummified foetuses, lesions on the cervix, uterus and placenta and maternal death were observed in  $\text{NO}_3$

exposed guinea pigs [14]. Further, higher risk of ovarian cancers in women drinking  $\text{NO}_3$  contaminated water [15] and incidences of abortions in potassium nitrate ( $\text{KNO}_3$ ) administered pregnant cows [16,17] were reported. In male rat and mice models, treatment of different doses of sodium nitrate ( $\text{NaNO}_3$ ), sodium nitrite ( $\text{NaNO}_2$ ) and  $\text{KNO}_3$  caused a significant reduction in weight of testis and epididymis, sperm count and sperm motility and increased sperm abnormality [9-12]. In addition  $\text{NO}_3$  causes lesions in spermatocytes and spermatids of germ layers, degenerations of leydig cells and arrest of spermatogenesis [9,10]. Further,  $\text{NO}_3$  affects steroidogenesis of male reproductive system by decreasing the activities of  $3\beta\text{HSD}$  (3-beta-hydroxysteroid dehydrogenase) and  $17\beta\text{HSD}$  (17-beta-hydrosteroid dehydrogenase) and serum concentration of testosterone [10,11,18].

Besides *in vivo* studies, few *in vitro* studies have focused on the toxic effect of NO on human spermatozoa. For instance, treatment of NO declined forward progressive movement of spermatozoa in *in vitro* [19]. Studies have reported that the reduced sperm DNA integrity with the association and high concentration of NO in seminal plasma [20]. In addition, high concentration of NO was reported in semen samples of patients suffering from asthenozoospermia than that of normozoospermic individuals [21]. Further, higher concentration of NO caused decreased zona binding capacity of sperm and embryonic development [22]. These results clearly indicate that NO has adverse effect on sperm functions. Though there are enough reports on effect of NO on human spermatozoa, the studies related to the action of  $\text{NO}_3$  on spermatozoa in *in vitro* is lacking. Hence, there is a need for investigation to assess the toxic effect of  $\text{NaNO}_3$  (source of  $\text{NO}_3$ ) on motility and abnormality of spermatozoa at different time intervals. with this background, the present study was conducted to investigate the dose and time dependent effect of two different concentrations of  $\text{NaNO}_3$  (10mg/ml and 100mg/ml) on sperm motility and sperm abnormality in *in vitro*.

## Materials and Methods

### Animals

The study included adult male wistar rats weighing 180-200g procured from central animal facility of University of Mysore, Mysore. Polypropylene cages were used to maintain the animals under standard laboratory conditions and were provided with rat chow and water *ad libitum* and relative 12h light/dark cycle. The ethical acceptance to conduct the experiment was obtained from Institutional Animal Ethics Committee of University of Mysore, India (reference number: UOM/IAEC/04/2018) and the study was carried out as per the guidelines of the committee.

## Experimental Design

An *in vitro* experiment was conducted to assess the effects of two different concentrations of  $\text{NaNO}_3$  on sperm motility and abnormality at different time intervals viz., 0, 5, 10, 15, 20, 25 and 30 min. The epididymal sperm suspension (60 million/ml) was incubated with different concentrations of  $\text{NaNO}_3$  (10mg/ml and 100mg/ml) at different time intervals. The group devoid of  $\text{NaNO}_3$  was treated as control. After incubation, sperm motility and sperm abnormalities were analysed at regular time intervals.

### Isolation of spermatozoa

Epididymal spermatozoa were isolated from cauda epididymis of adult male rats. The epididymis was minced in 1 ml of phosphate buffered saline and the suspension was filtered through muslin cloth. The filtered sperm suspension was used for the *in vitro* experiment.

### Estimation of sperm motility

Progressive sperm motility was considered for estimating sperm motility. Motility was estimated by placing the sperm suspension on a slide and number of motile spermatozoa was counted from three different microscopic fields under light microscope. The mean of the three estimations was calculated in percentage [23].

### Estimation of sperm abnormality

Sperm abnormality was analysed by staining the sperm suspension with eosin and a uniform smear was made on a glass slide. One thousand spermatozoa were observed under higher magnification (40X) and the number of spermatozoa showing head and tail abnormalities were counted. The aggregate of different types of spermatozoa showing abnormality were considered to compute percentage of spermatozoa with abnormal morphology [24,25].

### Statistical Analysis

The mean  $\pm$  standard error of each parameter was computed by considering the data and the mean values of each parameter of different groups were compared using one way analysis of variance followed by Duncan's multiple range test and judged significant if  $p < 0.05$ .

## Results

### Effect of different concentration of $\text{NaNO}_3$ on sperm motility at 0, 5, 10, 15, 20, 25 and 30min intervals

There was a significant decrease in sperm motility in

sperm suspension treated with 100 mg/ml of NaNO<sub>3</sub> at 5, 10, 15, 20, 25 and 30 min of time interval compared to control. However, 10mg/ml of NaNO<sub>3</sub> significantly decreased the sperm motility only at 25 and 30 min time interval compared to control. No significant changes in sperm motility was

observed at 5, 10, 15 and 20 min time interval of 10mg/ml of NaNO<sub>3</sub> treated groups compared to controls. Treatment of 100mg/ml of NaNO<sub>3</sub> caused complete loss of sperm motility at 15, 20, 25 and 30 min of incubation (Table 1).

Groups	Mean percentage of motility ± SE						
	0 min	5 min	10 min	15 min	20 min	25 min	30 min
Control	91.00±3.21 <sup>a</sup>	91.33±2.72 <sup>a</sup>	94.33±2.33 <sup>a</sup>	89.00±3.05 <sup>a</sup>	76.66±2.40 <sup>a</sup>	67.33±1.76 <sup>a</sup>	45.33±3.711 <sup>a</sup>
10mg/ml of NaNO <sub>3</sub>	92.33±1.76 <sup>a</sup>	95.33±0.88 <sup>a</sup>	91.66±0.88 <sup>a</sup>	82.00±3.05 <sup>a</sup>	69.33±2.96 <sup>a</sup>	45.66±2.33 <sup>b</sup>	27.66±4.33 <sup>b</sup>
100mg/ml of NaNO <sub>3</sub>	91.00±2.64 <sup>a</sup>	37.66±6.48 <sup>b</sup>	15.00±5.13 <sup>b</sup>	0.00±0.00 <sup>b</sup>	0.00±0.00 <sup>b</sup>	0.00±0.00 <sup>c</sup>	0.00±0.00 <sup>c</sup>
ANOVA F value(df=2,6)	0.87 *p>0.918	84.31 ***p<0.001	187.04 ***p<0.001	393.58 ***p<0.001	368.88 ***p<0.001	414.27 ***p<0.001	48.11 ***p<0.001

**Table 1:** Effect of different concentration of NaNO<sub>3</sub> on sperm motility.

**Note:** Mean values were compared by one-way ANOVA followed by Duncan's multiple range tests. Values with same superscript letters in the given column are not significantly different, whereas those with different superscript letters are significantly (\*p<0.05) different from each other. df - degree of freedom, NaNO<sub>3</sub> - sodium nitrate.

### Effect of different concentration of NaNO<sub>3</sub> on sperm abnormality at 0, 5, 10, 15, 20, 25 and 30 min interval

A significant increase in sperm abnormality was

observed in 100mg/ml of NaNO<sub>3</sub> treated groups compared to that of controls and 10mg/ml of NaNO<sub>3</sub>. However, no significant difference in sperm abnormality was observed between control groups and 10mg/ml of NaNO<sub>3</sub> with increase in duration of incubation (Table 2).

Groups	Mean percentage of abnormality ± SE						
	0 min	5 min	10 min	15 min	20 min	25 min	30 min
Control	0.00±0.00 <sup>a</sup>	0.66±0.33 <sup>a</sup>	0.66±0.33 <sup>a</sup>	0.66±0.33 <sup>a</sup>	0.66±0.33 <sup>a</sup>	0.66±0.33 <sup>a</sup>	0.66±0.33 <sup>a</sup>
10mg/ml of NaNO <sub>3</sub>	0.00±0.00 <sup>a</sup>	1.33±0.33 <sup>a</sup>	1.66±0.66 <sup>a</sup>	2.00±0.57 <sup>a</sup>	2.66±1.20 <sup>a</sup>	3.00±0.57 <sup>a</sup>	3.33±0.33 <sup>a</sup>
100mg/ml of NaNO <sub>3</sub>	1.00±0.57 <sup>a</sup>	11.00±1.15 <sup>b</sup>	14.66±1.76 <sup>b</sup>	23.66±3.48 <sup>b</sup>	23.66±3.48 <sup>b</sup>	23.66±3.48 <sup>b</sup>	23.66±3.48 <sup>b</sup>
ANOVA F value (df=2,6)	3 *p<0.125	64.5 ***p<0.001	49.9 ***p<0.001	39.83 ***p<0.001	35.63 ***p<0.001	38.29 ***p<0.001	38.49 ***p<0.001

**Table 2:** Effect of different concentration of NaNO<sub>3</sub> on sperm abnormality.

**Note:** Mean values were compared by one-way ANOVA followed by Duncan's multiple range tests. Values with same superscript letters in the given column are not significantly different, whereas those with different superscript letters are significantly (\*p<0.05) different from each other. df - degree of freedom, NaNO<sub>3</sub> - sodium nitrate.

## Discussion

Present study was undertaken to investigate the toxic effect of two different concentration of NaNO<sub>3</sub> on sperm motility and sperm abnormality at different time intervals. The study revealed that treatment of NaNO<sub>3</sub> decreased the sperm motility and increased the number of abnormal spermatozoa compared to controls.

Sperm motility is an important aspect of spermatozoa to reach the fallopian tube during fertilization. The quality

of spermatozoa depends on its motility and number of abnormal spermatozoa [26]. Earlier *in vivo* studies in rat and mice showed a decline in sperm motility under NO<sub>3</sub> treatment [9-12]. Similarly, present study showed NO<sub>3</sub> induced reduced sperm motility in *in vitro* condition. Between the two doses of NaNO<sub>3</sub>, a higher concentration that is 100mg/ml showed severe toxicity as it resulted in complete cessation of sperm motility at 15 min of incubation with epididymal spermatozoa. However, the lower dose of NaNO<sub>3</sub>, that is 10mg/ml showed a time dependent decrease in sperm motility from 5 min to 30 min compared to controls.

A significant dose dependent effect was observed between two doses of  $\text{NaNO}_3$  at different durations of incubations studied. Thus, the present study clearly demonstrates that treatment of  $\text{NaNO}_3$  affected sperm motility in dose and time dependent manner. Similarly, earlier study have reported the dose dependent inhibition of sperm motility by NO donors viz, nitroprusside and pure NO gas [27]. In contrast, few investigations reported the beneficial aspects of  $\text{NO}_3$  on spermatozoa wherein treatment with  $\text{NO}_3$  caused enhanced capacitation [28], motility, viability [29] and vigour [30] of spermatozoa. In addition, it has been reported that NO and nitric oxide synthase (NOS) is essential for capacitation and fertilization [31].

The  $\text{NO}_3$  induced decrease in sperm motility is due to increased level of NO, a product of  $\text{NO}_3$  and  $\text{NO}_2$  [2]. Higher concentration of  $\text{NO}_3$  enhances the synthesis of NO which in turn affects sperm motility. The peroxy nitrate ( $\text{ONOO}^-$ ) formed due to the interaction between NO and reactive oxygen species (ROS) can cause damage to lipids and thiol proteins of sperm membrane [32] thereby affects sperm motility. In addition, NO inhibit sperm motility by affecting mitochondrial electron transport protein and thereby obstruct cellular respiration and ATP production in spermatozoa [27,33]. Further,  $\text{NO}_3$  inhibit sperm motility by inducing oxidative stress [34]. Spermatozoa are very sensitive to ROS and  $\text{ONOO}^-$  which can readily damage the cell membrane [32,34]. These are the possible mechanisms through which  $\text{NO}_3$  can affect sperm motility.

The morphology of spermatozoa is another important aspect which plays a vital role in fertilization. Earlier *in vivo* studies have proved enhanced number of abnormal spermatozoa under  $\text{NO}_3$  treatment [9-12]. In the present *in vitro* study, treatment of 100mg/ml of  $\text{NaNO}_3$  caused a significant increase in the number of abnormal spermatozoa compared to 10mg/ml of  $\text{NaNO}_3$  and control groups. The  $\text{NaNO}_3$  induced increase in abnormal spermatozoa increased with increasing duration of exposure. These results clearly indicate that  $\text{NaNO}_3$  affect the sperm morphology by increasing head and tail abnormalities, thereby affect the fertilization capacity of spermatozoa. The  $\text{NO}_3$  induced sperm abnormality may be due to obstruction in cellular respiration and production of ATP and formation of ROS which can cause damage to the lipid and thiol proteins present in the sperm membrane [27,32-34]. Therefore,  $\text{NO}_3$  exposure increases the number of abnormal spermatozoa by affecting cell membrane.

The present *in vitro* study clearly demonstrated that  $\text{NaNO}_3$  has adverse effect on spermatozoa and the effect is time and dose dependent. The study reports the toxic effects of  $\text{NaNO}_3$  on spermatozoa in *in vitro* condition for the first time as earlier *in vitro* studies were focused on effect NO

on human spermatozoa [19,21,27]. The study also provides evidence that higher concentration of  $\text{NaNO}_3$  that is 100mg/ml has severe effect on sperm motility and abnormality. These results gain importance in male infertility of human beings residing in the areas where there is high level of  $\text{NO}_3$  in drinking ground water. Therefore, present investigation appears to be an evidence for the effect of  $\text{NO}_3$  on male reproduction. However, future studies are needed to reflect mechanistic action of  $\text{NO}_3$  in affecting sperm motility.

## Conclusion

The present study demonstrated dose and time dependent effect of  $\text{NO}_3$  on spermatozoa. The higher dose of  $\text{NaNO}_3$  that is 100mg/ml adversely affected sperm motility and abnormality compared to lower dose. The results of the present study for the first time showed reproductive toxicity of  $\text{NaNO}_3$  in *in vitro* condition.

## Conflicts of Interest

The authors declare no conflicts of interest.

## Acknowledgements

The first author is thankful to BCWD (Backward classes welfare department) Government of Karnataka. India.

## References

1. Hord NG, Tang Y, Bryan NS (2009) Food sources of nitrates and nitrites: the physiologic context for potential health benefits. *Am J Clin Nutr* 90(1): 1-10.
2. Lundberg J, Weitzberg E, Gladwin M (2008) The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 7(2): 156-167.
3. Nishiguchi M, Nushida H, Noriyuki O (2014) An Autopsy Case of Fatal Methemoglobinemia due to Ingestion of Sodium Nitrite. *J Forensic Res* 6(1): 1-3.
4. Ward MH, Jones RR, Brender JD, de Kok TM, Weyer PJ, et al. (2018) Drinking Water Nitrate and Human Health: An Updated Review. *Int J Environ Res Public Health* 15(7): 1557.
5. Drozd VM, Saenko VA, Brenner AV, Drozdovitch V, Pashkevich VI, et al. (2015) Major Factors Affecting Incidence of Childhood Thyroid Cancer in Belarus after the Chernobyl Accident: Do Nitrates in Drinking Water Play a Role? *PLoS One* 10(9): e0137226.
6. RuDusky BM (2001) Acute Myocardial Infarction Secondary to Coronary Vasospasm during Withdrawal from Industrial Nitroglycerin Exposure: A Case

- Report. *Angiology* 52(2): 143-144.
7. Bensoltane S, Messerer L, Youbi M, Houria B, Djekoun M, et al. (2006) Effects of acute and sub-chronic ammonium nitrate exposure on rat liver and blood tissues. *Afri J of Biotechnol* 5(9): 749-754.
  8. Adalakun SA, Ukwenya VO, Ogunlade BS, Aniah AJ, Ibiayo GA (2019) Nitrite-induced testicular toxicity in rats: therapeutic potential of walnut oil. *JBRA Assist Reprod* 23(1): 15-23.
  9. Aly HA, Mansour AM, Abo Salem OM, Abd-Ellah HF, Abdel-Naim AB (2010) Potential testicular toxicity of sodium nitrate in adult rats. *Food Chem Toxicol* 48(2): 572-578.
  10. Azza ME, El-Said ME, Waffa ME (2011) Evidence for Androgenic Activity of turmeric and curcumin in male rats exposed to water nitrate pollution. *Am J Sci* 7(9): 1016-1026.
  11. Bansode FW, Singh RK (2017) The reproductive toxicity assessment of Potassium nitrate in Sprague Dawley Rats. *Int J Innov Res Sci Eng Technol* 6(7): 15359-15370.
  12. Pant N, Srivastava S (2002) Testicular and spermatotoxic effect of nitrate in mice. *Hum Exp Toxicol* 21(1): 37-41.
  13. Guillette LJ, Edwards TM (2005) Is nitrate an ecologically relevant endocrine disruptor in vertebrates? *Integr Comp Biol* 45(1): 19-27.
  14. Sleight SD, Atallah OA (1968) Reproduction in the guinea pig as affected by chronic administration of potassium nitrate and potassium nitrite. *Toxicol Appl Pharmacol* 12(2): 179-85.
  15. Inoue-Choi M, Jones RR, Anderson KE, Cantor KP, Cerhan JR, et al. (2015) Nitrate and nitrite ingestion and risk of ovarian cancer among postmenopausal women in Iowa. *Int J Can* 137(1): 173-182.
  16. Case AA (1957) Some aspects of nitrate intoxication in livestock. *J Am Vet Med Assoc* 130(8): 323-329.
  17. Simon J, Sund JM, Douglas FD, Wright MJ, Kowalczyk T (1959) The effect of nitrate or nitrite when placed in the rumens of pregnant dairy cattle. *J Am vet Med Assoc* 135: 311-314.
  18. Panesar NS, Chan KW (2001) Decreased steroid hormone synthesis from inorganic nitrite and nitrate: Studies in vitro and in vivo. *Toxicol Appl Pharmacol* 169(3): 222-230.
  19. Rosselli M, Dubey RK, Imthurn B, Macas E, Keller PJ (1995) Effects of nitric oxide on human spermatozoa: evidence that nitric oxide decreases sperm motility and induces sperm toxicity. *Hum Reprod* 10(7): 1786-1790.
  20. Amiri I, Sheikh N, Najafi R (2007) Nitric oxide level in seminal plasma and its relation with sperm DNA damages. *Iran Biomed J* 11(4): 259-264.
  21. Balercia G, Moretti S, Vignini A, Magagnini M, Mantero F, et al. (2004) Role of nitric oxide concentrations on human sperm motility. *J Androl* 25(2): 245-249.
  22. Wu TP, Huang BM, Tsai HC, Lui MC, Liu MY (2004) Effects of nitric oxide on human spermatozoa activity, fertilization and mouse embryonic development. *Arch Androl* 50(3): 173-179.
  23. Saalu LC, Kpela T, Benebo A, Oyewopo AO, Anifowope EO, et al. (2010) The dose dependent testiculoprotective and testiculotoxic potentials of *Telfairia occidentalis* hook f. leaves extract in rat. *Int J Appl Res Nat Prod* 3: 27-28.
  24. Narayana K, D'Souza UJ, Seetharama Rao KP (2002) Ribavirin-induced sperm shape abnormalities in Wistar rat. *Mutat Res* 513(1-2): 193-196.
  25. Vega SG, Guzman P, Garcia L, Espinosa J, Cortinas de Nava C (1988) Sperm shape abnormality and urine mutagenicity in mice treated with niclosamide. *Mutat Res* 204(2): 269-276.
  26. Au DW, Chiang MW, Tang JY, Yuen BB, Wang YL, et al. (2002) Impairment of sea urchin sperm quality by UV-B radiation: predicting fertilization success from sperm motility. *Mar Pollut Bull* 44(7): 583-589.
  27. Weinberg JB, Doty E, Bonaventura J, Haney AF (1995) Nitric oxide inhibition of human sperm motility. *Fertil Steril* 64(2): 408-413.
  28. Zini A, De Lamirande E, Gagnon C (1995) Low levels of nitric oxide promote sperm capacitation in vitro. *J Androl* 16(5): 424-431.
  29. Zhang H, Zheng RL (1996) Possible role of nitric oxide on fertile and asthen-ozoospermic infertile human sperm functions. *Free Rad Res* 25(4): 347-354.
  30. Maciel VL, Caldas-Bussiere MC, Marín DFD, Paes de Carvalho CS, Quirino CR, et al. (2019) Nitric oxide impacts bovine sperm capacitation in a cGMP-dependent and cGMP-independent manner. *Reprod Domest Anim* 54(12): 1612-1620.
  31. Staicu F, Lopez-Úbeda R, Romero-Aguirregomez-corta J, Martínez-Soto JC, Parra MC (2019) Regulation of boar sperm functionality by the nitric oxide synthase/nitric

- oxide system. *J Assist Reprod Genet* 36(8): 1721-1736.
32. De Pinto MC, Tommasi F, De Gara L (2002) Changes in the antioxidant systems as part of the signaling pathway responsible for the programmed cell death activated by nitric oxide and reactive oxygen species in tobacco Bright-Yellow 2 cells. *Plant Physiol* 130(2): 698-708.
  33. Mitchell JA, Nelson L, Hafez ESE (1976) Motility of spermatozoa In Human semen and fertility regulation in men. St. Louis: C.V. Mosby Company, pp: 83-99.
  34. Doshi SB, Khullar K, Sharma RK, Agarwal A (2012) Role of reactive nitrogen species in male infertility. *Reprod Biol Endocrinol* 15: 10-109.

