

Diagnosis	References
Various eczemas	1-4
Allergic contact dermatitis	5-6
Psoriasis	1,4, 7-9
Photodermatitis	10-12
Alopecia areata	13-15
Drug eruption	4, 16
Tinea corporis	17
Crusted scabies	1,18,19
Hidradenitis suppurativa	20
Burns	21-23

Supplementary Table 1: Dermatologic conditions other than atopic dermatitis reported to have an increased serum total IgE levels.

References

1. Przybilla B, Ring J, Völk M (1986) [Total serum IgE in dermatological diseases]. *Hautartz* 37(2): 77-82.
2. Krueger GG, Kahn G, Weston WL, Mandel MJ IgE (1973) levels in nummular eczema and ichthyosis. *Arch Dermatol* 107(1): 56-58.
3. O'Loughlin S, Diaz-Perez JL, Gleich GJ, Winkelmann RK (1977) Serum IgE in dermatitis and dermatosis: an analysis of 497 cases. *Arch Dermatol* 113(3): 309-315.
4. Kim TY, Park HJ, Kim CW (1997) Eosinophil cationic protein (ECP) level and its correlation with eosinophil number or IgE level of peripheral blood in patients with various skin diseases. *J Dermatol Sci* 15(2): 89-94.
5. Peltonen L, Havu VK, Mattila L Serum (1988) IgE in non-atopic adults and in dermatitis patients. *Allergy* 43(2): 152-1518.
6. Welke S, Schimpf A (1975) Serum IgE-concentrations in patients with contact sensitization. *Arch Dermatol Res* 253(3): 319-328.
7. Chen ZY, Ainsworth SK, Khan T, Pilia PA, Dobson RL (1985) Immunoglobulin E in psoriasis evaluated by paper radioimmunosorbent and paper enzyme-immunosorbent tests. *Acta Derm Venereol* 65(1): 14-18.
8. Ding Y, Yi X, Yu N (2013) Serum IgE levels are increased in patients with generalized pustular psoriasis. *Clin Exp Dermatol* 38(5): 549-552.
9. Paparo SB, Guaragna MA, Albanesi M High (2014) IgE levels in patients affected by psoriasis: review of the literature and personal observations. *Clin Ter* 165(2): 91-93.
10. Vestey JP, Addo H, MacFarlane HA, Barnetson RS (1985) Patients with the chronic photosensitivity dermatitis with actinic reticuloid syndrome have high total serum IgE concentrations.

Photodermatology 2(2): 124-125.

11. Ko DY, Choi SH, Ha SM, Kim TH, Song KH, et al. (2016) The clinical severity score of chronic actinic dermatitis correlates with in vivo photoallergic reactions and the immunologic parameters related to a shift towards Th2 immunity from the Th2/Th1 balanced status in patients with chronic actinic dermatitis. *Photodermatol Photoimmunol Photomed* 32(4): 199-206.
12. Cuevas Gonzalez JC, Lievanos Estrada Z, Vega Memije ME, HojyoTomoka MT, Dominguez Soto L(2016) Correlation of serum IgE levels and clinical manifestations in patients with actinic prurigo. *An Bras Dermatol* 91(1): 23-26.
13. KasumagićHalilović E, Prohić A (2006) Serum levels of total immunoglobulin E in patients with alopecia areata: relationship with clinical type of the disease. *Acta Dermatovenerol Croat* 14 (3): 149-152.
14. Zhao Y, Zhang B, Caulloo S, Chen X, Li Y, et al. (2012) Diffuse alopecia areata is associated with intense inflammatory infiltration and CD8+ T cells in hair loss regions and an increase in serum IgE level. *Indian J Dermatol Venereol Leprol* 78(6): 709-714.
15. Bakry OA, El Shazly RM, Basha MA, Mostafa H (2014) Total serum immunoglobulin E in patients with alopecia areata. *Indian Dermatol Online J* 5(2): 122-127.
16. Mészáros C, Debreczeni M, Mahunka M (1985) IgE level in some dermatological diseases. *Acta Med Hung* 42(3-4): 125-128.
17. Hay RJ, Shennan G (1984) Antibody responses in tinea imbricata: the role of immunoglobulin E. *Trans R Soc Trop Med Hyg* 78(5): 653-635.
18. Roberts LJ, Huffam SE, Walton SF, Currie BJ (2005) Crusted scabies: clinical and immunological findings in seventy-eight patients and a review of the literature. *J Infect* 50(5): 375-381.
19. Walton SF, Beroukas D, Roberts-Thomson P, Currie BJ (2008) New insights into disease pathogenesis in crusted (Norwegian) scabies: the skin immune response in crusted scabies. *Br J Dermatol* 158(6): 1247-1255.
20. Pascual JC, García-Martínez FJ, Martorell A, González I, Hispan P (2016) Increased total serum IgE levels in moderate-to-severe hidradenitis suppurativa. *Br J Dermatol* 175(5): 1101-1102.
21. Gleich GJ, Dunnette SL, Volenec FJ, Mani MM (1979) Quantification of serum IgE in patients with burns. *Clin Allergy* 9(2): 133-139.
22. Polacek V, Jira M, Fara M, Strejcek J, Konigova R (1987) Immunoglobulin E (IgE) in patients with severe burns. *Burns Incl Therm Inj* 13(6): 458-461.
23. Smith CJ, Payne VM, Scott SM, Luterman A (1997) Immunoglobulin E levels and anticollagen antibodies in patients with postburn hypertrophic scars. *J Burn Care Rehabil* 18(5): 411-416.