




Supplementary Figures

Index


Figures titles

- Supplementary Figure 1: Sub-nephrotic proteinuria as indication for renal biopsy regarding the reports' world region
- Supplementary Figure 2: Sub-nephrotic proteinuria as indication for renal biopsy regarding the reports' populations' range of age
- Supplementary Figure 3: MGN diagnosis among patients with sub-nephrotic proteinuria regarding the reports' world region
- Supplementary Figure 4: IgA nephropathy diagnosis among patients with sub-nephrotic proteinuria regarding the reports' world region
- Supplementary Figure 5: FSGS diagnosis among patients with sub-nephrotic proteinuria regarding the reports' world region
- Supplementary Figure 6: LUPUS NEPHRITIS diagnosis among patients with sub-nephrotic proteinuria regarding the reports' world region
- Supplementary Figure 7: MINIMAL CHANGE DISEASE diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 8: CRESCENTRIC NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 9: MPGN diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 10: AMYLOIDOSIS diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 11: DIABETIC NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 12: MesPGN diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 13: TUBULOINTERSTITIAL DIS. diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 14: VASCULAR NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 15: HEREDITARY NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 16: NEPHROANGIOSCLEROSIS diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 17: HENOCH-SCHÖNLEIN PURPURA diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 18: IgM NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 19: HBV NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 20: UNSPECIFIC PROLIFERATIVE GLOMERULONEPHRITIS in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 21: UNSPECIFIC PARAPROTEINEMIA diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 22: PEGN diagnosis in patients with sub-nephrotic proteinuria regarding the reports' region
- Supplementary Figure 23: MGN diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 24: IgA NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 25: FSGS diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 26: LUPUS NEPHRITIS diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 27: MINIMAL CHANGE DISEASE diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 28: CRESCENTRIC NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 29: MPGN diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 30: PEGN diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 31: AMYLOIDOSIS diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses


- Supplementary Figure 32: DIABETIC NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 33: MesPGN diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 34: TUBULOINTERSTITIAL DISEASES diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 35: VASCULAR NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 36: NEPHROANGIOSCLEROSIS diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 37: HEREDITARY NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 38: HENOCH-SCHÖNLEIN PURPURA diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 39: HBV NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 40: IgM NEPHROPATHY diagnosis in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 41: UNSPECIFIC PROLIFERATIVE GLOMERULONEPHRITIS in patients with sub-nephrotic proteinuria regarding the age subclasses
- Supplementary Figure 42: UNSPECIFIC PARAPROTEINEMIA in patients with sub-nephrotic proteinuria regarding the age subclasses




Supplementary Figure 1: Frequency of sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 2: Frequency of sub-nephrotic proteinuria as indication for renal biopsy regarding the report's populations' range of age.




Supplementary Figure 3: Frequency of MGN diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 4: Frequency of IgA nephropathy diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 5: Frequency of FSGS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 6: Frequency of LUPUS NEPHRITIS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 7: Frequency of MINIMAL CHANGE DISEASE diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 8: Frequency of CRESCENTRIC NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 9: Frequency of MPGN diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 10: Frequency of AMYLOIDOSIS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 11: Frequency of DIABETIC NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 12: Frequency of MesPGN diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 13: Frequency of TUBULOINTERSTITIAL DISEASES diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 14: Frequency of VASCULAR NEPHROPATHIES diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 15: Frequency of HEREDITARY NEPHROPATHIES diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 16: Frequency of NEPHROANGIOSCLEROSIS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 17: Frequency of HENOCH-SCHÖNLEIN PURPURA diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 18: Frequency of IgM NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 19: Frequency of HBV NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 20: Frequency of UNSPECIFIC PROLIFERATIVE GLOMERULONEPHRITIS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 21: Frequency of UNSPECIFIC PARAPROTEINEMIA diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 22: Frequency of PEGN diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the report's world region.




Supplementary Figure 23: Frequency of MGN diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 24: Frequency of IgA NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 25: Frequency of FSGS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 26: Frequency of LUPUS NEPHRITIS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 27: Frequency of MINIMAL CHANGE DISEASE diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 28: Frequency of CRESCENTRIC NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 29: Frequency of MPGN diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 30: Frequency of PEGN diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 31: Frequency of AMYLOIDOSIS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 32: Frequency of DIABETIC NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 33: Frequency of MesPGN diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 34: Frequency of TUBULOINTERSTITIAL DISEASES diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 35: Frequency of VASCULAR NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 36: Frequency of NEPHROANGIOSCLEROSIS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 37: Frequency of HEREDITARY NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 38: Frequency of HENOCHE-SCHÖNLEIN PURPURA diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.




Supplementary Figure 39: Frequency of HBV NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.



Supplementary Figure 40: Frequency of IgM NEPHROPATHY diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.



Supplementary Figure 41: Frequency of UNSPECIFIC PROLIFERATIVE GLOMERULONEPHRITIS diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.



Supplementary Figure 42: Frequency of UNSPECIFIC PARAPROTEINEMIA diagnosis among patients with sub-nephrotic proteinuria as indication for renal biopsy regarding the age subclasses.