



# Fine Needle Aspiration Cytology Findings of Breast Lesions in Female Patients Presenting With Palpable Breast Lumps at Makerere University College of Health Sciences, Kampala-Uganda

Sulaiman K<sup>1</sup>, Bary AD<sup>2</sup> and Boaz M<sup>3\*</sup>

<sup>1</sup>Department of Pathology, Makerere University, Uganda

<sup>2</sup>Faculty of Biology, University of Manchester, United Kingdom

<sup>3</sup>Cytology Society of Uganda, Uganda

Research Article

Volume 5 Issue 1

Received Date: January 29, 2021

Published Date: March 03, 2021

\*Corresponding author: Mwesigwa Boaz, Cytology Society of Uganda, Kampala, Uganda, Tel: +256705482996; Email: mwesigwaboaz155@gmail.com

## Abstract

**Background:** Palpable breast lumps are a common manifestation encountered by physicians during clinical practice. The use of fine-needle aspiration cytology (FNAC) method has proven to be quick, simple and cost-effective in the evaluation of these lumps for benign, atypical and malignant changes. This study aims to determine the type of breast lesions diagnosed by FNAC and to determine the age-wise distribution for such lesions.

**Methodology:** A prospective study consisting of 291 females presenting with palpable breast lesions was carried out at Makerere College of Health Sciences (MakCHS) pathology laboratory from January 2019 to May 2019. FNAC results were grouped into tier five reporting categories as C1, C2, C3, C4 & C5 in accordance to the International Academy of Cytology (IAC).

**Results:** Out of the 291 cases examined, 14 (4.8%) were insufficient (C1), 192 (66%) were benign (C2), 9 (3.0%) were atypical (C3), 15 (5.2%) were suspicious of malignancy (C4) & 61 (21%) were malignant (C5). In the benign category, lesions of fibroadenoma were the most commonly diagnosed constituting 110/192 (57.3%) whereas under the malignant category lesions of invasive ductal carcinoma were the most commonly diagnosed constituting 27/61(44.3%). The peak age group for benign lesions was 21-30 years whereas the peak age group for malignant lesions was 41-50 years.

**Conclusion:** Fine-needle aspiration cytology (FNAC) was found to be an effective diagnostic tool in the categorization of palpable breast lumps into benign, malignant, atypical, suspicious and inadequate categories.

**Keywords:** Breast cancer; Fine needle aspiration cytology; Palpable breast masses

**Abbreviations:** ADH: Atypical Ductal Hyperplasia; AUS: Atypia of Undetermined Significance; CBE: Clinical Breast Examination; FA: Fibroadenoma; FCC: Fibrocystic Changes; FNAC: Fine Needle Aspiration Cytology; IAC: International Academy of Cytology; IDC: Invasive Ductal Carcinoma; MAKCHS: Makerere University College of Health Sciences;

SPSS: Statistical Package for Social Sciences.

## Introduction

Breast lump is the localized swelling, protuberance or mass in the breast and is one of the prime indicators of

breast disorders. A lump might be a cyst, a benign tumour or a malignancy [1]. Symptom misinterpretation, misdiagnosis and late presentation at the primary health care level are reported as one of contributors to the increased of progression to breast cancer [2]. Globally, over 2 million females are diagnosed with breast cancer accounting for an excess of 626,000 deaths [3]. In the 2018, the age-standardized incidence rate and mortality rate for breast cancer in Uganda was 21.3 per 100,000 and 10.3 per 100,000 population, respectively [4]. These high mortality rates were attributed to a late stage disease diagnosis of women with breast cancer with up to 89% presenting with stage disease that was too difficult to treat [5]. The recommended approach for the early detection and diagnosis of breast lumps involves the use of triple assessment which includes; clinical breast examination (CBE), imaging (ultrasound or mammography) and fine-needle aspiration cytology [6]. Fine-needle aspiration cytology (FNAC) has been in use since the 1930s and is simple, quick and a cost-effective method of diagnosing palpable breast lumps [7]. Whereas this method offers a reliable means of diagnosis during the assessment of breast diseases, its utility in this country is restricted to a few public or private health institutions and limited data is available regarding the prevalence of breast diseases in Uganda. The aim of this study was to document the type of breast diseases diagnosed by FNAC at the MakCHS pathology laboratory over a period of 5 months (January –May, 2019). Furthermore this study aimed at determining the age-wise distribution of breast diseases at the pathology laboratory over the said period.

## Materials and Methods

### Study Design

The study was a cross-sectional prospective study.

### Study Area

This study was conducted at the cytology clinic of the department pathology of Makerere University College of Health Sciences (MakCHS). It is located at Mulago National Referral and Teaching Hospital in Kampala. It is one of fewest public laboratory that offers cytology services.

### Study Duration

The study was carried out in five months starting from January 2019 to May 2019.

### Study Population

The study population consisted of all female patients

that presented with palpable breast lumps.

### Sample Size Estimation

For calculation of sample size, the formula below was used [8];

$$n = Z^2 P(1-P)/d^2$$

where Z is the value of 1.96 (z in normal distribution curve), n is the required sample size, p is the estimated prevalence, and d is precision error (5%). A prevalence p of 25.5% was used in the estimation of sample size, n. Using this equation, a total sample size of 291 individuals was computed [9].

### Selection Criteria

**Inclusion criteria:** All female patients who presented with palpable breast lumps. Patients who were able to give consent.

### Exclusion Criteria

The patients with an already known malignancy were excluded. Patients who were severely sick hence couldn't undergo FNAC procedure.

**Sample collection technique, processing and reporting:** Verbal consent was taken after explanation of the procedure and its benefits to the patient. In the FNAC technique, equipment used involved glass slides, disposable gloves, fixatives (70%, 95% alcohol), swabs, 23 gauge needle and 10 ml syringes. The patient was made to lie supine on an examination couch after exposure of upper half of the body. Gloving of hands was done by the resident pathologist in order to perform a breast examination. This examination involved;- an assessment of the size of the breast, overlying skin appearance, quadrants and presence or axillary lymph nodes. Thereafter the skin of the breast over the lump was disinfected with 70% alcohol in a swab.

The lump was immobilised and stabilised between the index finger and the thumb of the left hand. A 23gauge needle attached to a 10 ml syringe was inserted into the nodule. The plunger was retracted to create a vacuum in the needle for suction. Backward and forward movements were used under constant suction with the needle moved at different depths and angles within the confines of the nodule. The process was terminated when fluid appeared in the hub of the needle. The plunger was released to prevent aspiration of the material into the syringe. The needle was removed from the nodule and syringe detached. The syringe was re-attached after withdrawing the plunger and air was used with the needle tip close to the glass slide, the sample was expressed on the slide. At least two passes were made in two different

quadrants of the breast lump. With a sterile swab, pressure was applied over the FNA site for about five minutes.

The aspirated material was smeared on a slide labelled with the patient's laboratory number and another labelled slide was placed on the smear to evenly and thinly spread it between the two slides on pulling them apart, this made two smears per pass and therefore four smears per patient. Two slides were air dried whereas the other two were fixed immediately by immersion into 95% alcohol. The air-dried smears were stained with May-Grunwald-Giemsa (MGG) staining technique and the smears fixed in 95% alcohol were stained with Papanicolaou staining technique. The attending cytotechnologist took part in the assessment of specimen adequacy.

The cytological diagnoses were categorized into five standardized categories in accordance to International academy of cytology reporting criterion for breast cytology; Category I (Insufficient material), Category II (Benign), Category III (Atypical), Category IV (Suspicious, probably in situ or invasive) & Category V (Malignant) [10]. The breast samples from FNAC were further sub-classified into definite breast entities.

### Data Entry and Analysis

The data collected was entered using Epidata software and exported to the Statistical Package for Social Sciences (SPSS) version 17.0 for analysis.

### Results

Out of 291 female patients presenting with palpable breast lesions at the MakCHS pathology laboratory, the category-wise distribution was:-14 (4.8%) insufficient (C1), 192 (66%) benign (C2), 9 (3.0%) atypical (C3), 15 (5.2%) suspicious of malignancy (C4) & 61 (21%) malignant (C5). Thus the ratio of benign to malignant lesions was 3:1. In the benign category, fibroadenoma was the most commonly diagnosed lesion constituting 110/192 (57.3%) followed by fibrocystic changes constituting of 22/192 (11.5%). In the malignant category, invasive ductal carcinoma was the most commonly diagnosed lesion constituting 27/61 (44.3%) followed by Invasive ductal carcinoma of No Specific type (NST) constituting 14/61 (23.0%). The peak age group for benign lesions was 21-30 years whereas the peak for malignant lesions was 41-50 years. These findings were summarized in (Tables 1-3) respectively.

IAC Classification	Category	Frequency
Insufficient	C1	14 (4.8%)
Benign	C2	192 (66%)
Atypical	C3	9 (3.0%)
Suspicious of Malignancy	C4	15 (5.2%)
Malignant	C5	61 (21%)
<b>Total</b>		291

**Table 1:** Category-wise distribution of cases in accordance to International Academy of Cytology.

Cytological result	Frequency (%)
<b>Insufficient C1</b>	<b>n=14</b>
<i>No epithelial cells or histiocytes</i>	14(100.0)
<b>Benign C2</b>	<b>n=192</b>
<i>Acute mastitis</i>	7(3.6)
<i>Chronic mastitis</i>	8(4.2)
<i>Cystic degeneration</i>	8(4.2)
<i>Duct ectasia</i>	4(2.1)
<i>Fibroadenoma (FA)</i>	110(57.3)
<i>Fibroadenoma with chronic mastitis</i>	1(0.5)
<i>Fibrocystic changes (FCC)</i>	22(11.5)
<i>Galactocele</i>	9(4.7)

<i>Lactating adenoma</i>	1(0.5)
<i>Lipoma</i>	2(1.0)
<i>Normal breast aspirate (Negative for Malignancy)</i>	18(9.4)
<i>Tuberculous mastitis</i>	2(1.0)
<b>Atypical C3</b>	<b>n=9</b>
<i>Atypical ductal cells of undetermined significance (AUS)</i>	1(11.1)
<i>Atypical ductal hyperplasia (ADH)</i>	8(88.9)
<b>Suspicious of Malignancy C4</b>	<b>n=15</b>
<i>Low- grade ductal carcinoma in situ (Nuclear grade 1)</i>	1(6.7)
<i>High-grade ductal carcinoma in situ (Nuclear grade 3)</i>	14(93.3)
<b>Malignant C5</b>	<b>n=61</b>
<i>Adenocarcinoma of Breast</i>	10(16.4)
<i>Invasive ductal carcinoma (IDC)</i>	27(44.3)
<i>Invasive ductal carcinoma of No Specific type (NST)</i>	14(23.0)
<i>Invasive ductal papilloma</i>	9(14.8)
<i>Metaplastic carcinoma</i>	1(1.6)

**Table 2:** Type-specific distribution of cases as diagnosed by FNAC.

Age-group	Insufficient(C1)	Benign(C2)	Atypical(C3)	Suspicious(C4)	Malignant(C5)	Total
20-Nov	6	25	-	-	-	31
21-30	5	70	-	-	-	75
31-40	3	52	1	-	5	61
41-50	-	20	2	-	33	55
51-60	-	15	2	3	10	30
61-70	-	8	1	6	6	21
71-80	-	2	3	6	7	18
<b>Total</b>	<b>14</b>	<b>192</b>	<b>9</b>	<b>15</b>	<b>61</b>	<b>291</b>


**Table 3:** Age-wise distribution of cases according to breast categories.

## Discussion


Breast lumps are a common clinical presentation in women presenting with breast symptoms. Whether benign or malignant lumps cause anxiety and depression to the patient [11]. Nature of breast lump can be established by a triple test which includes clinical breast examination, imaging and fine-needle aspiration cytology [12]. FNAC is extremely recognized as a reliable procedure for the initial examination of palpable breast masses. It is minimally invasive, cost-effective, safe, simple, rapid and sensitive as compared to biopsy thus reduces the number of open breast biopsies [13].

In our study, 4.8% cases had inadequate aspirates and were placed in C1 category which was in concordance with studies done by Sudrat N, et al. [14] whereas Bajwa and Tariq [15] had a slightly higher rate (13.6%) of C1 cases. An “inadequate” report was issued as due to scanty or a cellular material with in the aspirates, Benign lesions under the C2 category were most commonly diagnosed with a percentage of 66% (192 cases) was congruent to the study done by Sunita, et al. [16] and by Ahmed, et al. [17]. Fibroadenoma (Figures 1A & 1B) and fibrocystic changes (Figures 2A & 2B) were the most common benign lesion observed in this study similar to that found by Smita and Suyrakant [18]. In this study, atypical lesions in the category C3 comprised of


3% (9 cases) was in agreement to a study done by Sudarat, et al. [19]. Smears in this category showed ductal epithelial cells benign in nature with overcrowding and mild nuclear enlargement. In the C3 category atypical ductal cells of undetermined significance (AUS) was the most diagnosed lesions followed by atypical ductal hyperplasia (ADH). This study showed 5.2% (15 cases) in suspicious for malignancy category (C4). Other studies showed variable results in C4 category, namely Rakhshinda, et al. [20] in their study found 10.3% in 464 cases and Yusuf, et al. [21] found 13.5% in 66 cases. Smears in this category showed cells. In the category C5, malignant lesions accounted for 21% (61 cases) which was in concordance to the study done by Yusuf, et al. [21]. Invasive ductal carcinoma (Figure 3) was the most commonly reported lesion which was also found by Bukhari, et al. [22]. There was a predominance of benign lesions in an age group of 21-40 years and increased malignancy in the age group of 41-50 years. Similar findings were found by Shirish, et al. [23] and Almobarak, et al. [24].




**Figure 1A:** Fibroadenoma showing papillary-like pattern (Papanicolaou stain,  $\times 50$ ).




**Figure 1B:** Fibroadenoma showing cohesive groups of ductal epithelial cells Admixed with myoepithelial cells (MGG,  $\times 200$ ).



**Figure 2A:** Fibrocystic change showing apocrine metaplasia (MGG,  $\times 400$ ).



**Figure 2B:** Fibrocystic changes showing apocrine metaplastic cells in a Honey-comb architecture (Papanicolaou stain,  $\times 200$ ).



**Figure 3:** Invasive ductal carcinoma of breast showing coarse chromatin pattern and intranuclear inclusions (MGG,  $\times 200$ ).

### Limitations of the Study

The occurrence of unsatisfactory (inadequate) smears in 14 cases was a limitation to this study. This was mainly due to the collagenous nature of the lesions and some lesions were less than 1cm in size which caused difficulties in sampling. In future studies, this may be improved through performing FNAC under ultrasound guidance.

### Conclusion and Recommendations

This study demonstrated a predominance of benign lesions in young age (reproductive-age) and an increased malignancy in older age (post-menopausal). Fine-needle aspiration cytology (FNAC) was also found to be an effective diagnostic tool in the categorization of palpable breast lumps into benign, malignant, atypical, suspicious and inadequate categories.

### References

- Shah T, Shrestha M (2008) Prevalence of Breast Lump and Risk Factors of Breast Cancer among Reproductive Aged Women of Jabalpur VDC of Sunsari District, Nepal. *Journal of Nepal Health Research Council* 2(1).
- Olayide A, Samuel O, Ganiyu R, Julius O, Oladapo K, et al. (2019) Impact of Primary Care Delay on Progression of Breast Cancer in a Black African Population: A Multicentered Survey. *Journal of Cancer Epidemiology* 2019.
- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre A, et al. (2018) Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68(6): 394-424.
- International Agency Research on Cancer (2018) Global cancer observatory: cancer today.
- Galukande M, Wabinga H, Mirembe F (2015) Breast cancer survival experiences at a tertiary hospital in sub-Saharan Africa: a cohort study. *World J Surg Oncol* 13: 220.
- Ankit S, Abhinav J, Kalayansunda S (2019) Epidemiological Study of Breast Lump Cases. *International Journal of Contemporary Surgery* 7(1).
- Das DK (2003) Fine-needle aspiration cytology: its origin, department, and present status with special reference to a developing country, India. *Diagn Cytopathol* 28(6): 345-351.
- Mohamad A, Mohsen V, Mitra R (2013) Sample size calculation in medical studies. *Gastroenterol Hepatol Bed Bench* 6(1): 14-17.
- David E, John A, Olufemi O (2017) Fine needle aspiration cytology of breast lumps with histopathologic correlation in Owo, Ondo State, Nigeria: a five-year review. *Ghana Med J* 51(1): 1-5.
- Field A, Raymond W, Rickard M, Arnold L, Brachtel E, et al. (2019) The International Academy of Cytology Yokohama System for Reporting Breast Fine-Needle Aspiration Biopsy Cytopathology. *Acta Cytol* 63(4): 257-273.
- Keyzer Dekker CMG, van Esch L, de Vries J, Ernst MF, Nieuwenhuijzen GAP, et al. (2012) An abnormal screening mammogram causes more anxiety than a palpable lump in benign breast disease. *Breast Cancer Res Treat* 134(1): 253-258.
- Ngotho J, Githaiga J, Kaisha W (2013) Palpable discrete breast masses in young women: two of the components of the modified triple test may be adequate. *S Afr J Surg* 51(2): 58-60.
- Nguansangiam S, Jesdapatarakul S, Tangjitgamol S (2009) Accuracy of fine needle aspiration cytology from breast masses in Thailand. *Asian Pacific Journal of Cancer Prevention* 10(4): 623-626.
- Sudarath N, Somneuk J, Siriwan (2009) Accuracy of fine needle aspiration cytology from breast masses in Thailand. *Asian Pac J Cancer Prev* 10(4): 623-626.
- Bajwa R, Tariq Z (2010) Association of fine needle aspiration cytology with tumor size in palpable breast lesions. *Biomedica* 26: 124-129.
- Sunita H, Urmila, Sharma DC (2015) Cytomorphological study of breast lesions with sonomammographic correlation. *J Evol Med Dent Sci* 4(81): 14137-14142.
- Ahmed HG, Ali AS, Almobarak A (2009) Utility of fine-needle aspiration as a diagnostic technique in breast lumps. *Diagn Cytopathol* 37(12): 881-884.
- Sankaye SB, Dongre SD (2014) Cytological study of palpable lumps presenting in an Indian rural setup. *Indian J Med Paediatr Oncol* 35(2): 159-164.
- Sudarath N, Somneuk J, Siriwan T (2009) Accuracy of Fine Needle Aspiration Cytology from Breast Masses in Thailand. *Asian Pacific Journal of Cancer Prevention* 10(4): 623-626.
- Bajwa R, Zulfiqar T (2010) Association of Fine Needle Aspiration Cytology with Tumor Size in Palpable breast

Lesions. *Biomedica* 26: 124-129.

21. Ibrahim Y, Atanda AT (2014) Validity of fine needle aspiration cytology of the palpable breast lesions: A teaching hospital experience. *Nigerian Journal of Basic and Clinical Sciences* 11(1): 36-40.
22. Bukhari MH, Arshad M, Jamal S, Niazi S, Bashir S, et al. (2011) Use of fine-needle aspiration in the evaluation of breast lumps. *Pathology Res Int* 2011: 689521.
23. Shirish SC, Kanika G, Arpana AD, Sukanya P, Archana CB (2014) Pattern of palpable breast lesions on fine needle aspiration: A retrospective analysis of 902 cases. *J Midlife Health* 5(4): 186-191
24. Almobarak AO, Elhassan TM, Elhoweris MH, Awadalla HM, Elmadhoun WM (2014) Cytomorphologic patterns of breast lesions in Sudanese patients: Lessons learned from fine needle aspiration cytology. *Asian Pacific Journal of Cancer Prevention* 15: 3411-3413.

