

## Pattern of Chronic Arterial Occlusive Diseases in Bangladesh and its Treatment

**Rahman MM\***

Department of Vascular Surgery, Bangabandhu Sheikh Mujib Medical University,  
Bangladesh

**\*Corresponding author:** Rahman MM, Department of Vascular Surgery,  
Bangabandhu Sheikh Mujib Medical University, Bangladesh, E-mail: profmahub@yahoo.com

### Short Communication

Volume 1 Issue 1

**Received Date:** March 01, 2017

**Published Date:** April 13, 2017

### Introduction

Chronic arterial occlusive diseases are not very much uncommon in Bangladesh and also are south asean population [1,2]. Many patients loss their limbs and become burden for his family and society and in the long run the patient succumbs to death [1,2]. So it is very much important to do accurate and timely diagnosis of the disease and offer appropriate treatment of this disease [3,4]. There are 3 (three) distinct variety of disease s which cause inhibition of arterial flow in the affected limbs thereby causing necrosis, gangrene and infection causing ever septicemia and death [1,4]. These three diseases are Atherosclerotic occlusive diseases, Obliterative endarteritis and Thrombo-angitis obliterans [4-6]. Although last one is not very common in our country. This was our aim to find out the cause of chronic arterial occlusive diseases and to formulate the type of treatment for the restoration of circulation in the affected limb [7]. For this reason, with the aim is to find out the cause of arterial occlusive disease in this group of patients by following methods:-

- 1) Ultrasonic duplex study of arterial system [8]
- 2) Serum lipid profile [9,10]
- 3) Selective arteriogram of the affected limb [6,8] and finally histopathological examination of the dissected segment of arterial lumen.

Mainly upon the basis of findings of above mentioned investigations, which included histopathological examination, the treatment options is to be redefined. Because, according to my opinion, there is great difference of operative techniques in the most common

type so arterial occlusive diseases seen in Bangladesh [4,8,10]. In case of obliterative endarteritis only Endarterectomy have good prognosis [4,8] while, in case of atherosclerotic occlusive disease angioplasty with stenting or placemat of arterial graft gives better result [10-21]. It is observed that, good long time patency /placement of arterial grafts in atherosclerotic occlusive diseases of male population in more than 50 years of age.

### References

1. Rahman M, Kadirov SI, Ismailova DA (1987) Long term intra-arterial catheter therapy in the treatment of chronic severe ischemia of lower limb. Medical Journal of Uzbekistan 45-48.
2. Karimov SI, Osipov NG, Rahman M (1988) Femoro-Distal by-pass grafting in patients with gangrene of an extremely. The Journal of cardiovascular surgery 29: 41.
3. Long term intra-arterial catheter therapy in the complex treatment of chronic severe lower limb ischemia "1988. Thesis for obtaining Ph.D in ex USSR.
4. Criqui MH, Vargas V, Denenberg JO, Ho E, Allison M, et al. ( 2005) Ethnicity and peripheral arterial disease: the soon Diego population study. Circulation 112: 2703-2707.
5. Harisson GK, Karin LA, Robertson, Nauc ler SC (2006) Inflammation and atherosclerosis. Annu Rev Pathol 1: 297-329.

6. Hiatt WR, Hoag S, Hamman RF (1995) Effect of diagnostic criteria on the prevalence of peripheral arterial disease: the San Luis Valley Diabetes Study. *Circulation* 91: 1472-1479.
7. Hobson RW, Mancel Dekkar (1995) *Vascular Surgery: principles and practice*, New York, 2004. *Circulation* 91: 1472.
8. Meyers MR, Gokce N (2007) Endothelial dysfunction in obesity: etiological role in atherosclerosis. *Curr Opin Endocrinol Diabetes obes* 14(5): 365-369.
9. Barter P (2005) the role of HDL-cholesterol in preventing atherosclerotic disease. *Euro Heart J suppl* 7: 4-8.
10. Selvin E, Erligen TP (2004) Prevalence of and risk factors for peripheral arterial disease in the limited states: results from the National Health Nutrition Examination Survey, 1999-2000. *Circulation* 110(6): 738-741.
11. Dennis FB, Thomas MB, Edward VK, Gary RS, Jonathan BT (1991) Replacement of vascular prostheses infected by bacterial biofilms. *J Vase Surg* 13(5): 575-583.
12. Kieffer E, Gomes D, Chiche L, Fléron MH, Koskas F, et al. (2004) Replacement for infrarenal aortic graft infection: early and late results in 179 patients. *J Vase Surg* 39(5): 1009-1017.
13. Daenens K, Fourneau I, Nevelsteen A (2002) Ten-year experience in autogenous reconstruction with the femoral vein in the treatment of aortofemoral prosthetic infection. *Eur J Vac Surg* 25(3): 240-245.
14. Young RM, , Cherry KJ, Davis PM, Gloviczki P, Bower TC, et al. (1999) The results of in situ prosthetic replacement for infected aortic graft. *Am J surg* 178(2): 136-140.
15. Vogt PR, Brunner La Rocca HP, Lachat M, Ruef C, Turina MI (2002) Technical details with the use of cryopreserved arterial allografts for aortic infection: influence on early and midterm mortality. *J Vac Surg* 35(1): 80-86.
16. Pruitt A, Dodson TF, Najibi S, Thourani V, Sherman A et al. (2002) Distal septic emboli and fatal brachiocephalic artery mycotic pseudoaneurysm as a complication of stenting. *J Vase Surg* 36(3): 625-658.
17. Laser A, Baker N, Rectenwald J, Eliason JL, Criado-Pallares E, et al. (2011) Graft infection after endovascular abdominal aortic aneurysm repair. *J Vase Surg* 54(1): 58-63.
18. Heyer KS, Modi P, Morasch, Matsumura JS, Kibbe MR, et al. (2009) Secondary infection of thoracic and abdominal aortic endografts. *J Vase Interv Radiol* 20(2): 173-179.
19. Alankar S1, Barth MH, Shin DD, Hong JR, Rosenberg WR (2003) Aortoduodenal fistula and associated rupture of abdominal aortic aneurysm after endoluminal stent graft repair. *J Vase Surg* 37(2): 465-468.
20. Eliason JL, Guzman RJ, Passman MA, Naslund TC (2002) Infected endovascular graft secondary to coil embolization of endoleak; a demonstration of the importance of operative sterility. *Ann Vase Surg* 16(5): 562-565.
21. Khaladj , Pichlmaier U, Stachmann A, Peterss S, Reichelt A, et al. (2013) Cryopreserved human allografts/homografts for the management of graft infections in the ascending aortic position extending to the arch. *Eur J Cardiothorac Surg* 43(6): 1170-1175.