



Comparative Study of Length-Weight-Relationship (LWR) of the Fishes and Fishing Gears use in Goronyo Reservoir and River Rima in Sokoto State, Nigeria

Gusau IA^{1*}, Muhammad U² and Bilbis BD²

¹Department of Fisheries and Aquaculture, Faculty of Agriculture, Usmanu Danfodiyo University, Sokoto, Nigeria

²National Biotechnology Development Agency (NABDA), Airport Road Lugbe, Abuja, Nigeria

*Corresponding author: Gusau IA, Department of Fisheries and Aquaculture, Faculty of Agriculture, Usmanu Danfodiyo University, Sokoto, Nigeria, Email: ibrahim.abdullahi@udusok.edu.ng

Research Article

Volume 4 Issue 2

Received Date: February 13, 2021

Published Date: March 17, 2021

DOI: [10.23880/izab-16000283](https://doi.org/10.23880/izab-16000283)

Abstract

Length - weight relationship is an important fishery management tool in estimating the average weight at a given length group and in assessing the relative wellbeing of a fish population. This study was conducted for six months, from April to October 2018 in both Goronyo reservoir (GR) and downstream River Rima (RR) to compare and evaluate the Length-Weight-Relationship (LWR) of some important commercial fishes and fishing gears used in the two water bodies. Five (5) and three (3) fish landing sites were purposively selected from Goronyo reservoir and River Rima because of their high fish landings and 10 fishermen were randomly selected and their catches monitored. The length and weight values were transformed into natural logs through the regression equation $\text{Log } W = a + b \text{ Log } L$ using SPSS computer software version 22. A total of 66 fish species were identified from the two water bodies, Goronyo reservoir accounted for 28 while River Rima accounted for 38 and 22 species were common in both water bodies. The mean b values of the LWR indicated that *Latesniloticus*, *Bagrusbayad macropterus* and *Auchenoglanisocci dentalis* exhibited positive allometric growth pattern in both water bodies, but slightly higher in Rima River (b = 3.9, 3.51, 3.7 in Rive Rima and 3.8, 3.04, 3.5 in Goronyo Reservoir) respectively. The least mean b values (b = 2.01) of LWR was recorded in respect of *Synodontisgobroni* in River Rima while *Labeocoubie* exhibited the highest b value of 3.93 in River Rima with SE value of 0.392. Gill nets were the most frequent gear used in Goronyo reservoir, and the least (3.14%) was cast net while hook and lines were almost equally used in both water bodies. In conclusion, the positive allometric growth pattern exhibited by *Latesniloticus*, *Bagrusbayad* and *Auchenoglanisocci dentalis* indicated favorable aquatic environments for the species to thrive well, therefore management practices should be centered on maintaining these aquatic environments. *Labeocoubie* should be re-introduced in both water bodies because they exhibited positive allometric growth pattern despite expected predation from *Latesniloticus* and fishing mortality. More attention should be given to the types of gears used in the water bodies in order to avoid catching smaller fishes. Selective gears like gillnet must conform to the Sokoto State Fisheries Edict of 1988 in order to protect heavily fished species such as *Synodontis* species from being over exploited. Finally, more studies are recommended to determine the effect of sex on LWR of some fish species in the two water bodies.

Keywords: Trophy Cascades; Fisheries Edict; Goronyo Reservoir; River Rima; Fishing Gears

Introduction

Stock assessment is a fundamental and important tool needed for generating inputs necessary in developing strategies for efficient and proper management, development and enhancement of the fish species in any water body [1,2]. The length-weight-relationship (LWR), length-length-relationship (LLR), weight-weight-relationship (WWR) and Fulton's condition factor ('k') are some examples of stock assessment tools use in fisheries management. These tools can be used to predict the health status, standing crop or biomass, growth pattern and yield of fish stock. Length-weight data of population are basic parameters for any monitoring study of fishes since it provides important information concerning the structure and function of populations [3]. In fish population studies, it is therefore imperative to study fish species diversity together with the fish stock assessments in order to have a clear understanding of the management strategies peculiar to the fish population in such a water body.

According to Ita, et al. [4], fishery management is water body specific. Fish growth is an addition in length and size as it ages with a growth pattern of either isometric or allometric [5]. The isometric growth pattern was when a fish shape did not change when growing up, while in an allometric pattern of growth, the fish either becomes slender as it grows or becomes deeper and bulkier.

Various scholarly works Benidittoceillio, et al. [6-12] were published which reported and described different means values for LWR, LLR, CF et cetera, for many important freshwater fish species. There was a paucity of information regarding fish LWR and fishing gears used in the Goronyo reservoir and downstream River Rima [13]. Thus, this study was aimed at bridging the gap and therefore provides information on LWR of some important fish species and the fishing gears used by the artisanal fishermen in the two water bodies.

Materials and Methods

Description of the Study Sites/Locations

The study sites were downstream River Rima (RR) and Goronyo reservoir (GR) all in Sokoto State, Northwestern Nigeria (Figure 1). The state lies between longitude 4o 8'E and 6o 5'E and Latitude 12o N and 13o 58' N [14]. The exact point on River Rima where this study was conducted was at latitude 13.0667° and Longitude 5.1667° E (Figure 2). The upper Rima is seasonal before it was dammed at Goronyo village (Figure 3) so it flows during the rainy season overflowing its banks in August and September. The reservoir (Figures 1-3) formed after damming the Rima river is known as Goronyo

reservoir; it lies on coordinates 13° 31'50"N 05° 52'56"E [15]. The reservoir has a storage capacity of 976,000,000 cubic meters [16]. It has river Bunsuru, Maradi and Gangare as its main tributaries. Some of the fish landing sites selected were in Keita, Katsira, Killaro, Marima and Gidanaryfara fishing villages surrounding it.



Figure 1: Map of Nigeria Showing Sokoto State.



Figure 2: Sattelite Map of River Rima in Sokoto along University road.



Figure 3: Goronyo Reservoir.

Data Collection and Sample Size

A total of eight (8) landing sites were purposively selected because of their high fish landings from the study area. Three (3) landing sites selected from River Rima were Usmanu Danfodiyo University major bridge (Kwalkwalawa), Dundaye village and Sokoto State Water treatment Plant along Illela road and designated as stations A, B, and C. Five (5) fish landing sites, Keta, Rimawa, Killaro, Katsira and Gidanyarfara at Goronyo reservoir were selected. Ten (10) fishermen were randomly selected and their catches monitored. A total of about 1,535 (GR, 848 and RR, 687) fish specimen samples were collected for data collection. The specimen samples were collected monthly between April and October, 2018 Appendix 1.

The Fish Length/ Size

The fishermen's fish catches were measured based on species. The length of the individual fish specimen was measured using a meter rule to the nearest mm expressed as total and standard lengths as described by Bagenal, et al. [17] and the same method carried out by Gray, et al. [18], Balogun, et al. [19] Shinkafi, et al. [20].

The Fish Weight

The individual fish of each species was measured using a weighing balance to the nearest gram (g) as described by Bagenal, et al. [17] and the same method carried out by Gray, et al. [18].

Fishing Gears and Equipment Survey

Information on types and kinds of fishing gears and equipment used by the fishermen from both water bodies was collected using a structured questionnaire. A total of

50 fishermen were selected randomly from both water bodies for an interview, using an open and closed structured questionnaire and display of their gears for inventory taking [21].

Data Analysis

Length-Weight-Relationship (LWR)

The LWR parameter of the fish species from the two water bodies was fitted into the following regression equation using SPSS computer software:

$$W = a L^b$$

The regression equation was further transformed as follows:

$$\ln W = a + b \ln L,$$

where

W = weight of the fish specimen

L = length of the fish specimen

a = intercept of the slope

b = slope of the regression

ln = natural log

Results

Growth and Growth Pattern

Table 1 showed results for means b values of the LWRs which indicated that *Latesniloticus*, *Bagrusbayad macropterus* and *Auchenoglanisocci dentalis* species from both water bodies had exhibited positive allometric (b = 3.8, 3.04 and 3.5) with those of downstream River Rima exhibiting slightly higher means values (b = 3.9, 3.51 and 3.7), respectively. The least mean b values of the LWRs were recorded in respect of specie *Synodontisgobroni* from Goronyo reservoir. *Mormyrus rume* and *Bagrusbayad macropterus* had the highest number of individuals accounting for 134 and 110 respectively while the least (4) individuals were recorded for *Labeocoubie*.

S/No.	Fish species	No. of individual		R		R2		A		b		SEE	
		GR	RR	GR	RR	GR	RR	GR	RR	GR	RR	GR	RR
1	Latesniloticus	31	4	0.841	1	0.713	1	-12.02	-21	3.8	3.9	0.9	1.138
2	Bagrusdocmac	47	16	0.81	0.707	0.66	0.499	-7	-3	3.04	3.51	0.7	0.65
3	Bagrusbayadmacropterus	110	44	0.805	0.828	0.649	0.685	-53	-11	3.2	3.82	0.98	1.3
4	Bagrus filamentous	4	Nil	0.539	-----	0.288	-----	-14.6	-----	2.4	-----	1.5	-----
5	Auchenoglanisoccidentalis	4	9	0.908	0.669	0.824	0.447	-28.5	-30	3.5	3.7	0.69	0.841
6	Auchenoglanisbiscutatus	5	7	0.868	0.959	0.753	0.92	-23.1	-51	2.09	2.3	0.7	0.844
7	Claroteslaticep	5	21	0.006	0.542	0	0.295	-12.1	3.83	2.01	2.8	10.7	0.66
8	Clarotesauratus	Nil	7		0.649	-----	0.418	-----	8	-----	2.3	-----	2.22
9	Synodontisclarias	66	58	0.827	0.645	0.693	0.416	-9.6	-1.5	3.77	3.8	1.8	1.318
10	Synodontisorex	4	5	0.999	0.313	0.998	0.098	-7.1	-5.32	2.35	2.5	1.45	0.701

11	Synodontisnigrata	12	16	0.343	0.118	0.295	0.14	-2.3	-2.046	2.51	2.45	1.92	1.53
12	Synodontismembraneus	5	Nil	0.992	-----	0.983	-----	-1.8	----	2.75	----	0.77	-----
13	Synodontisgobroni	5	Nil	0	----	0	-----	10.5	-----	3	-----	1.1	-----
14	Synodontiscourteti	9	4	0.115	1	0.013	1	-7	-6.47	2.35	3.09	0.9	1.091
15	Synodontisvermiculatus	Nil	6	-----	0.444	-----	0.19	----	-6.56	-----	2.1	-----	0.219
16	Synodontisbudgetti	Nil	5	----	0.98	----	0.98	----	2.143	----	3.8	----	0.53
17	Synodontis filamentous	Nil	5	----	0.992	----	0.983	----	-7.74	----	3.2	-----	0.711
18	Synodontismacropterus	Nil	5	-----	0.969	-----	0.939	-----	-8.23	----	3.6	-----	0.866
19	Schilbemystus	45	25	0.526	0.403	0.277	0.162	4.97	-1.66	3.65	3.1	0.74	0.87
20	Schilbeintermedius	Nil	24	----	0.212	----	0.045	----	10.613	----	2.4	----	0.951
21	Parailiapelucida	6	9	0.514	0.938	0.265	0.88	-3.67	6.55	2.53	2.09	0.886	1.078
22	Clariasgaripepinus	41	25	0.96	0.868	0.921	0.74	-598	-3.13	3.24	3.4	0.66	0.414
23	Clariasanguillaris	22	11	0.862	0.792	0.743	0.627	-60.6	-20.4	2.15	2.6	1.74	1.9
24	Malapterirus electricus	31	21	0.71	0.17	0.504	0.029	-12	51.319	3-2.53	2.3	1.29	1.16
25	Alestes nurse	Nil	11	-----	0.793	-----	0.593	-----	-3.05	-----	3.4	-----	0.65
26	Alestesbaremose	Nil	15	----	0.616	----	0.379	----	-1.04	----	3.5	-----	1.37
27	Hydrocynus forskalii	Nil	5	----	0.983	----	0.966	-----	-2.3	---	3.3	---	0.99
28	Mormyrus rume	134	102	0.735	0.833	0.54	0.695	-21	-8.6	3.48	3.7	2.31	0.99
29	Hyperopisus bebeo occidentalis	52	39	0.901	0.854	0.812	0.73	8.1	-2.4	2.42	3.5	0.85	1.069
30	Gnathonemus abadii	27	15	0.939	0.772	0.882	0.596	-3	-10.5	2.49	2.8	0.49	0.9
31	Marcusenussenegalensis	4	27	0.999		0.998	-----	-3	-----	2.1	-----	2.23	-----
32	Petrocephalus bovie	4	20	0.91	0.164	0.829	0.027	-12.9	14.56	2.9	3.1	3.995	1.38
33	Mormyrops macrophthalmus	7		0.557		0.311		30.35		2.61		1.42	
34	Mormyropstapirus	Nil	8	-----	0.982	-----	0.964	-----	-16.5	-----	2.3	-----	1.45
35	Tilapia zilli	5		0.984		0.968		-6		3.53		1.29	
36	Oreochromis niloticus	65	33	0.757	0.609	0.573	0.371	-7.5	-5.7	3.8	3.01	1.3	0.73
37	Hemichromis fasciatus	Nil	5	-----	0.998	-----	0.997	-----	-12.1	-----	2.6	-----	1.462
38	Labeosenegalensis	Nil	8	-----	0.462	-----	0.214	-----	-34.84	-----	3.1	-----	1.121
39	Labeocoubie	Nil	4	-----	0.999	-----	0.999	-----	-5.77	-----	3.93	-----	0.392
40	Polypterussenegalensis	21	Nil	0.355	----	0.128	----	-12	----	3.1	-----	0.23	-----
41	Polypterus ansorgie	Nil	12	-----	0.504	-----	0.254	-----	33.5	----	2.6	-----	0.84
42	Protopterus annectens	77	29	0.932	0.759	0.868	0.576	22.24	27.87	3.1	2.9	0.77	0.82
43	Distichodus rostratus	Nil	13	-----	0.754	-----	0.568	----	-4.5	-----	2.8	-----	1.1
44	Distichodus engycephalus	Nil	14	-----	0.376	-----	0.141	-----	-35.73	-----	2.7	-----	2.45

Table 1: The Mean b Values of the length-weight-relationships (LWRs) of some important fish species in Goronyo reservoir and downstream river Rima.

Source: The Study field work, 2018

Key: **R** = R value, **R²** = R-square value, **a** = intercept, **b** = slope of the relation and **SEE** = standard error of estimate, **RR**= River Rima, **GR** = Goronyo Reservoir

Fishing Gears and Equipment Survey

Results of the present study had indicated that gillnets and hook and line were the major gears used by the fishermen in the study area (Table 2). Gillnets account for 30.37% and

14.15% while hook and line recorded 29.28% and 29.87% in both Goronyo and river Rima respectively. The least gear (Yawa/bottom gillnet and screen fishing method) accounted for about 0.16 each in terms of usage and quantity in river Rima.

S/No.	Gear Type	Goronyo Reservoir (%)	Rima River (%)
1	Malian trap	17.35	26.73
2	Gillnet	30.37	14.15
3	Clap net	10.85	23.58
4	Cast net	Nil	3.14
5	Ndrutu	10.85	Nil
6	Fish barrier/Screen	Nil	0.16
7	Seine net	0.43	0.31
8	Hook and line	29.28	29.87
9	Pole and line	0.87	1.89
10	Yawa (bottom gillnet)	Nil	0.16
Total		100	100
	Equipment		
1	Canoe/boat	1.72	0.45
2	Fishing gourd	98.28	99.55
Total		100	100

Table 2: Fishing gears and equipment Used by the Fishermen in both Goronyo and downstream River Rima.

Source: The Study fieldwork, 2018

Discussion

The results showed most of the fish to be fairly exhibiting allometric growth patterns in both water bodies. For instance, the species *Latesniloticus* exhibited positive allometric growth pattern. This means as the fish were growing; they were adding weight and becoming bulkier and deeper. The same thing happened with other predators in both water bodies, such as *Bagrusdocmac*, *Bagrusbayad macropterus*, *Auchenoglanis occidentalis* and *Hydrocinus forskalii*.

The results obtained were probably due to good/favourable aquatic environments, slightly better in river Rima as reported by Malami, et al. [13] and Abubakar, et al. [22]. While some of our results are in agreement with other regional and international findings on biometrics of fish studied, others are contrary. Similar results were obtained in the studies conducted by Mansor, et al. [10] in Kerian river basin and Keru Lake all in Malaysia. Furthermore, the current study was in conformity with the positive allometric growth reported by Froese, et al. [9] among 1773 specimens of fish.

Several studies corroborate the negative allometric growth reported in our study. According to Reed, et al. [23], *Bagrus filamentosus*, *Auchenoglanis biscutatus* and *Clarotes laticeps* were relatively insectivorous or carnivores in

feeding habits. These species exhibited negative allometric growth patterns from both water bodies. This could be due to steep competition from other species of the same families who are favorably disposed in the Rima and the reservoir. Imaobong, et al. [24] also reported steep competition for food among species of same families in Qua Iboe River estuary, Akwalbom State, southeastern Nigeria. This could probably be responsible for them exhibiting negative growth pattern in the two water bodies. Correspondingly Waidi, et al. [25] reported negative allometric growth patterns ($b = 2.114$) for an African silver fish, *Chrysichthys nigrodigitatus* in Ogun State coastal estuary.

Synodontis clarias and others from Mochokidae family exhibited positive allometric growth patterns more than the *Synodontisgobroni* in the Rima which exhibited negative growth patterns. This finding was also reported by Beniditto-ccillio, et al. [6] in Itaipu reservoir, Parana, Brazil where the authors observed mean b values ranged from 2.34 to 3.35 for about 72 fish species. *Clarias gariepinus* exhibited positive allometric growth pattern in both Goronyo and River Rima more than *Clarias anguillaris* which might be related to their feeding habits and reproductive strategy [26]. *Mormyrus rume* which accounted for the highest individuals caught exhibited positive allometric growth pattern for most of the 134 specimens examined. However, *Mormyrops tapirus* exhibited negative allometric growth pattern ($b+ 2.3$) and was only found in Rima. By adapting to bottom-feeding habit, this species successfully colonized the bottom strata as was reported by Reed, et al. [23] in the northern Nigerian water bodies. The fishermen successfully caught these species using Ndrutu and other traps and bottom gill nets in the two water bodies. Similar findings were reported by Reed, et al. [23] in northern Nigeria.

Oreochromis niloticus exhibited a positive allometric growth pattern in both Goronyo and the Rima, indicating the adaptability of the species in the reservoir. But *Hemichromis fasciatus* was not found in the reservoir and had exhibited a negative growth pattern in the Rima. According to Reed, et al. [23], *Hemichromis fasciatus* is a predatory Cichlid and probably could not survive the steep competition for food with the other predators in the reservoir. Nehemiah, et al. [2] reported both positive and negative allometric growth patterns in *Tilapia zilli* and some *Oreochromis niloticus* species cultured in freshwater and seawater ponds in Tanzania. But, Dan-kishiya, et al. [11] reported mean negative allometric growth patterns for 5 tropical fish species which included *Tilapia zilli*, *Tilapia mariae* and *Oreochromis niloticus* in a tropical reservoir in Abuja. *Labeocoubie* and *Labeo senegalensis*, a very important fish species exhibited positive growth pattern in River Rima and were completely absent in the reservoir. This indicated the species to be suitable in lotic rather than lacustrine aquatic environment

of the reservoir. Ita, et al. [4] also reported the high number of this specie in river Niger prior to creating of Lake Kainji. *Distichodus rostratus* and *Distichodus biscutatus* all exhibited negative (2.8 and 2.7) allometric growth patterns in the reservoir. The reservoir had scanty macrophytes and other vegetation probably was responsible for the absence of these species. Reed, et al. [23] reported that this species lived on grasses and therefore could not thrive well in open water devoid of vegetation cover.

The agreements and disagreements with other works could be attributed to the condition of the species itself, its phenotype and its specific geographic location and hence its environment as several researchers has observed [27]. Phenotypic factors can play a role in affecting the allometric growth data. The bodyform and shape strongly affect the LWRs [28]. The variation in the obtained values might be correlated with many factors such as food availability, season and sex [29].

Survey on the fishing gears and equipment has shown that gillnets, hook and lines, and Malian traps are the most important fishing gears in both Goronyo and river Rima. Similar findings were reported by Reed, et al. [23] in Northern Nigeria and Gusau, et al. [21] in the Bakolori reservoir. Most of the fishing gears are seasonal and selective and is based on the type of fish species to be catch. The types of fishing gears (gillnet, cast net, hook and line, and Malian trap) found in this area were also reported by Adeyemi, et al. [30] as the common fishing gears in fish gear survey of Gbedikere Lake, Bassa, Kogi State, Nigeria.

Conclusion

This study revealed that most of the predatory fish species in both river Rima and Goronyo reservoir exhibited positive growth patterns, indicating the ability to adapt and colonize the different ecological niche in the water bodies. But few species with low resilience and ability to compete for food and space became slender as they were aging. The study recommended paying attention to *Labeocoubie* in the river Rima because of its ability to survive expected predation as they were growing. It's also very important for the gears to be regulated in order to protect the fishes in the two water bodies from been overfished.

Acknowledgment

The authors wish to acknowledge the support and encouragement of the management of the Sokoto-Rima Basin Development Authority for making their facilities available to us. We also wish to thank the fishermen who took part in the fishing gears and equipment survey as well those whose fishes were made available to us for stock assessment.

References

1. Gupta SR, Gupta PC (2008) General and Applied Ichthyology (Fish and Fisheries) (1st 2006), reprinted 2004, S. Chand and Company PVT, LTD, Ram nagar, New Delhi.
2. Nehemiah A, Justin D, Rumsha C (2012) Length-Weight-Relationships and Condition Factors of Tilapia Species Grown in Marine and Freshwater Ponds. Agriculture and Biological journal of North America 3(3): 117-124.
3. Anderson OR, Neumann RM (1996) Length, weight and associated structural indices, Fisheries Techniques. Bethesda A Fish Soc pp: 732.
4. Ita EO (1993) Inland Fisheries Resources of Nigeria. FAO of the United Nation Rome CIPA Occasional pp: 20.
5. Pauly D (2002) Reproductive Strategies of Fishes in a Tropical Temporary Stream of Upper Senegal Basin, Beole River in Mali. Aquatic Living Resources 15(1): 25-35.
6. Beneditto Cecillio E, Agostinho AA, Carnelos machadovelho RC (1997) Length-Weight-Relationships of Fishes Caught in the Itaipu Reservoir, Parana, Brazil. Naga Iclarm quarterly 20(3-4): 57-61.
7. Barham CP, Baxter A (2003) Condition Factor, K for Salmonid fish, Fisheries Notes pp: 1-3.
8. Fafioye OO, Oluaye OA (2005) Length-Weight-Relationships of Five Fish Species in Ife Lagoon, Nigeria. African Journal of Biotechnology 4(7): 741-751.
9. Froese R (2006) Cube Law, Condition Factor and Weight-Length- Relationship: History, metal-analysis and Recommendation. Journal of Applied Ichthyology 22(4): 241-253.
10. Mansor MI, Rawi CSM, Rosla R, Sha SAM, Sha ASRM (2010) The Length-Weight-Relationships of Fresh Water Fish Species in Kerian River Basin and Pedu Lake. Research Journal of Fisheries and Hydrobiology 5(1): 1-8.
11. Dan Kishiya A (2013) Length-Weight- Relationship and Condition Factor of Five Fish Species from a Tropical Water Supply Reservoir in Abuja, Nigeria. American Journal of Research Communication 1(9): 175-187.
12. Jin S, Yam X, Zhong H, Fan W (2015) Weight-Length-Relationships and Fulton's Condition Factors of Skip Jack Tuna (*Katsuwonus pelamis*) in the Western and Central Pacific Ocean. Pear J 3: e758.

13. Malami GZ, Magawata I (2010) Analysis of Food and Feeding Habits of Catfish (*Bagrus bayad macropterus*, Daget) in River Rima in Sokoto State, Nigeria. Nigerian Journal of Basic and Applied Sciences 18(2): 277-284.
14. Mamman AB, Oyebanji JO, Peters SW (2000) Nigeria: A People United A Future Assured (Sokoto State), Gabumo Publishing Company Limited, Lagos Nigeria pp: 298.
15. Google Earth (2016) Image Viewer.
16. Sembenelli Consulting (2010) Goronyo Main and Secondary Dam (pdf) Archived from the original.
17. Bagenal T, Tesch F (1978) Age and Growth in: T.B. Methods for Assessments of Fish Production in Freshwater 3rd (Edn.), Oxford: Blackwell Scientific Publications.
18. Gray CA, Jones M, Rotterdam D, Broadhurst M, Johnson D (2005) Utility and Efficiency and Multi mesh Gill net and Trammel nets for Sampling Assemblages Populations of Estuaries Fishes. Marine and Freshwater Research 56: 1077-1088.
19. Balogun JK (2005) Fish Distribution in a Small Reservoir: A case Study of Kaingimi Reservoir. Journal of Applied Science and Environmental Management 9(1): 93-97.
20. Shinkafi AS, Ukaola LN, Thekir NA (2013) Some Aspects of Growth and Reproduction in Nile Perch (*Latesniloticus*, Linne 1760) from Rivers Rima, North western Nigeria. Nigerian Journal of Fisheries and Agriculture 1(1): 31-41.
21. Gusau IA, Bilkisu AS, Ipinjolu JK (2016) Artisanal Fisheries of Bakolori Reservoir in Zamfara State, Northwestern Nigeria. Equity Journal of Science and Technology 3(2): 43-50.
22. Abubakar M, Ipinjolu JK, Magawata M (2012) Seasonal Variation in Some Chemical Parameters of the Sokoto-Rima Rivers In Northwestern Nigeria. Journal of Agriculture and Environment 3(2): 55-65.
23. Reed W, John B, Hopson AJ, Jennings J, Yaro I (1967) The Fish and Fisheries of Northern Nigeria. In: 1st (Edn.), Zaria, Gaskiya Corporation Limited, Zaria, Nigeria.
24. Imaobong EE, Mandu AE, Nkwoji JN (2014) Food and feeding habits and condition factor of fish species in Qua Iboe River estuary, Akwalbom State, southeastern Nigeria. International Journal of Fisheries and Aquatic Science 2(2): 38-46.
25. Waidi OA (2015) Life History Change in Silver- Catfish (*Chrysichthys nigrodigitatus*) family Bagridae, in Ogun State Coastal Estuary, Nigeria. Ife Journal of Sciences 17(3): 691-700.
26. Yusuf K, Dada SA, Abari MI (2013) Length-Weight-Relationship, Fecundity and Gonadal development of the African Catfish (*Clarias gariepinus*) from Dowa Dam, Nasarawa State, Nigeria. PAT June 9(1): 47-58.
27. Tsoumani M, Liasko R, Moutsaki P, Kagalou I, Leonardos I (2006) Length-weight relationships of an invasive cyprinid fish (*Carassius gibelio*) from 12 Greek lakes in relation to their trophic states. J Appl Ichth 22(4): 281-284.
28. Karachle PK, Stergiou KI, (2012) Morphometrics and Allometry in Fishes. In Agricultural and Biological Sciences Christina Wahl pp: 65-86.
29. Ali RAS, Elawad AN, Khalifa MM, El Mor M (2016) Length-weight relationship and condition factor of *Liza ramada* from Eastern coast of Libya. Int J Fish Aqua Res 2(2): 1-9.
30. Adeyemi S, Bankole N, Adikwu I (2009) Fish Gear Survey of Gbedikere Lake, Bassa, Kogi State, Nigeria. International Journal of Lakes and Rivers 2(1): 53-56.
31. Yilma S, Yazıcioglu O, Erbasaran M, Esen S, Zengin M, et al. (2012) Length weight relationship and relative condition factor of white bream, *Bliccabjoerkna* (L., 1758), from Lake Ladik Turkey. J Black Sea Medit Environ 18(3): 380-387.

