



First Record of the Nest Orientation of *Apis Dorsata* in Relation to Cell Tower

Pattazhy S*

Department of Zoology, Universty of Kerala, India

*Corresponding author: Sainudeen Pattazhy, Department of Zoology, Universty of Kerala, Kariavattom, Thiruvananthapuram, India, Tel: 9447243002; Email: sainudeenpattazhy@hotmail.com

Research Article

Volume 3 Issue 2

Received Date: March 31, 2020

Published Date: April 20, 2020

DOI: 10.23880/izab-16000218

Abstract

This study is aimed to study the nest orientation of *Apis dorsata* in relation to cell tower in the Kariavattom panchayat of Thiruvananthapuram district, Kerala. The distributions and orientation of nests were observed around 5 km of cell tower using a survey method. Results showed that 60 nests (40 active nests, 20 abandoned combs) of *A. dorsata* were found. Tree species used as nesting sites were *Ficus* sp.(Moraceae), *Adenanthera* sp.(Fabaceae), *Spondias pinnata* (Anacardiaceae), *Artocarpus sericoarpus* (Moraceae), *Alstonia scholaris* (Apocynaceae), *Knema cinerea* (Myristicaceae) and *Litsea mappacea* (Lauraceae). The nests in trees were found in 3-11 meters (5 nests), 11-25 meters (20 nests), and 20-35 meters on buildings (35 nests). Maximum nets on building were noted on the top of Greenfield International Stadium (30 nests). Most of the bee nests were oriented towards South east direction and minimum towards south west direction on buildings and trees. In the South east direction maximum bee hive orientation was noted both on trees and buildings where radiation exposure from cell towers was minimum (0.20 w/m^2) whereas in the South west direction minimum bee hive orientation was recorded where the radiation exposure was maximum (9.20 w/m^2).

Keywords: Nest Orientation; *A. dorsata*; Cell Tower; Building

Introduction

Apis dorsata construct nests on varieties of nesting structures that provide suitable habitat for leading successful life [1,2]. Neupane, et al. [3] found that greater preference of *A. dorsata* colonies towards strong-cemented water tanks and residential buildings for nesting. These colonies are often found in aggregations on a suitable nesting structure and their numbers vary from a few colonies to many hundreds colonies [4,5]. *A. dorsata* build nests by orienting towards suitable compass directions. They are found to orient their nests in large numbers towards North-South axis by minimizing the exposure to strong wind and bright sunlight [6].

The present study aims to study the nest orientation of *A. dorsata* in relation to cell towers. The forest may be especially

reliant on animal pollinators, such as honey bee because they are needed for long distance movement of pollen among spatially separated conspecific trees that are often self-incompatible or dioeciously [7,8]. Rapid development of telecommunication devices has caused enhancement of human interference with nature so that undesirable impacts were noticed on biological, physical and ecological systems. To combat the need of mobile connection large number of mobile towers is installed in urban and rural areas. The electromagnetic radiation emitted from the mobile towers, transmission lines and power lines induces undesirable impact on living beings in a multiple ways causing pollution known as electromagnetic smog.

Material and Methods

Observation of nest orientation of *A. dorsata* was carried

out from April 2019 to March 2020 in Kariavattom village (8.5678° N, 76.8908° E) and it is an area of Trivandrum city the capital of Kerala where Greenfield International stadium, the largest international stadium of Kerala is situated. The vegetation is of mixed type dominated by acacia plants. Other vegetation includes trees, shrubs, climbers, creepers etc. Interestingly, this region was abundant with buildings and trees as major nesting structures. Five mobile towers were installed in this village. These studies used survey and interview method [9,10]. The live and abandoned combs found were then recorded and the information about the existence of the nests in that location was also collected. The position of the trees and nesting site was measured by using GPS *Garmin map 62s*. Nesting site preference of *Apis dorsata* was described by descriptive analysis. The distribution and orientation of nests was visualized using ArcGIS 10.2 program. (<http://www.esri.com/landingpages/software/arcgis/arcgis-desktop-student-trial>). EMF (Electromotive field) power density was measured with the help of RF Power density meter and direction of area with magnetic pocket compass. The nest height from ground level (m) was measured by using hagameter. Characteristic, slope and direction of branching, as well as the protected and unprotected nest of honey bee by using binoculars and compass. The physical conditions around the nest observed were the type of habitat, water distance from the nest, relative humidity (%) and temperature (°C) by using termohygrometer, wind speed (km/h) using anemometer, and light intensity (lux) by using lux meter. Monthly average data of rainfall, humidity, and wind speed were collected from Meteorological Department [11].

Results

The comb of *Apis dorsata* is more or less semicircular. The width of the *Apis dorsata* comb was 98.5 ± 26.2 cm ($n = 158$), and the height was 45.8 ± 11.4 cm ($n = 146$). Results showed that 60 nests (40 active nests, 20 abandoned combs) of *A. dorsata* were found. Tree species used as nesting sites were *Ficus* sp. (Moraceae), *Adenantha* sp. (Fabaceae), *Spondias pinnata* (Anacardiaceae), *Artocarpus sericoarpus* (Moraceae), *Alstonia scholaris* (Apocynaceae), *Knema cinerea* (Myristicaceae) and *Litsea mappacea* (Lauraceae). The nests in trees were found in 3-11 meters (5 nests), 11-25 meters (20 nests), and 20-35 meters on buildings (35 nests). Maximum nests on building were noted on the top curved sloppy surface of Greenfield International Stadium (30 nests). Most of the bee nests were oriented towards South east direction and minimum towards south west direction on buildings and trees. In the present study, the orientation of *A. dorsata* nests on different nesting structures was recorded towards all compass directions. But they showed fluctuation in their numbers towards each direction. Generally selection of these nesting structures will be attributed to greater protection

of colonies against pests, predators, bright sunlight and even winds pressure. But in the present study it was shown that nest orientation was in accordance with intensity of Electromagnetic radiations from cell towers. Maximum nest orientation towards southeast direction was due to less radiation exposure (0.2 watt/m^2) whereas high radiation exposure (8.1 watts/m^2) was noted where nest construction was less. The abandoned bee hives were mostly found within 50–300 meters around the tower where radiation exposure was maximum (9.2 watts/m^2) (Table 1). The trees are the primary nesting structures of giant honeybees due to cool and pleasant environment for bee nests along with sufficient shade and moderate amounts of sunlight for normal colony activities [2]. But, in the present study *A. dorsata* selected buildings for maximum nest construction for availing radiation shielding from cell towers. 90 present nests was noted 500 meter to 2 Km away from cell towers could be due to less radiation exposure.

Buildings	Trees	Radiation Level (watts/m ²)	
East-West	4	3	5
North-South 3.00	3	6	
North-East	4	4	5
North-West	4	4	5.2
South-East	18	10	0.2
South-West	2	1	9.2

Table 1: Nest orientation of *Apis dorsata* colonies on different nesting structures.

Discussion

During the present study a total of 60 nests (40 active nests, 20 abandoned combs) of *A. dorsata* were found. The nests in trees were found in 3-11 meters (5 nests), 11-25 meters (20 nests), and 20-35 meters on buildings (35 nests). Maximum nests on building were noted on the curved sloppy surface of top of Greenfield International Stadium (30 nests). Most of the bee nests were oriented towards South east direction and minimum towards south west direction on buildings and trees. In the South east direction maximum bee hive orientation was noted both on trees and buildings where radiation exposure from cell towers was minimum (0.20 w/m^2) whereas in the South west direction minimum bee hive orientation was recorded where the radiation exposure was maximum (9.20 w/m^2). In the East west, North south, North east and North west directions radiation exposure varied from 5 to 6 w/m^2 where bee hive was ranged from 3 to 4 numbers. The abandoned hives were mostly around 50-300 meters around the cell towers where radiation exposure range was extreme (9.2 w/m^2).

All mobile phone towers emit microwave radiations, which is in the radiofrequency radiation (RFR), a part of the spectrum of electromagnetic waves. Though radiofrequency radiation is a source of non-ionizing radiation, these radiations together with ionizing radiation make up the electromagnetic spectrum. Radio frequency of the electromagnetic waves ranged from 100 kilo hertz (KHz) to 300 Giga hertz (GHz). Radio frequency radiation is a source

of thermal energy and in adequate doses, has all the known effects of heating on biological systems. Sharing of towers by different companies should not be encouraged otherwise the magnitude of radiation would be increased. Frequencies of different mobile tower companies are represented in Table 2. Independent monitoring of radiation levels and overall health of the community and nature surrounding towers is necessary to identify hazards early.

S. No.	Mobile Technology in India	Frequency Bands in India	Major Operators
1	GSM (2G)	900 MHz, 1800 MHz	Airtel, Idea, Vodafone, Aircel, BSNL
2	CDMA	850 MHz	Reliance, BSNL, Tata
3	WCDMA (3G)	2100 MHz, 900 MHz	Airtel, Idea, Vodafone
4	WiMAX	2300 MHz	BSNL
5	4G LTE (4G)	1800 MHz	Airtel, Idea, Vodafone, Jio
		850 MHz	Jio
		2300 MHz	Airtel, Idea, Vodafone, Jio
		2500 MHz	BSNL, Idea, Vodafone

Table 2: Table for the different frequency bands in India for mobile technology 2G, 3G and 4G.

References

- Deodikar GB, Ghatge AI, Phadke RP, Mahindre DB, Kshirsagar KK, et al. (1977) Nesting behaviour of Indian honeybees.
- Reddy CC, Reddy MS (1993) Studies on the distribution of nests of giant honeybee (*Apis dorsata* F.). Indian Bee J 55: 36-39.
- Neupane KR (2004) Nesting behaviour of giant honeybees. Proc 8th IBRA International Conference. Trop. Bees, Brazil, pp: 351-357.
- Kumar GMN, Reddy CC (2003) Nest density, dimension and distribution of *Apis dorsata* F. in relation to height and direction. Asian Bee J 5: 82-89.
- Nagaraja N, Rajagopal D (2009) Honeybees: Diseases, Parasites, Pests, Predators and their Management. MJP publishers, Chennai, India, pp: 210.
- Reddy MS (1983) Studies on the factors of selection of nesting site by *Apis dorsata* F. Ph.D. thesis, Bangalore University, Bangalore, India, pp: 146.
- Bawa KS (1990) Plant-pollinator interactions in tropical rain forest. Annual Review of Ecology Systematics 21: 399-422.
- Corlett RT (2004) Flower visitor and pollination in the oriental (Indomalayan) region. Biological Reviews 79: 497-532.
- Bookhout TA (1996) Research and Management Techniques For Wildlife And Habitats Kansas (US): Allen Press Inc.
- Tongco MDC (2007) Purposive sampling as a tool for informant selection. Ethnobotany Research and Applications 5: 147-158.
- Nesting behaviour of *Apis dorsata* Fab. Indian Bee J 39: 1-12.

