



Solitary Orbital Plasmocytoma: A Case Report and Review of Literature

Belgacem I*, Djouadi Y, Habchi N and Djafer M

Department of Neurosurgery, Mustapha Pacha University Hospital, Algeria

*Corresponding author: Belgacem I, Department of neurosurgery, Mustapha Pacha University Hospital, Algiers, Algeria, Email: nawelmedbio@yahoo.fr

Case Report

Volume 5 Issue 2

Received Date: July 17, 2020

Published Date: November 12, 2020

DOI: 10.23880/nnoaj-16000153

Abstract

Solitary orbital plasma cell neoplasm in the absence of multiple myeloma is very rare. We present here a case report of extramedullary plasmacytoma of the left orbital soft tissue in a 50-year-old woman whose thorough diagnostic workup was negative for multiple myeloma. The patient presented with rapidly increasing proptosis, swelling, and visual disturbances of the left eye for past 15 days. MRI of the orbit revealed an extra conal well-circumscribed homogeneously enhancing masse filling the superior and the temporal part of the left orbit without bone destruction Or intracranial extension. The patient underwent a subtotal resection and the pathological report revealed a plasma cell neoplasm. So, normal skeletal survey, absence of bone marrow involvement, absence of Bence-Jones protein, no anemia, no hypercalcemia, no renal disease, normal level of immunoglobulin's all exclude the diagnosis of multiple myeloma. Since the visual acuity of the patient is well-preserved, so we have planned to treat her with local radiotherapy alone with 50 Gy.

Keywords: Orbit; Plasmacytoma; Radiotherapy

Introduction

Plasmocytoma can be manifest as systemic disease, namely, multiple myeloma [1] and its variants (such as indolent myeloma, smoldering myeloma, osteosclerotic myeloma, plasma cell leukemia and non-secretory myeloma) or localized disease represented as solitary plasmacytoma of bone (SMB) or extramedullary plasmacytoma (EMP)[2,3]. Involvement of the orbit can occur in either of the two forms. Solitary orbital plasma cell neoplasm in the absence of multiple myeloma is very rare with less than 50 cases reported in the literature [1,4,5].

Case Report

A 50 year-old female; with a biliary cirrhosis history was referred from an ophthalmologist with rapidly progressive left proptosis for past 15 days and moderate visual disturbances. Physical examination showed with left-sided proptosis, swelling, limited ocular movement (an offense of

abduction) with sever impairment of vision. The visual acuity was 8/10 of the left eye. The patient orbital MRI showing, an extra conal well-circumscribed homogeneously enhancing masse filling the superior and the temporal part of the left orbit without bone destruction (Figure 1).



Figure 1 : MRI scan depicts left superotemporal orbital mass without intracranial extension.

The patient underwent subtotal resection (Figures 2 & 3) to protect and improve ocular function by preserving the surrounding anatomical structures through a sub frontal approach with removal of upper orbital rim (Figure 4), the result of the pathological report revealed a plasma cell neoplasm.



Figure 2: Exposed orbital contents.



Figure 3: Image of the lesion at the moment of being resected through a sub frontal supra orbital craniotomy.



Figure 4: Bone Flap.

Complete skeletal survey and bone marrow examination report were normal. So, normal skeletal survey, absence of bone marrow involvement, absence of Bence-Jones protein, no anemia, no hypercalcemia, no renal disease, normal level

of immunoglobulin's all exclude the diagnosis of multiple myeloma. The diagnosis of solitary orbital plasmacytoma is retained. And have planned to treat her with local radiotherapy alone with 50 Gy to stabilize the tumor volume after 04 months of evolution.

Discussion

Orbital plasmacytoma can be primary or secondary. Side is multiple myeloma events [2]; more frequent and tend to be more aggressive. The diagnosis of plasmacytoma requires a biopsy to distinguish several closely related pathologies, including plasma cell granuloma, plasmacytoid lymphoma and large cell lymphomas (immunoblastic type). The prognostic importance comes from a higher response rate to chemotherapy or radiotherapy [6-8], and a prolonged overall survival compared with those with the sole diagnosis of multiple myeloma. Radiotherapy as the sole treatment option, as given in this case is adequate in the absence of systemic involvement [6].

Plasmacytomas have been classified as low grade, intermediate or high grade using criteria originally intended for the grading of multiple myelomas. Chemotherapy was recommended after primary radiotherapy [8] for the latter two grades as these lesions had only 17% local control rate with radiation alone. However, EMPs with low-grade histologies were controlled in 83% cases external radiation only. Our case was a low-grade EMP of orbital soft tissue and well-preserved visual acuity treatment with radiation alone was justified.

Conclusion

An orbital plasma cell neoplasm with no systemic involvement at present may be the first of its kind to be reported to the best of our knowledge. However, the risk of progression to multiple myeloma is not yet documented, should always be kept in mind, mandating a close follow-up of this patient.

References

1. Matos A, Goulart A, Ribeiro A, Freitas R, Monteiro C, et al. (2020) Orbital Plasmacytoma, An Uncommon Presentation of Advanced Multiple Myeloma. *Eur J Case Rep Intern Med* 7(3): 001149.
2. Sen S, Kashyap S, Betharia SM (2003) Primary orbital plasmacytoma: a case report. *Orbit* 22(4): 317-319.
3. Pierre-Filho P de TP, do Carmo Carvalho F, Braga Perdigão F, Caldato R, Rolim da Paz A (2004) Orbital extramedullary plasmacytoma leading to exenteration. *Can J Ophthalmol J Can Ophtalmol* Août 39(5): 557-559.

4. Meyer HJ, Ullrich S, Hamerla G, Surov A (2018) Extramedullary Plasmacytoma. ROFO Fortschr Geb Rontgenstr Nuklearmed 190(11): 1006-1009.
5. Wang SSY, Lee MB, George A, Wang SB, Blackwell J, et al. (2019) Five cases of orbital extramedullary plasmacytoma: diagnosis and management of an aggressive malignancy. Orbit Amst Neth Juin 38(3):218-25.
6. Tsang RW, Campbell BA, Goda JS, Kelsey CR, Kirova YM, et al. (2018) Radiation Therapy for Solitary Plasmacytoma and Multiple Myeloma: Guidelines From the International Lymphoma Radiation Oncology Group. Int J Radiat Oncol Biol Phys 101(4): 794-808.
7. Ezra E, Mannor G, Wright JE, Rose GE (1995) Inadequately Irradiated Solitary Extramedullar Plasmacytoma of the Orbit Requiring Exenteration. Am J Ophthalmol 120(6): 803-805.
8. (2020) Plasmocytoma vs radiotherapy. Recherche Google.
9. Richard W Tsang, Belinda A Campbell, Jayant S Goda, Peter M Mauch, Lena Specht, et al. (2018) Radiation Therapy for Solitary Plasmacytoma and Multiple Myeloma: Guidelines From the International Lymphoma Radiation Oncology Group 101(4) : 794-808.

