

Food Additives and Food Acceptability

Shaltout F*

Department of food control, Faculty of Veterinary Medicine, Benha University, Egypt

***Corresponding author:** Fahim Shaltout, Food Control Department, Faculty of Veterinary Medicine, Benha University, Egypt, Email: fahim.shaltout@fvtm.bu.edu.eg

Review Article

Volume 2 Issue 1 Received Date: April 16, 2024 Published Date: June 10, 2024 DOI: 10.23880/oajfs-16000108

Abstract

The Producers and the manufacturers have been challenged by the increasing demand for the safe and high-quality meat and meat products over the past few decades. Particularly, the recent demand for minimally the processed, the easily prepared, and the ready-to-eat meat products combined with the novel concepts of all-natural and clean-label has rapidly increased. These products may contain natural or organic ingredients without artificial preservatives that do not trigger the common food allergies or the sensitivities. The meat and the meat products are highly prone to microbial contamination since they are rich in essential nutrients and perishable. This is further accelerated by some intrinsic factors including pH and water activity of the fresh meat. In general, the freshest meat has a water activity value higher than 0.85, and its pH value falls within the favorable pH range for spoilage bacteria of the meat. Hence, deterioration in quality and potential public health issues is common if these products are not properly handled and preserved. The significant spoilage of the meat and the meat products occurs every year at different levels of the production chain including the preparation, the storage, and the distribution. Besides the lipid oxidation and the autolytic enzymatic spoilage, the microbial spoilage plays a significant role in this deterioration process leading to a substantial economic and environmental impact.

Keywords: Essential Oils; Safe; High Quality; Meat; Meat Products

Introduction

The microorganisms associated with the spoilage of the meat and the meat products including bacteria such as the Pseudomonas, the Acinetobacter, the Brochothrix thermosphacta, the Lactobacillus spp., the Enterobacter, etc., and the yeast and the mold cause the quality defects such as of flavor, off-odor, etc. Additionally, the foodborne diseases have emerged as important and growing public health and economic problems in many countries over the last few decades [1-6]. The foodborne diseases are not limited to a particular age group or the country. The foodborne diseases occur each year resulting in the illnesses, with hospitalizations and deaths, leading to high medical costs and productivity losses. In regarding to the meat and the meat products, several pathogenic microorganisms including the Salmonella spp., the Campylobacter *jejuni*, the Escherichia coli O157:H7, the Listeria *monocytogenes*, the Clostridium spp. and the *Aeromonas hydrophila* can result in the foodborne illnesses to the consumers if the products are not preserved and handled properly [7-12].

The vegetative cells of the microbes are destroyed with the thermal processing, but not the spores of the food pathogens such as the Bacillus spp. and Clostridium spp. They can survive and need much harsher conditions to be inactivated.



Open Access Journal of Frailty Science

With the absence of the competitive microflora, these spores can germinate and grow under favorable conditions caused by mishandling of the heat-treated products. Hence, it is vital to apply other hurdles such as low-temperature storage to avoid this problem [13-19]. To extend the period of the refrigerated storage, which is the most common method used for preserving the fresh meat and the meat products, many synthetic additives have been used over the years. The synthetic additives have been accused for some carcinogenic and toxic properties. This increased the consumer concerns towards the healthier meat and the meat products and the demand for the natural food additives over the years, which led researchers to examine the natural alternatives to the synthetic food additives [20-26].

The meat and the meat products are highly subject to the microbial deterioration, which ultimately leads to the safety and the quality issues if the meat are not properly handled and preserved. Several plant-derived Essential oils can be effectively used in the meat and the meat products as natural alternatives to the synthetic food additives, particularly as the effective antimicrobial agents [27-33]. The Phenolic compounds, such as the carvacrol, the eugenol, and the thymol, are mainly responsible for the antimicrobial activity of the Essential oils to increase the permeability of cell membranes and leading to loss of the cellular constituents. These natural additives should improve the meat quality without leaving residues in the product or the environment [34-40].

The Microbial Deterioration of the Meat and the Meat Products

Depending on the preservation method used, the growth and the metabolism of the spoilage and the pathogenic microorganisms can cause rapid spoilage of the meat and the meat products and the serious foodborne intoxications .The Bacteria are generally considered as the principle agents responsible for these deteriorations and the health issues in addition to the molds and yeasts presents the major genera of bacteria, yeasts, and molds found in meat and meat products [41-48]. Main spoilage bacteria including the Pseudomonas, the Acinetobacter, the B. thermosphacta, the Moraxella, the Enterobacter, the Lactobacillus spp., the Leuconostoc spp., the Proteus spp. etc, the yeast and the mold, the decompose meat and the meat products and develop the unpleasant quality characteristics when they grow in large numbers in these perishable products [49-56]. This can be attributed to the degradation of the proteins and the lipids present in the meat and the meat products resulting in the off-odors, the off-flavors, the discoloration, the texture defects, the slime and the gas productions, and the changes in pH. Although the spoilage microorganisms normally do not cause the illness. they can result in the gastrointestinal disturbances when consumed in the high concentrations [57-64].

However, the rate of the meat spoilage is affected by several factors including the hygiene, the storage temperature, and the acidity of the meat and the meat products. The growth of the spoilage bacteria is favored at a pH range of 5.5-7.0. The Campylobacter jejuni, Salmonella serotype Typhimurium, Escherichia coli 0157:H7, other Enterohemorrhagic E. coli (EHEC), Listeria monocytogenes, Arcobacter butzleri, Mycobacterium avium subspp. paratuberculosis and Aeromonas hydrophila are the most prevalent and serious emerging pathogens in the meat and the meat-derived products [65-72]. The color, the odor, the taste or the texture of the meat and the meat products are not often changed by these pathogenic bacteria. Which are mainly responsible for the food poisoning and the food intoxications. The microbial deterioration of the meat and the meat products can be considered as one of the main limitations in the meat industry. The proper protective practices should be, therefore, applied for the meat and the meat products to produce the highest quality products possible [73-79].

The common preservation methods used against the microbial deterioration Due to the increasing demand for the precooked, the refrigerated, the ready-to-eat meat, and the meat products convenient for the modern and the busy lifestyles, the meat manufacturers must overcome many challenges, including the control of the microbial deterioration of these products [80-87]. Several factors such as the microbial growth, the color, and the lipid oxidation affect the shelf life and the consumer acceptance of the fresh meat and the meat products. Therefore, to produce the safest and the highest quality products possible, these factors must be controlled [88-95]. Several of the thermal and the nonthermal food preservation techniques have been used, alone or in combinations, to prevent or minimize the growth of spoilage and the pathogenic microorganisms in the meat and the meat products .The common physical and the chemical preservation techniques used to control the bacterial activity [96-102].

The Mode of Action of the Essential Oils

The antimicrobial activity of a given Essential oil can be attributed to its major constituents as well as their interaction with the minor constituents present in oils. However, the antimicrobial activity of the Essential oils has been consistently linked to phenolic constituents such as the carvacrol, the eugenol, and the thymol. The presence of the hydroxyl groups in the phenolic compounds is very vital for their antimicrobial activity [103-109].

The antimicrobial activity of the Essential oils is not attributable to one specific mechanism. Several locations or mechanisms in the microbial cells are supposed to be the sites of action for the Essential oil constituents. In brief, Essential oils can degrade the cell wall, disturb the phospholipid bilayer of the cytoplasmic membrane, and damage the membrane proteins leading to increased permeability of the cell membrane and loss of cellular constituents [110-116]. The Essential oils can further disrupt the proton motive force, electron flow, and active transport, and coagulate the cell contents .The Essential oils can impair a variety of enzyme systems including the enzymes involved in the energy regulation and synthesis of structural components and inactivate or destroy genetic material, strengthening their antimicrobial activities [117-122].

Limitations and Future Perspectives

Although good antimicrobial activities were observed for many Essential oils, some limitations have 446 also been identified in the application of the Essential oils in the meat and the meat products. The interaction of some Essential oils with the food ingredients and structure may decrease their effectiveness [123-129]. The markedly reduced activity of the Essential oils may result in the food systems such as the meat and the meat products when compared to in vitro results. This may be attributed to the presence of fats, carbohydrates, proteins, and salts in such systems. For instance, the mint and the cilantro Essential oils were not effective against the *L. monocytogenes* in the products containing high levels of the fat, such as the pâté and a coating for the ham containing the canola oil [130-136].

It may be difficult to maintain the quality consistency because the composition of an individual Essential oil can vary due to several factors including the time of harvesting, variety, the part of the plant used, and method of extraction. The antimicrobial potency of the Essential oil constituents depends on pH, temperature, and level of microbial contamination [137-143]. Further, use of the Essential oils as the preservatives in the food has been limited as they are required in high concentrations in order to achieve the sufficient antimicrobial activity. The use of some Essential oils are mainly restricted to the spicy foods including the meat products that are associated with the herbs, spices, or seasonings since they affect the organoleptic quality of the food [144-150].

Conclusion

The application of the Essential oils is partially limited due to their intense aroma which may cause negative organoleptic effects. Novel technologies such as encapsulation of the Essential oils into the Nano emulsions and the use of the Essential oils as part of the hurdle technology (combined processes with MAP, nisin, EDTA, lysozyme etc.) to improve the microbial stability and the sensory quality of the meat and the meat products are being used in the meat industry; traditional methods of adding the Essential oils directly into the meat batter during manufacturing of the meat products.

Conflicts of Interest

The author declare no conflicts of interest regarding the publication of this paper.

References

- 1. Nataro JP, Kaper JB (1998) Diarrheagenic Escherichia coli. Clinical Microbiology Reviews 11(1): 142-201.
- Shaltout FA, Riad EM, Elhassan AA (2017) Prevalence Of Mycobacterium Spp. In Cattle Meat And Offal's Slaughtered In And Out Abattoir. Egyptian Veterinary medical Association 77(2): 407-420.
- 3. McDonnell MJ, Rivas L, Burgess CM, Fanning S, Duffy G (2012) Evaluation of Carvacrol for Control of Escherichia coli 0157 on Cattle Hide and Carcass Cuts. Foodborne Pathogens and Disease 9(11): 1049-1052.
- Elaziz OA, Hassanin F, Shaltout F, Mohamed O (2021) Prevalence of Some Foodborne Parasitic Affection in Slaughtered Animals in Loacal Egyptian Abottoir. Journal of Nutrition Food Science and Technology 6(2): 37-42.
- 5. Cornforth D (1994) Color—Its Basis and Importance. In: Pearson AM, Dutson TR (Eds.), Quality Attributes and their Measurement in Meat, Poultry and Fish Products (Edn.). Springer, USA, pp: 34-78.
- 6. Elaziz OMA, Hassanin FS, Shaltout FA, Mohamed OA (2021) Prevalence of some zoonotic parasitic affections in sheep carcasses in a local abattoir in Cairo, Egypt. Advances in Nutrition & Food Science 2(3): 38.
- Fasseas MK, Mountzouris KC, Tarantilis PA, Polissiou M, Zervas G (2008) Antioxidant Activity in Meat Treated with Oregano and Sage Essential Oils. Food Chemistry 106(3): 1188-1194.
- 8. Al Shorman AAM, Shaltout FA, Hilat N (1999) Detection of certain hormone residues in meat marketed in Jordan. Jordan University of Science and Technology, 1st International Conference on Sheep and goat Diseases and Productivity, pp: 23-25.
- 9. Souza AA, Dias NAA, Piccoli RH, Bertolucci SKV (2016) Chemical composition and minimum bactericidal concentration of sixteen essential oils against enterotoxigenic Escherichia coli. Revista Brasileira de Plantas Medicinais 18(1): 105-112.

- Saleh E , Shaltout F , Elaal EA (2021) Effect of some organic acids on microbial quality of dressed cattle carcasses in Damietta abattoirs, Egypt. Damanhour Journal of Veterinary Sciences 5(2): 17-20.
- 11. Tserennadmid R, Takó M, Galgóczy L, Papp T, Vágvölgyi C, et al. (2010) Antibacterial Effect of Essential Oils and Interaction with Food Components. Central European Journal of Biology 5(5): 641-648.
- Edris A, Hassanin FS, Shaltout FA, Azza H Elbaba, Nairoz M Adel (2017) Microbiological Evaluation of Some Heat Treated Fish Products in Egyptian Markets. EC Nutrition 33(2): 305-316.
- 13. Chorianopoulos N, Kalpoutzakis E, Aligiannis N, Mitaku S, Nychas GJ, et al. (2004) Essential Oils of Satureja, Origanum, and Thymus Species: Chemical Composition and Antibacterial Activities against Foodborne Pathogens. Journal of Agricultural and Food Chemistry 52(26): 8261-8267.
- 14. Edris A, Hassan MA, Shaltout FA, Elhosseiny S (2013) Chemical evaluation of cattle and camel meat. Benha Veterinary Medical Journal 24(2): 191-197.
- 15. Hayouni EA, Chraief I, Abedrabba M, Bouix M, Leveau JY, et al. (2008) Tunisian Salvia officinalis L. and Schinus molle L. Essential Oils: Their Chemical Compositions and Their Preservative Effects against Salmonella Inoculated in Minced Beef Meat. International Journal of Food Microbiology 125(3): 242-251.
- Edris AM, Hassan MA, Shaltout FA, Elhosseiny S (2012) Detection of E.coli and Salmonella organisms in cattle and camel meat. Benha Veterinary Medical Journal 24(2): 198-204.
- 17. Govaris A, Solomakos N, Pexara A, Chatzopoulou PS (2010) The Antimicrobial Effect of Oregano Essential Oil, Nisin and Their Combination against *Salmonella* Enteritidis in Minced Sheep Meat during Refrigerated Storage. International Journal of Food Microbiology 137(2-3): 175-180.
- Edris AM, Hemmat MI, Shaltout FA, Elshater MA, Eman FMI (2012) Study on Incipient Spoilage of Chilled Chicken Cuts-up. Benha Veterinary Medical Journal 23(1): 81-86.
- Ramos EM, Gomide LAM (2009) Meat Quality Assessment: Fundamentals and Methodologies 1st (Edn.). UFV, Brazil, pp: 599.
- Edris AM, Hemmat MI, Shaltout FA, Elshater MA, Eman FMI (2012) Chemical Analysis Of Chicken Meat With Relation To Its Quality. Benha Veterinary Medical Journal

23(1): 87-92.

- 21. Dool HVD, Kratz PD (1963) A Generalization of the Retention Index System Including Linear Temperature Programmed Gas-Liquid Partition Chromatography. Journal of Chromatography A 11: 463-471.
- 22. Edris AM, Shaltout FA, Abd Allah AM (2005) Incidence of Bacillus cereus in some meat products and the effect of cooking on its survival. Zag Vet J 33(2): 118-124.
- 23. Adams RP (2007) Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy. Allured Publishing Corporation Illinois, pp: 803.
- 24. Edris AM Shaltout FA, Arab WS (2005) Bacterial Evaluation of Quail Meat. Benha Vet Med J 16(1): 1-14.
- 25. Aziz M, Karboune S (2016) Natural Antimicrobial/ Antioxidant Agents in Meat and Poultry Products as Well as Fruits and Vegetables: A Review. Critical Reviews in Food Science and Nutrition 58(3): 486-511.
- 26. Edris AM, Shaltout FA, Salem GH, El-Toukhy EI (2011) Incidence and isolation of *Salmonellae* from some meat products. Benha University, Egypt, pp: 172-179.
- 27. Souza AA (2015) Chemical analysis and antimicrobial potential of essential oils on enterotoxigenic Escherichia coli in the preservation of ground meat. Universidade Federal de Lavras, Brazil.
- Edris MA, Hassanin FS, Shaltout FA, Elbaba AH, Adel NM (2017) Microbiological Evaluation of Some Heat Treated Fish Products in Egyptian Markets. EC Nutrition 33(2): 305-316.
- 29. Banwart GJ (1998) Basic Food Microbiology. Springer, USA.
- Edris AM, Shaltout FA, Salem GH, El-Toukhy EI (2011) Plasmid profile analysis of Salmonellae isolated from some meat products. Benha University, Egypt, pp: 194-201.
- 31. Burt S (2004) Essential Oils: Their Antibacterial Properties and Potential Applications in Foods—A Review. International Journal of Food Microbiology 94(3): 223-253.
- 32. Ragab A, Edris AM, Shaltout FAE, Salem AM (2022) Effect of titanium dioxide nanoparticles and thyme essential oil on the quality of the chicken fillet. Benha Veterinary Medical Journal 41(2): 38-40.
- 33. De Oliveira TLC, de AraújoSoares R, Piccoli RH

Open Access Journal of Frailty Science

(2013) A Weibull Model to Describe Antimicrobial Kinetics of Oregano and Lemongrass Essential Oils against Salmonella Enteritidis in Ground Beef during Refrigerated Storage. Meat Science 93(3): 645-651.

- Hassan MA, Shaltout FA, Arfa MM, Mansour AH, Saudi KR (2013) Biochemical Studies On Rabbit Meat Related To Some Diseases. Benha Veterinary Medical Journal 25(1): 88-93.
- Gill AO, Delaquis P, Russo P, Holley RA (2002) Evaluation of Antilisterial action of Cilantro Oil on Vacuum Packed Ham. International Journal of Food Microbiology 73(1): 83-92.
- 36. Hassan MA, Shaltout FA (1997) Occurrence of Some Food Poisoning Microorganisms In Rabbit Carcasses. Alex J Vet Science 13(1): 55-61.
- Cho SH, Kim J, Oh KH, Hu JK, Seo J, et al. (2014) Outbreak of Enterotoxigenic Escherichia coli O169 Enteritis in Schoolchildren Associated with Consumption of Kimchi, Republic of Korea, 2012. Epidemiology & Infection 142(3): 616-623.
- Hassan M, Shaltout FA, Saqur N (2020) Histamine in Some Fish Products. Archives of Animal Husbandry & Dairy Science 2(1): 1-3.
- 39. Mentzer AV, Connor TR, Wieler LH, Semmler T, Iguchi A, et al. (2014) Identification of Enterotoxigenic Escherichia coli (ETEC) Clades with Long-Term Global Distribution. Nature Genetics 46(12): 1321-1326.
- 40. Hassan MA, Shaltout FA, Arafa MM, Mansour AH, Saudi KR (2013) Biochemical studies on rabbit meat related to some diseases. Benha Vet Med J 25(1): 88-93.
- 41. Jayari A, El Abed N, Jouini A, Abdul-Wahab OMS, Maaroufi A, et al. (2018) Antibacterial Activity of Thymus capitatus and Thymus algeriensis Essential Oils against Four Food-Borne Pathogens Inoculated in Minced Beef Meat. Journal of Food Safety 38: e12409.
- 42. Hassan MA, Shaltout FA, Maarouf AA, El-Shafey WS (2014) Psychrotrophic bacteria in frozen fish with special reference to pseudomonas species. Benha Vet MedJ 27(1): 78-83.
- 43. Pateiro M, Barba FJ, Domínguez R, Sant'Ana AS, Khaneghah AM, et al. (2018) Essential Oils as Natural Additives to Prevent Oxidation Reactions in Meat and Meat Products: A Review. Food Research International 113: 156-166.
- 44. Hassan MA, Shaltout FA, Arafa MM, Mansour AH, Saudi KR (2013) Bacteriological studies on rabbit meat related

to some diseases. Benha Vet Med J 25(1): 94-99.

- 45. Smith-Palmer A, Stewart J, Fyfe L (2001) The Potential Application of Plant Essential Oils as Natural Food Preservatives in Soft Cheese. Food Microbiology 18(4): 463-470.
- 46. Hassanin FS, Hassan MA, Shaltout FA, Nahla A, Shawqy, et al. (2017) Chemical criteria of chicken meat. Benha Veterinary Medical Journal 33(2): 457-464.
- 47. Santurio DF (2011) Antimicrobial activity of essential oils from condiments against samples of Escherichia coli isolated from poultry and cattle. Rural Science 41: 1051-1056.
- Hassanin FS, Hassan MA, Shaltout FA, Elrais-Amina M (2014) Clostridium Perfringens In Vacuum Packaged Meat Products. Benha Veterinary Medical Journal 26(1): 49-53.
- Jayasena DD, Jo C (2013) Essential Oils as Potential Antimicrobial Agents in Meat and Meat Products: A Review. Trends in Food Science & Technology 34(2): 96-108.
- 50. Hassanien FS, Shaltout FA, Fahmey MZ, Elsukkary HF (2020) Bacteriological quality guides in local and imported beef and their relation to public health. Benha Veterinary Medical Journal 39(1): 125-129.
- 51. Rasschaert G, Houf K, Zutter DL (2006) Impact of the Slaughter Line Contamination on the Presence of Salmonella on Broiler Carcasses. Journal of Applied Microbiology 103(2): 333-341.
- 52. Hassanin FS, Shaltout FA, Mostafa EM (2013) Parasitic affections in edible offal. Benha Vet Med J 25(1): 46-55.
- 53. Bakkali F, Averbeck S, Averbeck D, Idaomar M (2008) Biological Effects of Essential Oils-A Review. Food and Chemical Toxicology 46(2): 446-475.
- 54. Hassanin FS, Shaltout FA, Lamada HM, Allah EMA (2011) The Effect of Preservative (Nisin) on the Survival of Listeria Monocytogenes. Benha Veterinary Medical Journal, pp: 141-145.
- 55. Nazzaro F, Fratianni F, Martino DL, Coppola R, Feo VD (2013) Effect of Essential Oils on Pathogenic Bacteria. Pharmaceuticals 6(12): 1451-1474.
- 56. Khattab E, Shaltout F, Sabik I (2021) Hepatitis A virus related to foods. Benha Veterinary Medical Journal 40(1): 174-179.
- 57. Hoferl M, Buchbauer G, Jirovetz L, Schmidt E, Stoyanova

A, et al. (2009) Correlation of Antimicrobial Activities of Various Essential Oils and Their Main Aromatic Volatile Constituents. Journal of Essential Oil Research 21(5): 459-463.

- 58. Saad MS, Fahim AS, Amal AAF, Hashim FM (2022) Organophosphorus Residues in Fish in Rural Areas. Journal of Progress in Engineering and Physical Science 1(1): 27-31.
- Saif M, Saad S, Hassanin FS, Shaltout FA, Marionette Z (2019) Molecular detection of enterotoxigenic Staphylococcus aureus in ready-to-eat beef products. Benha Veterinary Medical Journal 37(1): 7-11.
- 60. Aureli P, Costantini A, Zolea S (1992) Antimicrobial activity of some plant essential oils against Listeria monocytogenes. Journal of Food Protection 55(5): 344-348.
- Saif M, Saad SM, Hassanin FS, Shaltout FA, Marionette Z (2019) Prevalence of methicillin-resistant Staphylococcus aureus in some ready-to-eat meat products. Benha Veterinary Medical Journal 37: 12-15.
- 62. Badi HN, Yazdani D, Ali SM, Nazari F (2004) Effects of spacing and harvesting time on herbage yield and quality/quantity of oil in thyme, *Thymus vulgaris L*. Industrial Crops and Products 19(3): 231-236.
- 63. Farag AA, Saad MS, Fahim AS, Hashim FM (2023) Studies on Pesticides Residues in Fish in Menofia Governorate. Benha Journal of Applied Sciences 8(5): 323-330.
- 64. Reza SMA, Rahman A, Lee J, Kang SC (2010) Potential roles of essential oil and organic extracts of Zizyphus jujuba in inhibiting food-borne pathogens. Food Chemistry 119(3): 981-986.
- 65. Farag AA, Saad SM, Shaltout FA, Mohammed HF (2023) Organochlorine Residues in Fish in Rural Areas. Benha Journal of Applied Sciences 8 (5): 331-336.
- 66. Ayala-Zavala JF, Oliu GO, Serrano IO, Aguilar GAG, Parrilla EA, Belloso OM (2007) Bio-preservation of fresh-cut tomatoes using natural antimicrobials. European Food Research and Technology 226: 1047-1055.
- 67. Shaltout FA, Hussein MN, Elsayed NK (2023) Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. Journal of Advanced Veterinary Research 13(2): 157-160.
- 68. Shaltout FA, Heikal GI, Ghanem AM (2022) Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to Aspergillus flavus virulent factors. benha veteriv medical journal

veterinary 42(1): 12-16.

- 69. Darwish W, Shaltout FA, Salem RM, Eldiasty EM, Diab FA (2022) Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. Journal of Advanced Veterinary Research 12(5): 641-644.
- Shaltout FA, Barr AAH, Abdelaziz ME (2022) Pathogenic Microorganisms in Meat Products. Biomedical Journal of Scientific & Technical Research 41(4): 32836-32843.
- 71. Shaltout FA, Thabet MG, Koura HA (2017) Impact of Some Essential Oils on the Quality Aspect and Shelf Life of Meat. Benha Veterinary Medical Journal 7(2): 647.
- 72. Shaltout FA, Mohammed IZ, Afify ESA (2020) Bacteriological profile of some raw chicken meat cuts in Ismailia city, Egypt. Benha Veterinary Medical Journal 39(1): 11-15.
- 73. Shaltout FA, Mohammed IZ, Afify ESA (2020) Detection of E. coli 0157 and Salmonella species in some raw chicken meat cuts in Ismailia province, Egypt. Benha Veterinary Medical Journal 39(1): 101-104.
- 74. Shaltout FA, El-diasty EM, Hassan MAS (2019) Hygienic quality of ready to eat cooked meat in restaurants at cairo governorate. Journal of Global Biosciences 8(12): 6627-6641.
- 75. Shaltout FA, Nasief MZ, Lotfy LM, Gamil BT (2019) Microbiological status of chicken cuts and its products. Benha Veterinary Medical Journal 37(1): 57-63.
- 76. Shaltout FA (2019) Poultry Meat. Scholarly Journal of Food and Nutrition 22: 1-2.
- 77. Shaltout FA (2019) Food Hygiene and Control. Food Science and Nutrition Technology 4(5): 1-2.
- Hassanin FS, Shaltout FA, Homouda SN, Arakeeb SM (2019) Natural preservatives in raw chicken meat. Benha Veterinary Medical Journal 37(1): 41-45.
- Hazaa W, Shaltout FA, El-Shate M (2019) Prevalence of some chemical hazards in some meat products. Con Dai & Vet Sci 3(4): 313-317.
- 80. Hazaa WMA, Shaltout FA, Shater ME (2019) Identification of Some Biological Hazards in Some Meat Products. Benha Veterinary Medical Journal 37(1): 27-31.
- 81. Gaafar R, Hassanin FS, Shaltout FA, Marionette Zaghloul (2019) Molecular detection of enterotoxigenic Staphylococcus aureus in some ready to eat meat-based sandwiches. Benha Veterinary Medical Journal 37(1):

Open Access Journal of Frailty Science

22-26.

- 82. Gaafar R, Hassanin FS, Shaltout FA, Zaghloul M (2019) Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubiya Governorate, Egypt. Benha Veterinary Medical Journal 37(1): 16-21.
- 83. Saad SM, Shaltout FA, Elroos NAA, El-nahas SB (2019) Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. Journal of Food Science and Nutrition Research 2(1): 13-21.
- 84. Saad SM, Shaltout FA, Elroos NAA, El-nahas SB (2019) Incidence of Staphylococci and E. coli in Meat and Some Meat Products. EC Nutrition 14(6).
- 85. Shaltout FA (2019) Pollution of Chicken Meat and Its Products by Heavy Metals. Research and Reviews on Healthcare Open Access Journal 4(3): 381-382.
- Shaltout FA, EL-diasty EM, Mohamed MSM (2018) Effects of chitosan on quality attributes fresh meat slices stored at 40C. Benha Veterinary Medical Journal 35(2): 157-168.
- Shaltout F (2004) Salmonella enterica serovar Enteritidis in poultry meat and their epidemiology. Vet Med J Giza 52: 429-436.
- Shaltout FA, El-Shorah HF, El Zahaby DI, Lotfy LM (2018) Bacteriological Profile of Chicken Meat Products. Food Nutr Current Res 1(3): 83-90.
- 89. Shaltout FA, El-Shater MAH, Abd El-Aziz WM (2015) Bacteriological assessment of Street Vended Meat Products sandwiches in kalyobia Governorate. Benha Veterinary Medical Journal 28(2): 58-66.
- 90. Shaltout FA, El shatter MA, Fahim HM (2019) Studies on Antibiotic Residues in Beef and Effect of Cooking and Freezing on Antibiotic Residues Beef Samples. Scholarly Journal of Food and Nutritionm 2(1): 1-4.
- 91. Shaltout FA, Zakaria IM, Nabil ME (2018) Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to Clostridium perfringens. Benha Veterinary Medical Journal 33(2): 292-304.
- 92. Shaltout FA, Maarouf AAA, ES Elkhouly M (2017) Bacteriological Evaluation of Frozen Sausage. Nutrition and Food Toxicology 1(5): 174-185.
- 93. Shaltout FA, Ei E, Mm A (2019) Molecular Diagnosis of Salmonellae in Frozen Meat and Some Meat Products. Nutrition and Food Technology Open Access 5(1): 1-6.

- 94. Shaltout FA, Ali AM, Rashad SM (2016) Bacterial Contamination of Fast Foods. Benha Journal of Applied Sciences (BJAS) 1(2): 45-51.
- 95. Shaltout FA, Zakaria IM, Eltanani J, Elmelegy AS (2015) Microbiological status of meat and chicken received to University student hostel. Benha Veterinary Medical Journal 29(2): 187-192.
- 96. Saad SM, Edris AM, Shaltout FA, Edris S (2012) Isolation and identification of salmonellae and E.coli from meat and poultry cuts by using A multiplex PCR. Benha Vet Med J special issue 16-26.
- 97. Saad SM, Shaltout FA (1998) Mycological Evaluation of camel carcasses at Kalyobia Abattoirs. Veterinary Medical Journal Giza 46(3): 223-229.
- 98. Saad SM, Shaltout FA, Elroos NAA, El-nahas SB (2019) Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. J Food Sci Nutr Res 2(1): 012-020.
- 99. Saad SM, Hassanin FS, Shaltout FA, Nassif MZ, Seif MZ (2019) Prevalence of Methicillin-Resistant Staphylococcus Aureus in Some Ready-to-Eat Meat Products. American Journal of Biomedical Science & Research 4(6): 460-464.
- Saad SM, Shaltout FA, Elroos NAA, El-nahas SB (2019) Incidence of Staphylococci and E. coli in Meat and Some Meat Products. EC Nutrition 14(6).
- 101. Shaltout FA, Riad EM, Ahmed TES, AbouElhassan A (2017) Studying the Effect of Gamma Irradiation on Bovine Offal's Infected with Mycobacterium tuberculosis Bovine Type. Journal of Food Biotechnology Research 1(6): 1-5.
- 102. Shaltout FA, Zakaria IM, Nabil ME (2018) Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to Clostridium perfringens. Nutrition and Food Toxicology 2(5): 429-438.
- 103. Shaltout FA, Mohamed, Hassan A, Hassanin FS (2004) Thermal Inactivation of Enterohaemorrhagic Escherichia Coli O157:H7 and Its Senstivity to Nisin and Lactic Acid Cultures. The First Scientific Conference, Moshtohor, Zagazig University Benha branch, Egypt.
- 104. Shaltout FA, El-diasty EM, Elmesalamy M, Elshaer M (2014) Study on fungal contamination of some chicken meat products with special reference to 2 the use of PCR for its identification. Conference, Veterinary Medical Journal – Giza 60: 1-10.

- Shaltout FA (2002) Microbiological Aspects of Semi-cooked chicken Meat Products. Benha Veterinary Medical Journal 13(2): 15-26.
- 106. Shaltout FA, Thabet MG, Koura HA (2017) Impact of some essential oils on the quality aspect and shelf life of meat. Benha Veterinary Medical Journal 33(2): 351-364.
- 107. Shaltout FA, Farouk M, Ibrahim HAA, Afifi MEM (2017) Incidence of Coliform and Staphylococcus aureus in ready to eat fast foods. Benha Veterinary Medical Journal 32(1): 13-17.
- 108. Shaltout FA, Zakaria IM, Nabil ME (2017) Detection and typing of Clostridium perfringens in some retail chicken meat products. Benha Veterinary Medical Journal 33(2): 283-291.
- 109. Shaltout FA (1992) Studies on Mycotoxins in Meat and Meat by Products. M.V.Sc Thesis Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
- 110. Shaltout FA (1996) Mycological and Mycotoxicological profile of Some Meat products. Ph.D.Thesis, Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
- 111. Shaltout FA (1998) Proteolytic Psychrotrophes in Some Meat products. Alex Vet Med J 14(2): 97-107.
- 112. Shaltout FA (1999) Anaerobic Bacteria in Vacuum Packed Meat Products. Benha Vet Med J 10(1)1-10.
- 113. Shaltout FA (2000) Protozoal Foodborne Pathogens in some Meat Products. Assiut Vet Med J 42 (84): 54-59.
- 114. Shaltout FA (2001) Quality evaluation of sheep carcasses slaughtered at Kalyobia abattoirs. Assiut Veterinary Medical Journal 46(91): 150-159.
- Shaltout FA (2002) Microbiological Aspects of Semicooked Chicken Meat Products. Benha Vet Med J 13(2): 15-26.
- 116. Shaltout FA (2003) Yersinia Enterocolitica in some meat products and fish marketed at Benha city. The Third international conference Mansoura, pp: 29-30.
- 117. Shaltout FA (2020) Microbiological quality of chicken carcasses at modern Poultry plant. Nutrition and Food Processing 3(1).
- 118. Shaltout FA, Abdel Aziz AM (2004) Salmonella enterica Serovar Enteritidis in Poultry Meat and their Epidemiology. Vet Med J Giza 52(3): 429-436.
- 119. Shaltout FA, Abdel Aziz AM (2004) Escherichia Coli

Strains in Slaughtered Animals and Their Public Health Importence. J Egypt Vet Med Association 64(2): 7-21.

- Shaltout FA, Amin R, Nassif MZ, Shimaa A, Wahab A (2014) Detection of aflatoxins in some meat products. Benha veterinary medical journal 27(2): 368-374.
- 121. Shaltout FA, Jehan A Riad EM, Elhasan A, Asmaa A (2012) Improvement of microbiological status of oriental sausage. Journal of Egyptian Veterinary Medical Association 72(2): 157-167.
- 122. Shaltout FA, Daoud JR (1996) Chemical analytical studies on rabbit meat and liver. Benha Vet Med J 8(2): 17-27.
- 123. Shaltout FA, Edris AM (1999) Contamination of shawerma with pathogenic yeasts. Assiut Veterinary Medical Journal 40(64): 34-39.
- 124. Shaltout FA, Eldiasty E, Mohamed MS (2014) Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. Animal Health Research Institute, pp: 79-89.
- 125. Shaltout FA, El-diasty EM, Salem RM, Asmaa MAH (2016) Mycological quality of chicken carcasses and extending shelf life by using preservatives at refrigerated storage. Veterinary Medical Journal 62(3): 1-7.
- 126. Shaltout FA, Salem R, Eldiasty E, Fatema D (2016) Mycological evaluation of some ready to eat meat products with special reference to molecular chacterization. Veterinary Medical Journal 62(3): 9-14.
- 127. Shaltout FA, Elshater M, el-Aziz AMW (2015) Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate. Benha Vet Med J 28(2): 58-66.
- 128. Shaltout FA, Gerges MT, Shewail AA (2014) Impact of Organic Acids and Their Salts on Microbial Quality and Shelf Life of Beef meat. Assiut veterinary medical journal 1(2): 360-370.
- 129. Shaltout FA, Ghoneim AM, Essmail ME, Yousseif A (2001) Studies on aflatoxin B1 residues in rabbits and their pathological effects. J Egypt Vet Med Association 61(2): 85-103.
- 130. Shaltout FA, Hanan MT, Lawendy El (2003) Heavy Metal Residues In Shawerma. Beni-Suef Vet Med J 13(1): 213-224.
- 131. Shaltout FA, Hashim MF (2002) Histamine in salted,

Smoked and Canned Fish products. Benha Vet Med J 13 (1): 1-11.

- 132. Shaltout FA, Hashim MF, Elnahas S (2015) Levels of some heavy metals in fish (tilapia nilotica and Claris lazera) at Menufia Governorate. Benha Vet Med J 29(1): 56-64.
- Shaltout FA, Ibrahim HM (1997) Quality evaluation of luncheon and Alexandrian sausage. Benha Vet Med J 10(1): 1-10.
- 134. Shaltout FA, Nassif M, Shakran A (2014) Quality of battered and breaded chicken meat products. Global Journal of Agriculture and Food Safety Science 1(2): 283-299.
- 135. Shaltout FA, Amani M, Salem AH, Mahmoud KA (2013) Bacterial aspect of cooked meat and offal at street vendors level. Benha veterinary medical journal 24(1): 320-328.
- 136. Shaltout FA, Salem RM (2000) Moulds, aflatoxin B1 and Ochratoxin A in Frozen Livers and meat products. Vet Med J Giza 48(3): 341-346.
- 137. Yasser H, Al-Tarazi A, Al-Zamil A, Shaltout FA, Abdel-Samei H (2002) Microbiological status of raw cow milk marketed in northern Jordan. AVMJ 49(96): 180-194.
- 138. Shaltout FA, Zakaria IM, Nabil ME (2018) Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to Clostridium perfringens. Nutrition and Food Toxicology 2(5): 429-438.
- 139. Shaltout FA, El-diasty EM, Mohamed MS (2014) Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. 1st Scientific conference of food safety and Technology, pp: 79-89.
- 140. Shaltout FA, El-diasty EM, Salem RM, Asmaa MA, Hassan MT (2016) Mycological quality of chicken carcasses and extending shelf-life by using preservatives at refrigerated storage. Veterinary Medical Journal – Giza 62(3): 1-10.
- 141. Shaltout FA, Salem RM, El-Diasty EM, Hassan WIM

(2019) Effect of Lemon Fruits and Turmeric Extracts on Fungal Pathogens in Refrigerated Chicken Fillet Meat. Global Veterinaria 21(3): 156-160.

- 142. Shaltout FA, El-diasty EM, Elmesalamy M, Elshaer M (2014) Study on fungal contamination of some chicken meat products with special reference to 2 the use of PCR for its identification. Conference, Veterinary Medical Journal – Giza 60: 1-10.
- 143. Shaltout FA, Salem RM, El-diasty E, Fatema AH, Diab R (2016) Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. Veterinary Medical Journal – Giza 62(3): 9-14.
- 144. Shaltout FA, Ahmed AA, Maarouf E, Ahmed MK (2018) Heavy Metal Residues in chicken cuts up and processed chicken meat products. Benha veterinary medical journal 34(1): 473-483.
- 145. Shaltout FA, Hanan M, Lamada E, Edris AM (2020) Bacteriological examination of some ready to eat meat and chicken meals. Biomed J Sci & Tech Res 38(2): 76-79.
- 146. Sobhy A, Shaltout F (2020) Prevalence of some food poisoning bacteria in semi cooked chicken meat products at Qaliubiya governorate by recent Vitek 2 compact and PCR techniques. Benha Veterinary Medical Journal 38(2): 88-92.
- 147. Shaltout FA, Riad EM, Abou E, Asmaa A (2017) prevalence Of Mycobacterium Tuberculosis In Imported cattle Offals And Its lymph Nodes. Veterinary Medical Journal -Giza (VMJG) 63(2): 115 - 122.
- 148. Sobhy A, Shaltout F (2020) Detection of food poisoning bacteria in some semi-cooked chicken meat products marketed at Qaliubiya governorate. Benha Veterinary Medical Journal 38(2): 93-96.
- 149. Shaltout FA (2024) Abattoir And Bovine Tuberculosis as A Reemerging Foodborne Diseas. Clinical Medical Reviews and Report 6(1): 1-7.
- Shaltout FA (2023) Viruses in Beef, Mutton, Chevon, Venison, Fish and Poultry Meat Products. Food Science & Nutrition Technology 8(4): 1-10.

9