

# Sugarcane Bagasse and Passion Fruit Rind Flour as Substrates for Cellulase Production by *Bacillus* Sp. SMIA-2 Strain Isolated from Brazilian Soil

Costa EA<sup>1</sup>, Nunes R<sup>2</sup>, Cruz E<sup>1</sup>, Ladeira SA<sup>1</sup>, Moraes LP<sup>1</sup>, Carvalho RV<sup>2</sup> and Martins MLL<sup>1\*</sup>

<sup>1</sup>Food Technology Laboratory, Universidade Estadual do Norte Fluminense, Brazil

<sup>2</sup>Food Microbiology Laboratory, Universidade Federal do Espírito Santo. Alto Universitário, Brazil

**\*Corresponding author:** Meire Lelis Leal Martins, Food Technology Laboratory, Universidade Estadual do Norte Fluminense, Avenida Alberto Lamego 2000, 28013-602, Campos dos Goytacazes, Rio de Janeiro, Brazil, E-mail: meire@uenf.br

## Research Article

Volume 2 Issue 1

**Received Date:** February 18, 2017

**Published Date:** March 13, 2017

## Abstract

The application of wastes from the food processing industry as carbon sources in enzyme production processes reduces the cost of production, and helps solving problems related to their disposal. In this work, we demonstrated that the use of sugarcane bagasse as cellulosic substrate, in combination with passion fruit rind flour, as co-substrate and corn steep liquor as nitrogen source can be successfully employed for the production of avicelases (avicel-hydrolyzing enzymes) by *Bacillus* sp. SMIA-2. This would promote the use of these agricultural byproducts as novel and cost-effective culture media for the production of the enzyme. The maximum avicelase activity was obtained when *Bacillus* sp SMIA-2 was grown in liquid medium containing 0.625% (w/v) sugarcane bagasse, 0.625% (v/v) corn steep liquor and 0.75% (w/v) passion fruit peel flour for 168 hours.

**Keywords:** Bacillus Sp; Cellulase; Corn Steep Liquor; Passion Fruit Peel Flour; Sugarcane Bagasse

## Introduction

The complete enzymatic hydrolysis of cellulosic materials requires at least three different types of cellulases: endoglucanase (CM cellulase, EC 3.2.1.4), exoglucanase (Avicelase, EC 3.2.1.91) and  $\beta$ -D-glucosidase ( $\beta$ -D-glucoside glucohydrolase, EC 3.2.1.21) [1]. Among these types of cellulases, the avicelases seem to catalyze most of the bond-cleavages in the saccharification of crystalline cellulose and are usually the major component of cellulase preparations, especially in the case of current fungus-derived commercial enzymes [2]. Cellulases are used in different industrial applications such as pulp industry, food and animal feed, textile and detergent

industry [3]. They are also applied in the production of bioethanol from lignocellulosic materials [4, 5].

The present cellulase toolbox is not sufficient to meet the industrial demand. For this reason, the major challenge in the development of economically feasible bioprocess for cellulase production is to identify the potential microorganisms, composition of media and the optimization of various process parameters that affect microbial growth and enzyme production [6]. The carbon source used in cultivations is one of the most important factors affecting the cost and yield of cellulase production. In view thereof, biosynthesis of cellulase has also been conducted on lignocellulosic materials, such as sugarcane bagasse [6], which is abundantly available at low cost as a

byproduct from the sugar industry in Brazil [7]. Since the cellulose component of sugarcane bagasse is a structural polymer and is protected against enzymatic attack by the surrounding matrix of lignin and hemicellulose, its pretreatment removes lignin, reduces crystallinity of cellulose, which may enhance cellulase production by microorganisms [8].

Employment of Response Surface Methodology (RSM) is considered the most promising tool in the optimization of process parameters for better yield [9,10]. In this work, the Central Composite Design (CCD) was adopted to maximize the production of cellulase by cultures of thermophilic *Bacillus* sp SMIA-2 containing sugarcane treated with alkali and corn steep liquor. Subsequently, it was evaluated the use of passion fruit rind flour as co-substrate, as an alternative to the conventional rich medium.

## Material and Methods

### Organism

The present study used a thermophilic *Bacillus* sp strain SMIA-2, previously isolated from a soil sample collected in the city of Campos dos Goytacazes, Rio de Janeiro, Brazil. Phylogenetic analysis revealed that the bacteria were closely related to *Bacillus caldoolyolyticus* and *Bacillus* sp strain AK1, and these three organisms exhibited levels of ribosomal DNA sequence homology of 94% [11].

### Enzyme Production

The culture medium used in this work for cellulase production contained (g/L): KCl-0.3, MgSO<sub>4</sub>-0.5, K<sub>2</sub>HPO<sub>4</sub>-0.87, CaCl<sub>2</sub>-0.29, ZnO-2.03x10<sup>-3</sup>, FeCl<sub>3</sub>.6H<sub>2</sub>O-2.7x10<sup>-2</sup>, MnCl<sub>2</sub>.4H<sub>2</sub>O-1.0x10<sup>-2</sup>, CuCl<sub>2</sub>.2H<sub>2</sub>O-8.5x10<sup>-4</sup>, CoCl<sub>2</sub>.6H<sub>2</sub>O-2.4x10<sup>-3</sup>, NiCl<sub>3</sub>.6H<sub>2</sub>O-2.5x10<sup>-4</sup> and H<sub>3</sub>BO<sub>3</sub>-3.0x10<sup>-4</sup>. In first set of experiments, sugarcane bagasse (SCB) treated with alkali (81.05% cellulose, 18.75% hemicellulose, 5.45% lignine) was used as a source of cellulose [12] and commercial corn steep liquor (Sigma Aldrich), as a nitrogen source. The SCB and corn steep liquor (CSL) concentrations were adjusted for each value according to central composite design (CCD), as presented in (Table 1).

The pH was adjusted to 7.2 with 1.0 M NaOH and the medium was sterilized by steam-autoclaving at 121°C, 1 atm for 15 minutes. The medium (50 mL in 250 mL Erlenmeyer flasks) was inoculated with 1 mL of an standard overnight culture (initial number of cells 10<sup>4</sup>) and incubated at 50°C in an orbital shaker (Thermo

Forma, Ohio, USA) operated at 150 rpm. Triplicate flasks were withdrawn at regular intervals and the contents were then centrifuged (HERMLEZ 382K, Wehingen, Germany) at 15,500 g for 15 min, at 4 °C, and the cell free supernatant was used as crude enzyme preparation.

The effect of different co-substrates (0.5%, w/v) on avicelase secretion was assessed by supplementing the culture medium with passion fruit rind flour (obtained from a local market), apple pectin, fructose, glucose, xylose, lactose and cellobiose. In addition, the concentration of passion fruit rind flour (PFRF) in the culture medium ranged from 0.25% to 1.5% (w/v).

### Enzyme Assay

The cellulolytic enzyme activities were determined using the dinitrosalicylic acid method [13], which measures reducing sugars. The reaction mixture containing 0.5 mL of 1% (w/v) avicel, PH-101 prepared in 10 mM sodium phosphate buffer, pH 7.5, and 0.5 mL of appropriate concentration of enzyme solution, was incubated at 70°C. After 10 min of reaction, 1 mL of dinitrosalicylic acid reagent was added and boiled in water bath for 5 min. The resulting samples were then cooled to room temperature, and the absorbance was measured at 540 nm. One unit (U) of activity was defined as 1 μmole of glucose equivalent released per minute under the above assay conditions, by using a glucose standard curve. Appropriate controls were conducted in parallel with all assays. Enzyme blank containing 0.5 mL of 10 mM sodium phosphate buffer and 0.5 mL of 1% (w/v) substrate solution were run. To exclude the background of reducing sugars found in the enzyme supernatant from the results, a substrate blank containing 0.5 mL of 10 mM sodium phosphate buffer and 0.5 mL enzyme solution was also run. The absorbance of the enzyme blank sets and the substrate blank were subtracted from the absorbance of the activity assay. All of the samples were run in triplicate, while the blanks were run in duplicate.

### Experimental Design and Statistical Analysis

The surface-response methodology (SRM) was used to obtain a model for cellulase activity. To evaluate the effects of sugarcane bagasse and corn steep liquor concentration and culture time on the production of cellulase, a central composite design (CCD) 2<sup>3</sup> was constructed. The factorial planning had three central points and yielded a total of 17 treatments. The factors and levels studied are described in Table 1.

Treatments	SCB (% w/v)	CSL (% v/v)	Culture time (h)	Avicelase activity (U.mg protein <sup>-1</sup> )
1	0.4 (-1)	0.4 (-1)	139 (-1)	1.97
2	0.85 (+1)	0.4 (-1)	139 (-1)	0.44
3	0.4 (-1)	0.85 (+1)	139 (-1)	0.13
4	0.85 (+1)	0.85 (+1)	139 (-1)	0
5	0.4 (-1)	0.4 (-1)	197 (+1)	1.46
6	0.85 (+1)	0.4 (-1)	197 (+1)	0.7
7	0.4 (-1)	0.85 (+1)	197 (+1)	0
8	0.85 (+1)	0.85 (+1)	197 (+1)	0
9	0.25 (-1.68)	0.625 (0)	168 (0)	2.29
10	1.0 (+1.68)	0.625 (0)	168 (0)	0.09
11	0.625 (0)	0.25 (-1.68)	168 (0)	0.06
12	0.625 (0)	1.0 (+1.68)	168 (0)	0
13	0.625 (0)	0.625 (0)	120 (-1.68)	2.47
14	0.625 (0)	0.625 (0)	216 (+1.68)	0.25
15	0.625 (0)	0.625 (0)	168 (0)	2.45
16	0.625 (0)	0.625 (0)	168 (0)	2.73
17	0.625 (0)	0.625 (0)	168 (0)	2.41

Table 1: Matrix of CCD 2<sup>3</sup> (real and coded values) used and its response (avicelase activity).

The results were analyzed using the Statistical software system, version 5.0. In this context, the F test was used as a validation criterion of statistical significance of the models obtained at a confidence level of 95%.

The optimization of condition was performed using CCD and surface-response was produced with fixed central points of 0.625% sugarcane bagasse, 0.625% corn steep liquor and incubation time of 168 hours. The experimental model can be expressed as follows:

$$Y = bo + \sum_{i=1}^4 bi xi + \sum_{i=1}^4 bii xi^2 + \sum_{i \neq j=1}^4 \sum bij xi xj \quad (1)$$

Where *bo*, *bi*, *bii* and *bij* are the intercept terms, linear, quadratic coefficient and interactive coefficient, respectively, and *xi* and *xj* are coded independent variables.

### Scanning Electron Microscopy

The scanning electron microscopy (SEM) was used to obtain the degree of degradation of SCB and PFPF caused by *Bacillus* sp SMIA-2. Briefly, 72h bacterial cultures were harvested from culture media with 0.5% (v/v) CSL and

0.5% (w/V) of each of these substrates and fixed for 2 hours in 5% glutaraldehyde, 0.05 M sodium phosphate buffer, pH 7.1. Subsequently, the samples were washed 3 times in the same buffer and dehydrated in a graded alcoholic series (20, 40, 60, 70, 80, 90, 96 and 100%) for 15 min each. After dehydration, the samples were dried with CO<sub>2</sub> using a BAL-TEC CPD 030 Critical Point Dryer and then mounted in an aluminum sample holder and covered with a layer of palladium in a Bal-tec SCD 050 on atmosphere of argon for 120 seconds. Thereafter, the samples were observed on a ZEISS DSEM 962 scanning electron microscope (Lichtenstein, Germany) at the UENF Center for Bioscience and Biotechnology.

### Results

Initially, a central composite design (CCD) 2<sup>3</sup> was constructed to evaluate the effects of sugarcane bagasse and corn steep liquor concentration and culture time on avicelase production by *Bacillus* sp SMIA-2. According to Table 1, a variation in avicelase activity was observed from 0.0 (Treatments 4, 7, 8 and 12) to 2.73 U.mg protein<sup>-1</sup> (Treatment 16). Comparing the parameters studied in these experiments, it seems that the maximum avicelase activity was obtained when *Bacillus* sp SMIA-2 was grown

in liquid medium containing 0.625% (w/v) SCB and 0.625% (v/v) CSL for 168 hours.

The statistical significance of the model equation was assessed by an F-test (ANOVA) and the data are shown in Table 2. An equation for avicelase activity (Equation 2)

was developed based on a regression analysis of the following experimental data:

$$Y = 2.544 - 0.354 x_1 - 0.401 x_1^2 - 0.325 x_2 - 0.959 x_2^2 - 0.300 x_3 - 0.4778 x_3^2 + 0.272 x_1 x_2 + 0.113 x_1 x_3 + 0.015 x_2 x_3 \quad (2)$$

Variable	Mean square	(Degrees of freedom)	Sum of squares	Fcal	Ftab (0.5)	p < 0.05
Avicelase activity	19.83568	9	2.203964	5.696323	3.68	
Regression	2.70837	7	0.38691			
Residues	2.64722	5	0.529444	17.31624	19.3	
Lack of adjustment	0.06115	2	0.030575			
Pure error	22.54405	16				
Total error				R <sup>2</sup> =	87.98632	

Table 2: ANOVA for the variables of response surface quadratic model for avicelase production

The outcome of ANOVA analysis revealed that the adjusted model was significant, according to the analysis of the F test. The regression model for avicelase production was highly significant ( $p < 0.05$ ), with a satisfactory value of determination coefficient ( $R^2 = 87.98$ ). The response surface was produced according to Rodrigues and Lemma [14].

The response surface and contour plot figures obtained by the analysis of the experimental data of CCD showed a relationship between two variables at time. The non-explicit variables were fixed at the central point (level 0) for the surface construction. As showed in Figure 1 an initial increase in incubation time with simultaneous increase in SCB concentration, when keeping the CSL concentration constant at 0.625% (v/v), resulted in an increase of avicelase production. However, increasing SCB concentration beyond this limit decreased the avicelase production. In addition, an initial increase in incubation time, with simultaneous increase in CSL concentration, at constant SCB concentration (0.625%, w/v), increased avicelase production. However, when the concentration was higher than 0.625% (w/v), the enzyme production decreased. Finally, an increase in SCB concentration with simultaneous increase in CSL, with the fixed coded value of incubation time (168 h) led to an initial increase in avicelase production until reaching the optimal enzyme, 0.625% (w/v), for both ingredients, which was followed by a decrease.




Figure 1: Three-dimensional response surface plot for: Effect of incubation time and SCB concentration on avicelase production by *Bacillus* sp SMIA-2 at a constant CSL concentration (A); Effect of incubation time and CSL concentration on avicelase production by *Bacillus* sp SMIA-2 at a constant SCB concentration (B); Effect of SCB and CSL concentration on avicelase production by *Bacillus* sp SMIA-2 at a constant incubation time (C). Dark red color indicates high activity, while green and yellow color indicates low avicelase activity.

The addition of different co-substrates to the optimized medium improved avicelase production. The supplementation of fructose, xylose, lactose and cellobiose had little effect on avicelase production, while glucose, pectin and PFRF enhanced enzyme production. Among those, PFRF provided the highest levels of cellulase activity, around 30%, compared to the control.

Figure 2 shows the effect of different incubation periods on the production of avicelase by *Bacillus* sp SMIA-2 grown in the optimized culture medium supplemented with PFRF (0.5%, w/v). The maximum avicelase activity ( $3.6 \text{ U.mgprotein}^{-1}$ ) was obtained after 168 h of incubation culture. The visualization of *Bacillus* sp SMIA-2 cells growing on SCB and PFRF was performed using scanning electron microscopy (SEM) studies (Figure 3). *Bacillus* sp SMIA-2 attached firmly to both SCB and PFRF, which is an important characteristic in the degradation process of these substrates. In addition, the images provide qualitative observations of the cell population. Despite the difficulty in differentiating microbial cells from SCB, the number of cells grown in this insoluble substrate (Figure 4A) was much lower compared with PFRF (Figure 4B).




Figure 2: Effect of different fermentable sugars on avicelase production by *Bacillus* sp SMIA-2 grown in submerged culture containing 0.625% (w/v) SCB and 0.625% (v/v) CSL at 50°C for 168 h.




Figure 3: Effect of incubation time on avicelase production by *Bacillus* sp SMIA-2 grown in submerged culture containing 0.625% (w/v) SCB and 0.625% (v/v) CSL and supplemented with 0.5% (w/v) PFRF at 50°C.




Figure 4: Scanning electron microscopy (SEM) images showing cells of *Bacillus* sp SMIA-2 grown on 0.5% (v/v) CSL and 0.5% (w/v) SCB (A) or 0.5% (v/v) PFRF (B) at 50°C for 72 hours.

In order to find the best concentrations of PFRF for avicelase production, different amounts were used in the culture medium, ranging from 0.25% to 1.5% (w/v). The activity of the enzyme increased at PFRF concentrations between 0.25% and 0.75% (w/v) and then dropped at higher concentration levels (Figure 5). Thus, the medium containing 0.625% (v/v) CSL, 0.625% (w/v) SCB and 0.75% (w/v) PFRF was considered the most effective for avicelase production by *Bacillus* sp strain SMIA-2.




Figure 5: Effect of different concentrations of PFRF on avicelase production by *Bacillus* sp. strain SMIA-2 grown in submerged culture containing 0.625% (w/v) SCB and 0.625% (v/v) CSL at 50 °C for 168 h. Results represent the means of three separate experiments, and bars indicate  $\pm 1$  standard deviation.

## Discussion

The optimization study of avicelase (avicel-hydrolyzing enzymes) production by thermophilic *Bacillus* sp SMIA-2 showed that the amounts of SCB and CSL in the culture medium and fermentation time strongly affected the activity of this enzyme. The microbial production of cellulases is usually induced by SCB. Therefore, cellulose-rich wastes are exploited as abundant and inexpensive growth substrates to obtain cellulases [8]. CSL is a source of amino acids, vitamins and metal ions and has been successfully used in culture media for cellulase production [15,16] instead of meat and yeast extracts, which are expensive nitrogen sources. The highest avicelase activity ( $2.73 \text{ U}\cdot\text{mg protein}^{-1}$ ) was obtained when *Bacillus* sp SMIA-2 was grown in liquid medium containing 0.625% (w/v) treated SCB and 0.625% (v/v) CSL for 168 hours, which was the central point chosen in the experimental design approach used in this study. The central point was properly chosen based on previous studies carried out in our laboratory [17]. Thus, the experimental design approach used in this study was found to be efficient for the rapid optimization of the process parameters for cellulase production by *Bacillus* sp SMIA-2. Several published works have employed the response surface methodology as a tool for optimization of the cellulase production process. However, comparison between the activities obtained with *Bacillus* sp. SMIA-2 and the published literature was hindered by the different definitions of enzymatic activity and different levels of enzyme purity used.

Cirigliano et al. [16] optimized the production of endoglucanase by *Streptomyces misionensis* strain PESB-25 using response surface methodology (RSM). A peak of endoglucanase accumulation ( $1.01 \text{ U} \times [\text{mL}\cdot\text{sup}\cdot^{-1}]$ ) was observed in a medium with SCB 1.0% (w/v) and CSL 1.2% (w/v), within three days of cultivation. The maximum CMCase activity (2040 IU/L) was achieved when *Bacillus* sp. JS14 was grown on a wheat bran concentration of 400 g/L, pH 6.5, at 40 °C and incubation period of five days [9]. The optimization of cellulase production by *Bacillus amyloliquefaciens* UNPDV-22 using RSM resulted in a 96% increase in the enzyme activity over the control of non-optimized basal medium. Optimum cellulase production of 13 U/mL was obtained at 42.24 °C, pH of 5.25, and inoculum size of 4.95% (v/v) in a fermentation medium containing wheat bran, soybean meal and malt dextrin as major nutritional factors [10]. The optimum parameters for cellulase production by *Bacillus pumilus* EWBCM1 using RSM based on CCD model were galactose of 1.0 g/L, malt extract of 0.5 g/L and incubation time of 72 hrs. Optimization with coded factor provided the maximum

cellulase production observed by the model, 0.5751 IU/ml [18].

A successful and significant improvement (1.3-fold) in avicelase production by *Bacillus* sp. SMIA-2 was obtained when the optimized culture medium was supplemented with 0.5% (w/v) PFRF. It demonstrates the benefits of using co-substrates for increasing enzyme activity. Simple sugars have often been reported to work as repressors of enzyme activity. However, the mixture of substrates, including simple sugars as co-substrates, had been evaluated for the production of crude extract containing the enzyme pools with cellulolytic activity [19]. Muthuvelayudham and Viruthagiri [20] analyzed the effect of different carbon sources such as glucose, lactose and xylose on submerged fermentation, using isolates of *T. reesei*, and verified that the combination of cellulose and xylose led to the best level of cellulase production.

The highest avicelase activity ( $3.6 \text{ U}\cdot\text{mgprotein}^{-1}$ ) secreted by *Bacillus* sp. SMI-2 grown in liquid medium containing SCB (0.625%, w/v), CSL (0.625%, v/v) and PFRF (0.5%, w/v) was reached after 168 h of culture time. This incubation period was longer than that reported for maximum avicelase production by *Bacillus* sp SMIA-2 grown on untreated SCB [17] and shorter than that reported for cellulase production by *Bacillus* sp [21].

The presence of 0.75% (w/v) PFRF supported maximal avicelase ( $4.0 \text{ U}\cdot\text{mgprotein}^{-1}$ ). Passion fruit processing generates a substantial amount of residues, including peel, which is an environmental problem [22]. Its rich composition might supply nutrients that work as growth enhancers for microbial processes. The peel of passion fruit is rich in fiber, minerals, and especially pectin. In fact, qualitative observations from SEM images of *Bacillus* sp SMIA-2 grown on SCB or PFRF showed the potential of PFRF to be used as co-substrate to increase cell population. According to Maurosa et al. [23], measurements of bacterial growth on insoluble substrates are difficult for several reasons. For example, viable counts on plates of agar media are complicated by the propensity of cellulose-decomposing cells to adhere to the substrate, which hinders the accurate dilution of cultures. These authors described a dual-staining procedure to visualize and enumerate cells in the presence of insoluble plant biomass substrates.

In conclusion, SCB combined with PFRF can be successfully used for the production of avicelase (avicel-hydrolyzing enzymes) by *Bacillus* sp. SMIA-2, opening perspectives for the use of these agricultural byproducts

as novel and cost-effective culture media for the production of the enzyme.

### Acknowledgments

The authors thank the FAPERJ — Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro for financial support and Dr Fábio Olivares for providing support in analyzing the samples using SEM.

### References

1. Singhanian RR, Sukumaran RK, Patel AK, Larroche C, Pandey A (2010) Advancement and comparative profiles in the production technologies using solid-state and submerged fermentation for microbial cellulases. *Enzyme Microbial Technology* 46(7): 541-549.
2. Liu YS, Jo JB, Zeng Y, Himmel ME, Haas T, et al. (2011) Cellobiohydrolase hydrolyzes crystalline cellulose on hydrophobic faces. *J Biol Chem* 286(13): 11195-11201.
3. Sadhu S, Maiti TK (2013) Cellulase Production by Bacteria: A Review. *British Microbiology Research Journal* 3(3): 235-258.
4. Karmakar M, Ray RR (2011) Current Trends in Research and Application of Microbial Cellulases. *Research Journal of Microbiology* 6(1): 41-53.
5. Shi J, Ebrik MA, Yang B, Garlock RJ, Balan V, et al. (2011) Application of cellulase and hemicellulase to pure xylan, pure cellulose, and switchgrass solids from leading pretreatments. *Bioresour Technol* 102(24): 11080-11088.
6. Sajith S, Priji P, Sreedevi S, Benjamin S (2016) An Overview on Fungal Cellulases with an Industrial Perspective. *Journal of Nutrition and Food Science* 6: 461.
7. Castro AM, Pereira JRN (2010) Production, properties and application of cellulases in the hydrolysis of agroindustrial residues. *Química Nova* 33(1): 181-188.
8. Ashfaque M, Solomon S, Pathak N (2014) Optimization of pretreatment and fermentation conditions for production of extracellular cellulase complex using sugarcane bagasse. *Bioinformation* 10(10): 606-610.
9. Singh J, Kaur P (2012) Optimization of Process Parameters for Cellulase Production from *Bacillus* sp. JS14 in Solid Substrate Fermentation Using Response Surface Methodology. *Braz Arch Biol Technol* 55(4): 505-512.
10. Vasudeo Z, Lew C (2011) Optimization of Culture Conditions for Production of Cellulase by a Thermophilic *Bacillus* Strain. *Journal of Chemistry Engineering* 5: 521-527.
11. Souza AN, Martins MLL (2001) Isolation, properties and kinetics of growth of a thermophilic *Bacillus*, *Brazilian Journal of Microbiology* 32(4): 271-275.
12. Paixão SM, Ladeira SA, Silva TP, Arez BF, Roseiro JC, et al. (2016) Sugarcane bagasse delignification with potassium hydroxide for enhanced enzymatic hydrolysis. *International Journal to Further the Chemical Sciences* 6(2): 1042-1052.
13. Miller GL (1959) Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry* 31(3): 426-428.
14. Rodrigues MI, Lemma AF (2009) Planejamento de Experimentos e otimização de processos. (2<sup>nd</sup> Ed), Campinas SP, pp. 358.
15. Oliveira LRC, Barbosa JB, Martins MLL, Martins M (2014) Extracellular production of avicelase by the thermophilic soil bacterium *Bacillus* sp. SMIA-2. *Acta Scientiarum* 36(2): 215-222.
16. Cirigliano F, Novaes M, Rezende RC, Gravina MPO, Pereira PHF, et al. (2013) *Streptomyces misionensis* PESB-25 produces a thermoacidophilic endoglucanase using sugarcane bagasse and corn steep liquor as the sole organic substrates. *BioMed Research International*.
17. Ladeira SA, Cruz E, Delatorre AB, Barbosa JB, Martins MLL (2015) Cellulase production by thermophilic *Bacillus* sp. SMIA-2 and its detergent compatibility. *Electronic Journal of Biotechnology* 18: 110-115.
18. Shankar T, Isaiarasu L (2012) Statistical Optimization for Cellulase Production by *Bacillus pumilus* EWBCM1 Using Response Surface Methodology. *Global Journal of Biotechnology & Biochemistry* 7(1): 1-6.
19. Mesa L, Salvador CA, Herrera M, Carrazana DI, Gonzalez E (2016) Cellulases by *Penicillium* sp. in different culture conditions. *Bioethanol* 2(1): 84-93.

20. Muthuvelayudham R, Viruthagiri T (2006) Fermentative production and kinetics of cellulase protein on *Trichoderma reesei* using sugarcane bagasse and rice straw. *African Journal of Biotechnology* 5(20): 1873-1881.
21. Sadhu S, Ghosh PK, De TK, Maiti TK (2013) Optimization of cultural condition and synergistic effect of lactose with carboxymethyl cellulose on cellulase production by *Bacillus* sp. isolated from fecal matter of elephant (*elephas maximus*). *Advances in Microbiology* 3(3): 280-288.
22. Panda SK, Mishra SS, Kayitesi E, Ray RC (2016) Microbial-processing of fruit and vegetable wastes for production of vital enzymes and organic acids: Biotechnology and scopes. *Environmental Research* 146: 161-172.
23. Maurosa AJG, Lee SJ, Hazen SP, Leschine SB (2016) Direct imagebased enumeration of *Clostridium* phytofermentans cells on insoluble plant biomass growth substrates. *Applied and Environmental Microbiology* 82(3): 972-978.