



# A Review of Endophytic Fungi and their Applications in Different Fields of Biotechnology

Patyal U<sup>1</sup>, Kaur M<sup>2\*</sup>, Faizan M<sup>3†</sup>, Soysal S<sup>4</sup>, Djalovic I<sup>5</sup> and Trivan G<sup>6</sup>

<sup>1</sup>Department of Biosciences and Technology, Maharishi Markandeshwar deemed to be University, India

<sup>2</sup>Department of Biotechnology, Chandigarh Group of Colleges, Jhanjeri, Mohali, India

<sup>3</sup>Botany Section, School of Sciences, Maulana Azad National Urdu University, India

<sup>4</sup>Department of Field Crops, Faculty of Agriculture, Siirt University, Turkey

<sup>5</sup>Institute of Field and Vegetable Crops, National Institute of the Republic of Serbia, Serbia

<sup>6</sup>Institute for Multidisciplinary Research, University of Belgrade, Serbia

## Review Article

Volume 9 Issue 2

Received Date: June 13, 2024

Published Date: June 27, 2024

DOI: 10.23880/oajmb-6000299

\*Corresponding author: Manpreet Kaur, Department of Biotechnology, Chandigarh Group of Colleges, Jhanjeri, Mohali, India, Email: kaur64084@gmail.com

†Equally contributed towards this manuscript.

## Abstract

Endophytic fungus is an intriguing group of organisms that live inside their host's living tissues, which are primarily higher plants. Endophytes generate naturally occurring bioactive chemicals that are thought to operate as an elicitor for the synthesis of secondary metabolites in plants, without causing any disease symptoms in the host cells. By imparting stress tolerance and enhancing host resistance to a range of diseases, this incredibly varied collection of fungus can have a significant influence on plant ecosystems. They are known to improve nutrition through the reversible exchange of nutrients and protect plants from infections, which in turn affects development by releasing plant growth hormones. Endophytic fungus-infected plants exhibit notable increases in biomass, enhance commercial plant output, and are therefore beneficial to the agricultural industry. The potential uses of endophytic fungi as biological vectors, biological control agents, and insecticidal products, sources of secondary metabolites, antimicrobial agents, antitumor compounds, antibiotics, immunosuppressants, and antidiabetic agents make them of biotechnological interest. The current study concentrated on the numerous uses of fungal endophytes in several biotechnology fields.

**Keywords:** Endophytes; Secondary Metabolites; Plant Growth; Biofuel; Drug Discovery; Natural Products

**Abbreviations:** PGPE: Promote Plant Growth; IAA: Indole-3-acetic Acid; ROS: Reactive Oxygen Species.

## Introduction

Endophytic microorganisms, such as bacteria and fungi, are ubiquitous in all plant species and inhabit the

internal tissues of the plant without causing any apparent detrimental effects [1]. Endophytic microorganisms that promote plant growth (PGPE) reside within the tissues of plants, and their robust symbiotic relationship enhances the exchange of nutrients and enzymatic activity. Microbial endophytes can infiltrate plant tissues in a latent manner, thereby exhibiting no discernible symptoms [2]. Endophytic


fungi are a potential source of biocontrol agents by research that has been conducted on them [3]. These organisms spend part or all their life cycles living inside the otherwise healthy internal plant tissues of their hosts, yet they do not seem to do any noticeable damage to those tissues. They have a significant impact on the physiological processes that occur inside their host plants. By creating a diverse array of fungal metabolites, fungal endophytes improve their host's tolerance to abiotic stress, disease, insects, and mammalian herbivores [4].

Endophytes may either directly or indirectly activate these plant growth-promoting features in plants [5]. Endophytes are also known to have antimicrobial properties. These fungi frequently play a role in nutrient intake, boosting plant growth, improving plants' ability to withstand stress, limiting the growth of rivals and soil-borne pathogenic organisms, and improving disease resistance. Several techniques must be used in sustainable agriculture to improve food production while minimizing damage to the crop, the environment, and public health. An effective

substitute for conventional chemical farming is the use of fungal endophytes to enhance plant development. When crops are exposed to harmful biotic and abiotic stress factors including drought, cold, salinity, and disease, poisonous and lethal chemical pesticides and fertilizers, endophytic fungi help the plant to cope up with the stress [6]. An alternative to conventional chemical plant growth boosters is the use of endophytic fungi. Using mechanisms including triggered systemic resistance, bioremediation, and biological control, endophytic fungi help plants more effectively endure the impacts of abiotic stress [3].

### Applications of Fungal Endophytes

Beneficial secondary metabolites have been isolated from endophytic fungi, which have been used in biotechnological breakthroughs in the medicine, agricultural, nutrition, and environmental processing industries. Microorganisms called endophytes reside inside plant and are essential to their growth and well-being (Figure 1).



### Protection of the Host Plant Against Herbivorous Animals

There has been a significant amount of research and literature on the extremely specialised nature of the symbiotic relationship between plants and endophytes, as well as the impact that fungal alkaloids have on both vertebrate and invertebrate herbivores. When herbivores consume plant tissue, they are certain to come into contact with fungal metabolites if the host tissue has been pervasively and systemically colonised with endophyte hyphae. Herbivores may range from huge animals to tiny arthropods. Insect

resistance is increased by the majority of type C endophytes. Plants' resistance to insect herbivory is shown to depend on both the kind of fungal strain and the stage of growth, as determined by Rashid and Chung [7]. Evidence of the antinematode activity of class 1 endophytes has also been given by other investigations [8]. Although many endophytes are thought to confer resistance to insects and nematodes on their host plants, studies have shown that this is not always the case. These findings emphasise the need to investigate endophytes in their natural habitats among native plant species [9]. While C-endophytes have traditionally been regarded as beneficial mutualistic organisms, recent research

has revealed that a significant number of endophytes do not confer any protective advantages to their host plants [10]. Class IV endophytes, according to some studies, help the host plant defend itself from herbivores by secreting secondary compounds.

Plants with endophytes have been shown to be more resistant to fungal infection, and herbivores that nourish on infected plants have lower productivity. Endophytic fungi mitigate the negative effects of insect herbivores through a variety of mechanisms, such as diminished feeding, slowed development, stunted population growth and mortality [11]. Vertebrate herbivores exhibit similar avoidance and diminished performance patterns, including birds, rabbits, and deer. Endophyte infection can even lessen the impact of herbivores that live below ground, like nematodes and root-feeding insects. The strongest example for the anti-herbivore advantages of endophytic fungi comes from studies that indicate herbivore populations are destroyed when permitted to graze solely on plants infected with fungal endophytes [12].

Endophytic fungi create a wide variety of chemical defences, but not all of them are equally efficient, and many insect herbivores develop immunity to a specific substance at one point or another in their life cycles. As a general rule, larvae are more susceptible to poisons than adults. Geographic variation in consumption patterns means that even endophytic fungi that are supposed to be protective for their hosts, like *Neotyphidium* in plants, may not always cause avoidance or negative effects on herbivores [13].

### Influence of Plant's Ability to Withstand Stress

Plants are encountered with increasingly hostile environments for growth and maintenance as a result of environmental deterioration caused by agricultural operations and the changing climate. Equally difficult for a crop is that ever-increasing demands for agricultural output just further exacerbate the issue. Endophytic fungi are a potential solution for protecting plants from different stresses, and in this case, assistance is clearly required for healthy plant development. Endophytic fungi can protect host plants from water scarcity, extreme heat or cold, high or low salinity, and even poisonous metals [14]. Several dysfunctions in plant structure and function may be traced back to the altered genetic regulation of cellular pathways that results from exposure to abiotic stresses [15]. Endophytic fungus provides several benefits to host plants in a number of ways that increase their chances of success in the face of adversity. In response to oxidative stress, plants raise their catalase and peroxidase activity, which generates reactive oxygen species and initiates membrane damage and lipid peroxidation.

Endophytic fungi increase resistance to ROS and decrease lipid peroxidation [16]. Growth of the plants can be initiated directly by phytohormones, and indirect plant benefits can be achieved through the phytohormones' role in modulating the adaptation process to abiotic stresses. The hormone abscisic acid regulates stress-response gene expression and stomatal closure, both of which serve to decrease transpiration and so fluid loss. ABA is suppressed by endophytic fungi's presence [17]. Khan, et al. [18] investigated the effects of inoculating saline-stressed cucumber plants with the endophytic fungus *Paecilomyces* spp. Vegetative growth was higher in inoculation salinity-adapted plants. The favourable endophytic interaction of *Yarrowia lipolytica* was also reported by Jan, et al. [19], which reduced the negative effects of salt on maize plants. The effectiveness of the endophytic fungus *Porostereum spadiceum* in alleviating salt stress and enhancing the growth of soybean plants was observed. To achieve this, the phytohormone levels of both inoculated and control seedlings were compared [20]. Through regulating phytohormones, endophytic colonisation mitigated the negative effects of salt. The endophytic fungus *Aspergillus niger* was shown to give sunflower and soybean with resistance to high temperatures in another investigation. Plant length and biomass were all significantly raised by fungal inoculation [21]. Fungal endophytes produce bioactive metabolites and activate systemic resistance as the primary defence against pathogens, herbivores, and nematodes [22].

Because of its protective benefits on plants against biotic stresses, endophytic colonisation by a variety of fungi has been the centre of a significant amount of study. Several of these studies focused on *Serependita indica*, an endophyte that can protect many different types of crops from many different diseases [23]. Endophytic colonisation of cotton plants by *Phialemonium* sp., as described in a 2018 study by Zhou, et al. [24] inhibited root penetration by *Meloidogyne incognita* worms, gall development, and nematode reproduction. Endophytic colonization's impact on plant development and resistance to pests has also been investigated. One study by Dash, et al. [25] used an inoculation of endophytic fungi. The objective of the study was to investigate the impact of endophytic invasion on the fitness of the host plant and its resistance against *Tetranychus urticae*, commonly known as the two-spotted spider mite. To achieve this, seeds of *Phaseolus vulgaris* were subjected to inoculation with *Bacillus thuringiensis*, *Isaria fumosorosea*, and *Lecanicillium* sp. All strains examined by the scientists were able to successfully colonise bean plants as endophytes, and they were retrieved from the different parts of plants. The length of the plants and the fresh weight of the shoots and roots were both improved by the treatment of the seeds. Endophytic associations between *Penicillium* sp.

and *Phoma glomerata*, found in cucumber plants, resulted in considerably greater biomass and enhanced growth despite salt and water stress. The friendly relationship enhanced potassium, calcium, and magnesium intake while reducing sodium toxicity during saline stress. Reprogramming of plant growth in response to abiotic challenges by fungi via regulation of abscisic, jasmonic, and salicylic acid production has also been identified [26]. Studies by Jaber and Enkerli [27] demonstrated the ability of various strains of *B. bassiana* and *M. brunneum* to establish fungal colonies in broad bean plants, resulting in increased plant growth in seed-treated individuals. The same species were later reported by Jaber [28] as having successfully colonized the wheat (*T. aestivum*) plant's aerial parts and roots to stimulate plant development. Due of the impressive ability of the fungus to regenerate from both leaves and stems, suggested the capacity of systemic

colonization by *M. endophytica*, which was identified in 91% of the plants grown from fungal-inoculated seeds. These researches emphasized the possibility of endophytic interactions between several fungal species and their hosts. Research into the relationships between helpful microbes and their plant hosts has led to the discovery of plant-microbe interactions, opening up a new avenue in the quest for environmentally friendly farming practices [29].

### Production of Insecticidal Agents by Endophytes

Many endophytic fungi have anti-insect properties and are proved active against insects by producing some toxic insect repellent substances, some of which are given below in Table 1.

Sr. No.	Producing Fungus	Compound	References
1	<i>Geotrichium candidum</i>	Neofrapeptin A	Fredenhagen, et al. [30]
2	<i>Cordyceps heteropodac</i>	Cicadapeptin I	Krasnoff, et al. [31]
3	<i>Omphalotus olearius</i>	Omphalontins E-I	Liermann, et al. [32]
4	<i>Fusarium sp.</i>	Apicidin	Singh, et al. [33]
5	<i>Penicillium citrinum</i>	Quinolactide	Abe, et al. [34]
6	<i>Aspergillus niger</i>	Nafuredin- $\gamma$	Ômura, et al. [35]
7	<i>Beauveria bassiana</i>	Beauverolide N	Kuzma, et al. [36]

**Table 1:** Insecticidal compounds produced by endophytic fungi.


### Plant Growth Promotion

Endophytes, in addition to coping with host defence, are required to release compounds that are antibacterial and antifungal to maintain a healthy balance of antagonism with other competing organisms. The synthesis of phytohormones, which encourage the development of plants, is an additional

advantage of this process. Endophytes can indirectly or directly increase plant growth and yield (through biological nitrogen fixation, phytohormone production, siderophore synthesis, and phosphate solubilization), or indirectly through a variety of mechanisms (Table 2). Endophytes also have the potential to remove soil contaminants through the enhancement of phytoremediation (Figure 2).

Sr. no.	Activity Exhibited by Endophytic Fungi	References
1	IAA production	Bric, et al. [37]
2	Phosphate solubilization	Talukdar, et al. [38]
3	Cellulolytic activity	Dar, et al. [39]
4	Ammonia detection	Szilagyi-Zecchin, et al. [40]
5	HCN production	Donate-Correa, et al. [41]
6	Amylase production	Hankin, et al. [42]
7	Siderophore production	Schwyn, et al. [43]

**Table 2:** Different types of plant growth promoting activities exhibited by endophytic fungi.



**Figure 2:** Plant growth promotions by endophytic fungi in different ways [44].

Several fungal endophytes can stimulate growth in the host plant either directly or indirectly. This is achieved by processes such as habitat restriction and competition, direct antagonism of pathogens, and changes in the levels of plant hormones such as gibberellins and auxin in the host plant. The term “major nutrient” usually refers to this substance. It keeps plants in good overall health and vigour by doing things like encouraging the growth of roots, boosting the strength of stalks and stems, encouraging the production of flowers, improving crop quality, and boosting the resistance of plants to illnesses. Phosphorus is an important component of ATP. It is an essential component in passing on information about a plant’s genetic makeup from one generation to the next, acting as blueprint that details all aspects of a plant’s development as well as any changes to its DNA or RNA [45].

Because phosphorus is essential to the growth of seeds, it is abundant in seeds and fruits. Phosphorus is also present in high concentrations in fruits. Phytin is the primary type of phosphorus storage that is found in seeds. Phosphorus deficiency causes plants to mature later, produce lesser quality fodder, fruits, vegetables, and grains, and be more susceptible to disease; it slows the growth of the stem more than that of the roots and causes a purple hue to emerge in the leaves of some plants [46].

The organic form and the inorganic form of phosphorus are the two categories that may be found in the ground. It may be found in its organic form in decomposing plant matter, compost, and the tissues of microbes [47]. Plants are unable

to make use of phosphorus in an insoluble state because it cannot be dissolved in water [48]. Fungi are the organisms most capable of releasing vital nutrients from soil, according to a great number of studies. It was found that *Penicillium* and *Aspergillus* were two key species that were discovered with strong mineral-solubilizing activity, and this activity was associated to a lowering in the pH of culture medium.

Inorganic phosphate salts break down because of microorganisms, which leads to the release of organic acids [49,50]. Sustainable agriculture benefits from the wide variety of fungi that are known to reside in endophytic interactions with plants and are generally considered helpful to plant growth and development. *Penicillium* sp. was isolated from tea leaves, and their phosphate-solubilizing activity was investigated by Nath, et al. Both isolates that have demonstrated exceptionally high phosphate solubilization activity have contributed to a gradual elevation in the acidity of the culture medium up to day 8. These isolates may find widespread use as biofertilizers. Isolation of *Trichoderma gamsii* from *Lens esculenta* lateral root endophytes was confirmed by microscopic analysis, morphological analysis, and 18S rDNA sequencing. With a drop in pH, this fungus in the growth medium was able to solubilize up to 17% of the phosphate. *Trichoderma gamsii* demonstrated its capacity to promote plant development in a greenhouse bioassay utilizing four different test crops [51].

Research that was carried out by Nath, et al. [52] focuses on isolating fungal endophytes from tea (*Camellia sinensis*)

roots, stems, and leaves that were collected from tea gardens in Assam, India. These endophytes were then subjected to *in vitro* testing to determine their PGP activity levels. The fungus *Aspergillus niger* was shown to have the highest level of IAA activity, followed by the fungus *Penicillium sclerotiorum*. The highest activity was demonstrated by *Fusarium oxysporum* followed by *Penicillium chrysogenum* F1. *Penicillium sclerotiorum* is the most efficient phosphate solubilizer. Phytohormones are natural compounds that are released in certain organs of plants. These substances can be transported to other locations, where they stimulate specific reactions in terms of the plant's biology, physiology, and morphology. Aside from that, plant microbiomes synthesize growth regulators including gibberellic acid, auxins, and cytokinin [50]. The microbiomes of plants have been found to play a significant role in the development and proliferation of various plant organs, including but not limited to flowers, stems, leaves, and fruits.

Plant growth hormones can enhance and actually impact cell growth, development, and differentiation even at low concentrations. The following are the five major classes of phytohormones: abscisic acid (a) auxins, (b) cytokinins, (c) ethylene, (d) Gibberellins, (e) abscisic all are important plant growth regulators. According to the research conducted by Hamayun, et al. [53], certain fungal endophytes facilitate the growth of their host by stimulating the synthesis of gibberellins (GAs), indole-3-acetic acid (IAA), and cytokinins. The endophytic fungus, *Cladosporium sphaerospermum*, was identified in the roots of *Glycine max* (L.) and has been demonstrated to produce gibberellins. Rice was able to flourish when it was exposed to the culture filtrate, which had higher concentrations of the growth factors GA3, GA4, and GA7 respectively. The roots of *Calystegia soldanella* were examined, and a fungal endophyte called *Cadophora malorum* that produces GAs was discovered [54]. RSF-4L and RSF-6L are fungal endophyte species that have been identified as belonging to the genus *Fusarium* and a species of *Alternaria* were isolated from the leaves of the *Solanum nigrum* plant. According to the results of the Salkowski experiments, both *Fusarium* sp. & *Alternaria* sp. produced IAA, respectively. After being treated with fungal CFs, plant growth characteristics were enhanced.

The two categories of fungal endophytes play a crucial role in facilitating the growth and development of plants, as noted by Khan, et al. [55]. The presence of fungal endophytes significantly enhances a plant's capacity to endure biotic and abiotic stresses. To aid in relationships with plant hosts and to provide resistance to pests and diseases, endophytic fungi produce bioactive compounds. These bonds are mediated and protected by bioactive chemicals. There is enough proof to demonstrate that endophytic fungi interact with their

hosts for mutual advantages, including the improvement of host health, defence against pests, grazers, and rhizosphere nematodes, and improvement in drought tolerance and root development. In exchange, endophytic fungi acquire habitation and diet.

Endophytes are microorganisms that synthesise bioactive compounds, enabling plants to defend themselves and providing them with the requisite energy to do so. Endophytic microorganisms create mycotoxin, which defends the host plant against attack by pathogenic microbes caused by the plant's natural predators [56]. The 18S rRNA gene sequencing shows that the strain belonged to the genus *Penicillium* and produced gibberellins with the GA5 designation [53]. Fungal extracellular enzymes are finding more and more applications in fields as diverse as biotechnology, medicines, the food and beverage industries, leather production, agriculture, bioremediation, and chemical compound biotransformation [57,58].

In addition to their role as phytoremediators, endophytes perform the functions of main saprobic decomposers, mutualists, and latent pathogens. There is a wide variety of endophytes that can perform the function of phytoremediation. The majority of these endophytes are heavy metal-resistant endophytes, both hyper-accumulators and non-hyperaccumulators and organic contaminant-degrading endophytes [59,60]. Bioremediation using these endophytes increases plant growth (via regulation of phytohormones [61], enzyme production, and nitrogen fixation), decreases phytotoxicity (via production of iron chelators, siderophores, and enzymes that degrade iron), and ultimately increases plant metal tolerance [62].

### Antimicrobial and Antidiabetic Activity

Endophytes are capable of producing several hydrolytic enzymes in addition to peptide and polyene macrolide antibiotics like amphotericin B and nystatin [63]. Penicillin and cephalosporins are, without a doubt, the most significant anti-infective medications now available on the market. Endophytic fungi produce a number of antibiotics, including Daptomycin [64] and Valinomycin, a strong antibiotic that is effective against acute respiratory syndrome Coronavirus [65,66]. An endophytic fungus *Cryptosporiopsis quercina* is renowned for its active antifungal action against *C. albicans* and other species of Trichophyton. These fungi are considered to be human fungal infections. A molecule known as cryptocandin [66], which was discovered from *C. quercina*, is renowned for its efficacy as an antimycotic drug. A compound called cryptocin was isolated from the same fungus and has been demonstrated to effectively combat *Pyricularia oryzae* [67].

Pseudomyces, also known as lipopeptides, have been shown to be active against human pathogenic fungi such as *Candida albicans* and *Cryptococcus neoformans* [68,69]. In order to develop NMR approaches, ambuic acid, which is generated by *Pestalotiopsis microspore*, in conjunction with another endophyte called terrein, is employed [70]. A molecule called pestalosite [71], which was isolated from *P. microspore*, has been shown to have antimicrobial characteristics, but the two sesquiterpenes known as pestalotiopsins A and B are known to have phytotoxic effects. *Muscodora albus*, in addition to endophytic fungus, is responsible for the production of volatile chemicals that have antimicrobial and antifungal properties [72]. The isoamyl acetate that is generated by this fungus is the most useful substance that it produces. There is evidence that some species of *Gliocladium* generate the volatile antibiotic chemical annulene [73]. Some bioactive agents produced by the genus *Xylaria*, such as sordaricin and multiplolides A and B, have been shown to have biological activity against *C. albicans* [74]. On the other hand, mellisol and 1,8-dihydroxynaphthol 1-O- $\alpha$ -glucopyranoside have been shown to have activity against HSV type 1 [75]. An endophytic *Streptomyces* species that was isolated from *Grevillea pteridifolia* generates a new antibiotic known as 'Kakadumycin A,' which is both an effective antibacterial agent and an antimalarial agent [76].

The antimicrobial spectrum of *Phomopsis sp.*, which was isolated from the host plant *Rhizophora apiculata*, was more effective against gram-positive bacteria [77]. *D. zingiberensis* culture filtrates inhibited *Bacillus*, *Staphylococcus* and *E. Coli* [78]. It was discovered that the fungal endophytes found in *Eucalyptus exserta* have strong antibacterial effects when tested as raw extracts [79].

### Anticancer Compounds Produced by Endophytes

It is possible for endophytes to produce secondary metabolic products that are identical to or equivalent to those generated by their hosts (Table 3). Anticancer agents are examples of the bioactive substances that are created simultaneously by plants and the endophytes that are linked with them. Endophytic fungi have been shown to be a rich and consistent source of naturally occurring chemicals that have anticancer properties. These natural products have the potential to have a significant effect on the development of anticancer medications in contemporary medicine [80]. Endophyte-derived natural compounds have the potential to be employed as an alternative source in the research that ultimately leads to the creation of innovative treatment options for cancer [81-83]. Many endophytes have been identified as the sources of new chemicals that have shown promising results in anticancer testing.

Sr. No.	Host plant	Fungal Endophyte	Anticancer Agent	References
1	<i>Taxus brevifolia</i>	<i>Taxomyces andreanae</i>	Taxol (Palcitaxel)	Stierle, et al. [84]
2	<i>Camptotheca acuminata</i>	<i>Fusarium solani</i>	Camphothecin	Ran, et al. [85]
3	<i>Podophyllum. Hexandrum</i>	<i>Tramete shirsuta</i>	Podophyllotoxin	Giri, et al. [86]
4	<i>Adenophoreaxiliflora</i>	<i>Chaetomium sp.</i>	Chaetominine	Jiao, et al. [87]
5	<i>Torreya. Taxifolia</i>	<i>Pestalotiopsis microspore</i>	Cytochalasins Torreyanic acid	Sappapan, et al. [88]

**Table 3:** Anticancer agents produced by fungal endophytes.

### Bioactive Substances Produced by Endophytic Fungi

Interaction of plant with the endophytic fungus naturally results in the biosynthesis of secondary metabolites (Table 4). Genes from both the plant and the endophyte species regulate and modulate the nature of the connection between them [89]. Endophytic fungi in medicinal plants produce an abundance of secondary metabolism products [88]. Secondary metabolite production in plants may also be influenced by the symbiotic interaction between endophytes

and their host plants. Plants developing in adverse environments must be tested for endophyte isolation and its metabolites [88]. Alkaloids, benzopyranones, chinones, flavonoids, phenolic acids, and many others are only some of the beneficial secondary metabolites that endophytes produce [90]. Such bioactive metabolites are commonly used as agrochemicals, antibiotics, antiparasitics and antioxidants [88]. Biotechnological methods that employ certain organisms show promise as viable substitutes for developing an infinite, low-cost, and sustainable source of high-quality bioactive products as well as aromatic compounds.

Bioactive Compound	Plant Species	Endophyte	References
Podophyllotoxin and its analogues	<i>Sinopodophyllum hexandrum</i> ,	<i>Alternaria</i> sp.	Giri, et al. [86]
	<i>Diphylleia sinensis</i>	<i>Penicillium</i> sp.	
	<i>Dysosmaveitchii</i>	<i>Monilia</i> sp.	
Camptothecin and its analogues	<i>Camptotheca acuminata</i> ,	<i>Fusarium solani</i>	Kusari, et al. [91]
	<i>Nothapodytes foetida</i>	<i>Botryosphaeria parva</i>	
Huperzine A	<i>Huperzia serrata</i>	<i>Acremonium</i>	Kusari, et al. [91]
	<i>Lycopodium serratum</i>	<i>Penicillium chrysogenum</i>	
Vinblastine and its analogues	<i>Catharanthus roseus</i>	<i>Alternaria</i> sp.	Huang, et al. [92]
Cytoskyrin a	<i>Conocarpus erecta</i>	<i>Cytospora</i> sp.	Brady, et al. [73]
Phomoxanthone a	<i>Lucila spinosa</i>	<i>Xylariasp.</i>	Isaka, et al. [93]
Phomoxanthone b	<i>Tectona grandis</i>	<i>Phomosissp.</i>	
Rubrofusarin b	<i>Cynodondactylon</i>	<i>Aspergillus niger</i>	Guo, et al. [94]
Emindole DA	<i>Mediterranean green alga</i>	<i>Emericellanidulans</i>	Kralj, et al. [95]
Paclitaxel and its analogues	<i>Taxus cuspidata</i>	<i>Alternaria</i> sp.	Wani, et al. [95]

**Table 4:** Bioactive compounds produced by Endophytic Fungi.

### Production of Antioxidants

ROS, or reactive oxygen species, are by-products of the aerobic process found in all living organisms that serve a number of functions. The production of free radicals in the form of various ROS is triggered when the light absorption capacity of photosynthetic cells exceeds their usage [97,98]. This happens because oxygen molecules can quickly accept electrons from the body's electron transport system [99]. Herbivorous and pathogenic plant stress causes the production of ROS via oxidative bursts in the host plant [100,101]. During these bursts, chemicals are synthesised that may be employed to either block the spread of invading hyphae physically or to have direct, damaging effects on the cells of these organisms. Many studies [102,103] corroborate this notion. This study's results provide credence to the hypothesis that reactive oxygen species serve a useful role in preparing the plant's response to abiotic stresses, which has been proposed after a thorough reevaluation of the role of ROS in the abiotic stress response of plants [99,104]. The plant produces many singlet oxygen species as part of this function, and these species travel extensive distances inside the tissue of the plant to provide systemic communication throughout the plant.

An arsenal of antioxidants is produced during the activation of the plant's stress response. These antioxidants then regulate the degree of ROS build-up in plant cells, hence lowering the risk of cell damage and apoptosis [101]. ROS have a mechanistic role in apoptosis, nonspecific physiological stress, and systemic signalling, all of which

influence the effectiveness of fungal infection or endophytic fungi colonisation and the plant's reactions, which may be resistance, acceptance, or sanctions. In addition, via the interaction between oxidants and antioxidants, antioxidants may play a role in the transmission of stress signals. As a result, the host may be better able to respond rapidly to pathogenesis and tell the difference between a mutualist and a pathogen through chemical communication with an avirulent pathogen or an asymptomatic endophyte.

When cultures of *Phyllosticta* sp. were subjected to ROS, Srinivasan, et al. found that the organisms exhibited a high level of antioxidant activity. The interaction between fungal endophytes and host plants is a dynamic process that involves the potential involvement of reactive oxygen species (ROS) and antioxidants in the modulation of the host's hypersensitivity and systemic developed resistance mechanisms.

### Biofuel Production

Researchers looked at the possibility of producing lipid biodiesel precursors using endophytic fungus that had been identified from a variety of different tropical plant species. Widespread interest is already being shown in the production of next-generation fuels using endophytic fungus as a hitherto identified reservoir of low - molecular weight hydrocarbons and lipids. These microbes will be directly responsible for the production of these forthcoming biofuels, which will result in a simpler and shorter biosynthetic process. In addition, they will be well suited for instant use in




current combustion engines without requiring fundamental modifications to vehicle structure. Endophytes that have optimal lipid profiles have been reported to have been isolated by research organisations operating in different parts of the globe. These endophytes may be used as precursors in the production of biofuels. Researchers have discovered that some types of fungal endophytes may create volatile organic molecules, which can then be used to make traditional diesel fuel. The genomic sequence of one of these endophytic fungi that produces biofuel was only published not too long ago. Understanding of the biosynthetic mechanism by which endophytes are able to accomplish the difficult task of producing volatile hydrocarbons will be improved thanks to the discovery of *Ascocorynesarcooides*. Research in this vitally essential subject of alternative bio-energy for the purpose of ensuring India's energy supply has gradually begun to acquire pace. Fungal endophytes live inside plants and are used in the production of biodiesel. The plants *Jatropha curcas*, *Pongamia pinnata*, *Sapindusmukorossi*, *Mesuaferrea*, *Terminalia bellerica*, *Casabelathevetia*, and *Ricinus communis* are some of the plants that contain these fungi [105].

### Discovery of New Drugs

Natural substances have been shown in recent research to have potential in the pharmaceutical industry as a source of innovative and diversified pharmacological templates for future drug development and discovery [106-109]. Research into endophytes for natural products is expected to play a significant role in the future of drug discovery and development since it is becoming increasingly clear that many natural compounds result from microbial associations with their hosts [110-111].

### Natural Products Derived from Fungi Endophytes

Endophytes of fungi are well-established sources of natural compounds and produce particular plant-derived metabolites that are beneficial from a pharmaceutical standpoint. Many kinds of plant-derived medicinal chemicals (Figure 3), which are generated by fungal endophytes, are discussed in Table 5.



**Figure 3:** Natural products derived from fungi endophytes.

Endophytic Fungus	Plant Source	Plant Derived Compound	Activity	References
<b>Alkaloids</b>				
<i>Cladosporium cladosporioides</i>	<i>Aconitum</i> spp.	Aconitine	Anticancer	Yang, et al. [112]
<i>Alternaria alternata</i>	<i>Capsicum annuum</i>	Capsaicin	Anti-inflammatory, gastro-stimulatory	Devari, et al. [113]
<i>Fusarium oxysporum</i> ,	<i>Amoora rohituka</i> <i>Dysoxylum binectariferum</i>	Rohitukine	Anticancer, CDK inhibitor	Kumara, et al. [114]
<b>Coumarins</b>				
<i>Botrydia plodiatheobromae</i>	<i>Citrus bergamia</i> , <i>Grapefruit peel</i>	Bergapten, Meranzin	Antioxidant	Zaher, et al. [115]
<i>Annulohyphoxylona bovei</i> var. <i>microspora</i>	<i>Acanthopanax senticosus</i> , <i>Sarcandra glabra</i>	Isofraxidin	Anticancer	Yamazaki, et al. [116]; Cheng, et al. [117]
<i>Penicillium</i> sp., <i>Xylaria</i> sp.	<i>Alibertia macrophylla</i>	Mellein	Antibacterial, antifungal	Oliveira, et al. [118]
<i>Penicillium</i> sp.	<i>Artemisia scoparia</i> , <i>Scopolia carniolica</i>	Scopoletin, Umbelliferone	Antifungal, antioxidant, anti-inflammatory	Huang, et al. [92]
<b>Flavanoids</b>				
<i>Colletotrichum</i> sp	<i>Cajanus cajan</i>	Apigenin	Antibacterial, anticancer, antioxidant,	Shukla, et al. [119]
<i>Hypocrealixii</i>	<i>Cajanus cajan</i>	Cajanol	Anticancer, antimicrobial	Zhao, et al. [120]
<i>Curcuma wenyujin</i>	<i>Curcuma</i> spp	Curcumin	Anti-inflammatory, antioxidant, antitumor	Wang, et al. [121]
<b>Lignans</b>				
Members of <i>xylariaceae</i>	<i>Angelica archangelica</i>	Coniferin	Antidiabetic	Chapela, et al. [122]
<i>Alternaria neesex</i>	<i>Forsythia suspensa</i>	Podo-phyllotoxin	Antitumor, antiviral	Kong, et al. [123]
<b>Saponins</b>				
<i>Penicillium oxalicum</i>	<i>Gymnema sylvestre</i>	Gymnemagenin	Anti-diabetic	Parthasarathy, et al. [124]
<i>Aspergillus</i> sp.	<i>Panax</i>	Ginsenoside	antioxidation, antitumor	Wu, et al. [125]
<b>Terpenes</b>				
<i>Penicillium</i>	<i>Azadirachta indica</i>	Azadirachtin	Insecticidal	Kusari, et al. [91]
<i>Nodulisporium</i> sp.	<i>Cinnamomum camphora</i>	Camphor	Antimicrobial, topical skin preparations	Suwannarach, et al. [126]
<b>Quinones and Xanthenes</b>				
<i>Chaetomium globosum</i>	<i>Hypericum perforatum</i>	Hypericin	Anti-depressant, antimicrobial	Kusari, et al. [91]
<i>Gibberella moniliformis</i>	<i>Lawsonia inermis</i>	Lawsonone	Cytotoxic	Sarang, et al. [127]
<b>Miscellaneous Plant-derived Compounds</b>				
<i>Aspergillus unguis</i>	<i>Wheat, rye, barley</i>	Azelaic acid	Antimicrobial, anticancer	Kamat, et al. [128]
<i>Muscodor vitigenus</i>	<i>Ancistrocladus tectorius</i>	Naphthalene	Antibacterial, insect repellent	Ruangrungsi, et al. [129]

**Table 5:** A list of plant-derived compounds from fungal endophytes.

## Production of Immunosuppressive Compounds

Immunosuppressive drugs are offered to patients undergoing organ transplantation in order to suppress, minimise, or avoid allograft rejection. Therefore, they are very important in medical management of autoimmune diseases like SLE etc. Because of the rising demand for immunosuppressive treatments, there is a pressing need to accelerate the search for pharmaceuticals that are not only safer but also more trustworthy. This is necessary in order to investigate the issues that are now plaguing the efficacy of these medications. Many investigations have shown that endophytes are capable of producing bioactive chemicals that may suppress the immune system [65,130]. Chemical analysis is often employed to identify the presence of (-) mycosanine enamine. The endophyte *Mycosphaerella* sp. was isolated from the china root leaves plant to produce this biomolecule [131]. Cyclosporin A and (-) mycosanine enamine were able to decrease the proliferation of T lymphocytes by reducing the expression of CD25 and CD69 surface activation antigens. This resulted in the inhibition of T cell proliferation [131]. In conjunction to this, nine polyketides were isolated from the mangrove leaves that harbour the endophytic fungus *Penicillium* sp. ZJ-SY2. These compounds showed promising immunosuppressive properties [130]. *Fomitopsisbetulinus*, a basidiomycete often linked with wood decay, was the source of *Xylarialongipes* HFG1018. This fungus was responsible for the production of eighteen new nor-isopimarane diterpenes, which were designated as xylarinorditerpenes A–R (1–18). Some of these compounds had the ability to suppress the immune system [132].

## Production of Siderophores

Siderophores are created by many species with the ability to scavenge iron from the environment they are in to render the essential substance easily accessible to the cell. Various species are capable of producing siderophores. To facilitate uptake and utilisation of ferric iron, siderophores are secreted out into the environment to form soluble ferric complexes. Siderophores have very complicated chemical structures, which enables them to create the most powerful iron-chelating complexes. Siderophores are engaged in the process of virulence and play an important role in the mobilisation of iron and other components. Recent research has also shed light on the significant connection between siderophores and the capacity to withstand oxidative stress. Both in agricultural and medical contexts, their applications have been the subject of much research. On the other hand, new research areas are beginning to focus on the

use of siderophores as green-iron chelators. Specifically, siderophores are being considered as a potential solution for the protection of cultural assets.

Many microorganisms that live on or around plants, both beneficial and harmful, share a common trait: the synthesis of extracellular siderophores. Siderophores are produced by a number of different strains of fungus. Scots pine and Labrador tea foliar endophytic fungi were studied for their ability to produce antimicrobial and antioxidant compounds via extracellular siderophore production. In vitro production of the siderophore ferricrocin resulted in concentrations ranging from 7.9 to 17.6 micrograms per litre. Ferricrocin was only created by the fungus that also had antibacterial action, and none of the well-known siderophores were found in the broths made by the fungi that produced antioxidants. As a result, the synthesis of ferricrocin is characteristic of certain foliar endophytic fungus, but not all of them. The discovery of ferricrocin in the leaves of Labrador tea lends credence to the hypothesis that this compound may play a function in vivo in the dynamic relationship that exists between the endophyte and the plant host [133].

## Conclusion

New and exciting avenues for applied research into plant-microbe interactions can be found by studying the function of fungal endophytes, as these microbes can boost germination rates, enhance seedling health, and increase plant growth. The production of extracellular enzymes, phytohormones, and secondary metabolites can be linked to these abilities. Because of their potential to be used as a biofertilizer to improve plant growth, we should encourage further research into these microbes. An enormous portion of the endophytic population in terrestrial plants is probably still undiscovered. Metagenomic research into plant endophyte populations will reveal new applications and details about these microorganisms, such as the genera, phenotypic traits, and potential roles in seed germination and plant development. There is a need for more study to determine the genetic determinants involved in plant growth promotion and the molecular mechanisms by which plant-endophyte interactions induce defence resistance mechanisms against different types of stresses.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

1. Grabka R, d'Entremont TW, Adams SJ, Walker AK, Tanney JB, et al. (2022) Fungal Endophytes and Their Role in Agricultural Plant Protection against Pests and Pathogens. *Plants (Basel)* 11(3): 384.
2. Hassan SED (2017) Plant growth-promoting activities for bacterial and fungal endophytes isolated from medicinal plant of *Teucrium polium* L. *Journal of Advanced Research* 8: 687-695.
3. Fontana DC, Paula SD, Torres AG, Souza VHMD, Pascholati SF, et al. (2021) Endophytic Fungi: Biological Control and Induced Resistance to Phytopathogens and Abiotic Stresses. *Pathogens* 10(5): 570.
4. Wen J, Okyere SK, Wang S, Wang J, Xie L, et al. (2022) Endophytic Fungi: An Effective Alternative Source of Plant-Derived Bioactive Compounds for Pharmacological Studies. *J Fungi (Basel)* 8(2): 205.
5. Baron NC, Rigobelo EC (2021) Endophytic fungi: a tool for plant growth promotion and sustainable agriculture. *Mycology* 13(1): 39-55.
6. Chaudhary P, Agri U, Chaudhary A, Kumar A, Kumar G (2022) Endophytes and their potential in biotic stress management and crop production. *Front Microbiol* 13: 933017.
7. Rashid MdHO, Chung YR (2017) Induction of Systemic Resistance against Insect Herbivores in Plants by Beneficial Soil Microbes. *Front Plant Sci* 8: 1816.
8. Strobel G, Daisy B (2003) Bioprospecting for Microbial Endophytes and Their Natural Products. *Microbiol Mol Biol Rev* 67(4): 491-502.
9. Saikkonen K, Wali P, Helander M, Faeth SH (2004) Evolution of endophyte-Plant symbioses. *Trends in Plant Science* 9(6): 275-280.
10. Hawas UW, El-Beih AA, El-Halawany AM (2012) Bioactive anthraquinones from endophytic fungus *Aspergillus versicolor* isolated from red sea algae. *Arch Pharm Res* 35(10): 1749-1756.
11. Samal I, Bhoi TK, Majhi PK, Murmu S, Pradhan AK, et al. (2023) Combatting insects mediated biotic stress through plant associated endophytic entomopathogenic fungi in horticultural crops. *Front Plant Sci* 13: 1098673.
12. Bamisile BS, Dash CK, Akutse KS, Keppanan R, Wang L (2018) Fungal Endophytes: Beyond Herbivore Management. *Front Microbiol* 9: 544.
13. Koh S, Hik DS (2007) Herbivory Mediates Grass-Endophyte Relationships. *Ecology* 88(11): 2752-2757.
14. Zhang J, Lu J, Zhu Y, Shen X, Zhu B, et al. (2024) Roles of endophytic fungi in medicinal plant abiotic stress response and TCM quality development. *Chinese Herbal Medicines* 16(2): 204-213.
15. Egamberdieva D, Wirth SJ, Alqarawi AA, Allah EFA, Hashem A (2017) Phytohormones and Beneficial Microbes: Essential Components for Plants to Balance Stress and Fitness. *Front Microbiol* 8: 2104.
16. Yan L, Zhu J, Zhao X, Shi J, Jiang C, et al. (2019) Beneficial effects of endophytic fungi colonization on plants. *Appl Microbiol Biotechnol* 103(8): 3327-3340.
17. Khan AL, Hussain J, Al-Harrasi A, Al-Rawahi A, Lee IJ (2015) Endophytic fungi: resource for gibberellins and crop abiotic stress resistance. *Crit Rev Biotechnol* 35(1): 62-74.
18. Khan AL, Hamayun M, Kang SM, Kim YH, Jung HY, et al. (2012) Endophytic fungal association via gibberellins and indole acetic acid can improve plant growth under abiotic stress: an example of *Paecilomyces formosus* LHL10. *BMC Microbiol* 12: 3.
19. Jan FG, Hamayun M, Hussain A, Jan G, Iqbal A, et al. (2019) An endophytic isolate of the fungus *Yarrowia lipolytica* produces metabolites that ameliorate the negative impact of salt stress on the physiology of maize. *BMC Microbiol* 19(1): 3.
20. Hamayun M, Hussain A, Khan SA, Kim HY, Khan AL, et al. (2017) Gibberellins Producing Endophytic Fungus *Porostereum spadiceum* AGH786 Rescues Growth of Salt Affected Soybean. *Front Microbiol* 8: 686.
21. Hamayun MI, Hussain A, Iqbal A, Khan SA, Lee IJ (2020) *Aspergillus niger* boosted heat stress tolerance in sunflower and soybean via regulating their metabolic and antioxidant system. *J Plant Interactions* 15(1): 223-232.
22. Poveda J, Abril-Urias P, Escobar C (2020) Biological Control of Plant-Parasitic Nematodes by Filamentous Fungi Inducers of Resistance: Trichoderma, Mycorrhizal and Endophytic Fungi. *Front Microbiol* 11: 992.
23. Lugtenberg BJJ, Caradus JR, Johnson LJ (2016) Fungal endophytes for sustainable crop production. *FEMS Microbiol Ecol* 92(12): fiw194.
24. Zhou W, Wheeler TA, Starr JL, Valencia CU, Sword GA (2018) A fungal endophyte defensive symbiosis affects plant-nematode interactions in cotton. *Plant Soil* 422:

- 251-266.
25. Dash CK, Bamisile BS, Keppanan R, Qasim M, Lin Y, et al. (2018) Endophytic entomopathogenic fungi enhance the growth of *Phaseolus vulgaris* L. (Fabaceae) and negatively affect the development and reproduction of *Tetranychus urticae* Koch (Acari: Tetranychidae). *Microb Pathog* 125: 385-392.
  26. Wang Y, Mostafa S, Zeng W, Jin B (2021) Function and Mechanism of Jasmonic Acid in Plant Responses to Abiotic and Biotic Stresses. *Int J Mol Sci* 22(16): 8568.
  27. Jaber LR, Enkerli J (2017) Fungal entomopathogens as endophytes: can they promote plant growth? *Biocontrol Science and Technology* 27(1): 28-41.
  28. Jaber LR (2018) Seed inoculation with endophytic fungal entomopathogens promotes plant growth and reduces crown and root rot (CRR) caused by *Fusarium culmorum* in wheat. *Planta* 248(6): 1525-1535.
  29. Card S, Johnson L, Teasdale S, Caradus J (2016) Deciphering endophyte behaviour: the link between endophyte biology and efficacious biological control agents. *FEMS Microbiol Ecol* 92(8): fiw 114.
  30. Fredenhagen A, Molleyres LP, Böhlendorf B, Laue G (2006) Structure Determination of Neofrapeptins A to N: Peptides with Insecticidal Activity Produced by the Fungus *Geotrichum candidum*. *J Antibiot (Tokyo)* 59(5): 267-280.
  31. Krasnoff SB, Keresztes I, Gillilan RE, Szebenyi DME, Donzelli BGG, et al. (2007) Serinocyclins A and B, Cyclic Heptapeptides from *Metarhizium anisopliae*. *J Nat Prod* 70(12): 1919-1924.
  32. Liermann JC, Opatz T, Kolshorn H, Antelo L, Hof C, et al. (2009) Omphalotins E-I, Five Oxidatively Modified Nematicidal Cyclopeptides from *Omphalotus olearius*. *Eur J Org Chem* 2009(8): 1256-1262.
  33. Singh SB, Zink DL, Liesch JM, Dombrowski AW, Darkin-Rattray SJ, et al. (2001) Structure, Histone Deacetylase, and Antiprotozoal Activities of Apicidins B and C, Congeners of Apicidin with Proline and Valine Substitutions. *Org Lett* 3(18): 2815-2818.
  34. Abe M, Imai T, Ishii N, Usui M (2006) Synthesis of Quinolactamide *via* an Acyl Migration Reaction and Dehydrogenation with Manganese Dioxide, and Its Insecticidal Activities. *Biosci Biotechnol Biochem* 70(1): 303-306.
  35. Ōmura S, Miyadera H, Ui H, Shiomi K, Yamaguchi Y, et al. (2001) An anthelmintic compound, nafuredin, shows selective inhibition of complex I in helminth mitochondria. *Proc Natl Acad Sci USA* 98(1): 60-62.
  36. Kuzma M, Jegorov A, Kacer P, Havlicek V (2001) Sequencing of new beauverolides by high-performance liquid chromatography and mass spectrometry. *J Mass Spectrom* 36(10): 1108-1115.
  37. Bric JM, Bostock RM, Silverstone SE (1991) Rapid in Situ Assay for Indoleacetic Acid Production by Bacteria Immobilized on a Nitrocellulose Membrane. *Appl Environ Microbiol* 57(2): 535-538.
  38. Talukdar R, Tayung K (2019) Antimicrobial Activity of Endophytic Fungi Isolated from *Eryngium foetidum*, an Ethnomedicinal Plant of Assam. *IJPSSDR* 11(6).
  39. Dar RA, Saba I, Shahnawaz M, Sangale MK, Ade AB, et al. (2013) Isolation, purification and characterization of carboxymethyl cellulase (CMCase) from endophytic *Fusarium oxysporum* producing podophyllotoxin. *AER* 1(4): 91-96.
  40. Szilagyi-Zecchin VJ, Ikeda AC, Hungria M, Adamoski D, Kava-Cordeiro V, et al. (2014) Identification and characterization of endophytic bacteria from corn (*Zea mays* L.) roots with biotechnological potential in agriculture. *AMB Expr* 4: 26.
  41. Donate-Correa J, León-Barrios M, Pérez-Galdona R (2004) Screening for plant growth-promoting rhizobacteria in *Chamaecytisus proliferus* (tagasaste), a forage tree-shrub legume endemic to the Canary Islands. *Plant and Soil* 266(1/2): 261-272.
  42. Hankin L, Anagnostakis SL (1975) The Use of Solid Media for Detection of Enzyme Production by Fungi. *Mycologia* 67(3): 597-607.
  43. Schwyn B, Neilands JB (1987) Universal chemical assay for the detection and determination of siderophores. *Anal Biochem* 160(1): 47-56.
  44. Carvalho JOD, Broll V, Martinelli AHS, Lopes FC (2020) Endophytic fungi: positive association with plants. In: Sharma V, et al. (Eds.), *Molecular Aspects of Plant Beneficial Microbes in Agriculture*. Elsevier, pp: 321-332.
  45. Nyoki D, Ndakidemi PA (2014) Effects of Bradyrhizobium japonicum Inoculation and Supplementation with Phosphorus on Macronutrients Uptake in Cowpea (*Vigna unguiculata* (L.) Walp). *AJPS* 5(4): 442-451.
  46. Rodríguez H, Fraga R (1999) Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol Adv* 17(4-5): 319-339.

47. Silva AGD (2017) Forms of soil organic phosphorus at black earth sites in the Eastern Amazon. *Revista Ciência Agronômica* 48(1).
48. Priyadharsini P, Muthukumar T (2017) The root endophytic fungus *Curvularia geniculata* from *Parthenium hysterophorus* roots improves plant growth through phosphate solubilization and phytohormone production. *Fungal Ecology* 27(Part A): 69-77.
49. Wakelin SA, Warren RA, Harvey PR, Ryder MH (2004) Phosphate solubilization by *Penicillium* spp. closely associated with wheat roots. *Biology and Fertility of Soils* 40: 36-43.
50. Yadav AN, Sharma D, Gulati S, Singh S, Dey R, et al. (2015) Haloarchaea Endowed with Phosphorus Solubilization Attribute Implicated in Phosphorus Cycle. *Sci Rep* 5: 12293.
51. Rinu K, Sati P, Pandey A (2014) *Trichoderma gamsii* (NFCCI 2177): A newly isolated endophytic, psychrotolerant, plant growth promoting, and antagonistic fungal strain. *J Basic Microbiol* 54(5): 408-417.
52. Nath R, Sharma GD, Barooah M (2015) Plant Growth Promoting Endophytic Fungi Isolated From Tea (*Camellia Sinensis*) Shrubs Of Assam, India. *AEER* 13(3): 877-891.
53. Khan SA, Hamayun M, Yoon H, Kim HY, Suh SJ, et al. (2008) Plant growth promotion and *Penicillium citrinum*. *BMC Microbiol* 8: 231.
54. You YH, Kwak TW, Kang SM, Lee MC, Kim JG (2015) *Aspergillus clavatus* Y2H0002 as a New Endophytic Fungal Strain Producing Gibberellins Isolated from *Nymphoides peltata* in Fresh Water. *Mycobiology* 43(1): 87-91.
55. Khan AR, Ullah I, Waqas M, Shahzad R, Hong SJ, et al. (2015) Plant growth-promoting potential of endophytic fungi isolated from *Solanum nigrum* leaves. *World J Microbiol Biotechnol* 31(9): 1461-1466.
56. Carroll G (1988) Fungal Endophytes in Stems and Leaves: From Latent Pathogen to Mutualistic Symbiont. *Ecology* 69(1): 2-9.
57. Yadav C (2018) New and Future Developments in Microbial Biotechnology and Bioengineering: penicillium system properties and applications. Elsevier, Amsterdam.
58. Benjamin S, Pandey A (1998) *Candida rugosa* lipases: Molecular biology and versatility in biotechnology. *Yeast* 14(12): 1069-1087.
59. Rosa GDI, Peralta-Videa JR, Montes M, Parsons JG, Cano-Aguilera I, et al. (2004) Cadmium uptake and translocation in tumbleweed (*Salsola kali*), a potential Cd-hyperaccumulator desert plant species: ICP/OES and XAS studies. *Chemosphere* 55(9): 1159-1168.
60. Soleimani M, Afyuni M, Hajabbasi MA, Nourbakhsh F, Sabzalian MR, et al. (2010) Phytoremediation of an aged petroleum contaminated soil using endophyte infected and non-infected grasses. *Chemosphere* 81(9): 1084-1090.
61. Zhang W, Niu Y, Li YX, Zhang F, Zeng RJ (2020) Enrichment of hydrogen-oxidizing bacteria with nitrate recovery as biofertilizers in the mixed culture. *Bioresour Technol* 313: 123645.
62. Ma Y, Rajkumar M, Luo YM, Freitas H (2011) Inoculation of endophytic bacteria on host and non-host plants—Effects on plant growth and Ni uptake. *J Hazard Mater* 195: 230-237.
63. Martin JF, McDaniel LE (1977) Production of Polyene Macrolide Antibiotics. *Adv Appl Microbiol* 21: 1-52.
64. Kirkpatrick P, Raja A, LaBonte J, Lebbos J (2003) Daptomycin. *Nat Rev Drug Discov* 2(12): 943-944.
65. Wang WX, Cheng GG, Li ZH, Ai HL, He J, et al. (2019) Curtachalasin, immunosuppressive agents from the endophytic fungus *Xylaria cf. curta*. *Org Biomol Chem* 17(34): 7985-7994.
66. Strobel GA, Miller RV, Martinez-Miller C, Condrón MM, Teplow DB, et al. (1999) Cryptocandin, a potent antimycotic from the endophytic fungus *Cryptosporiopsis cf. quercina*. *Microbiology (Reading)* 145(Pt 8): 1919-1926.
67. Li JY, Harper JK, Grant DM, Tombe BO, Bashyal B, et al. (2001) Ambuic acid, a highly functionalized cyclohexenone with antifungal activity from *Pestalotiopsis* spp. and *Monochaetia* sp. *Phytochemistry* 56(5): 463-468.
68. Ballio A, Barra D, Bossa F, Collina A, Grgurina I, et al. (1991) Syringopeptins, new phytotoxic lipodepsipeptides of *Pseudomonas syringae* pv. *syringae*. *FEBS Lett* 291(1): 109-112.
69. Harrison L, Teplow DB, Rinaldi M, Strobel G (1991) Pseudomycins, a family of novel peptides from *Pseudomonas syringae* possessing broad-spectrum antifungal activity. *J Gen Microbiol* 137(12): 2857-2865.
70. Harper JK, Mulgrew AE, Li JY, Barich DH, Strobel GA, et al. (2001) Characterization of Stereochemistry and

- Molecular Conformation Using Solid-State NMR Tensors. *J Am Chem Soc* 123(40): 9837-9842.
71. Lee JC, Yang X, Schwartz M, Strobel G, Clardy J (1995) The relationship between an endangered North American tree and an endophytic fungus. *Chem Biol* 2(11): 721-727.
  72. Strobel GA, Dirkse E, Sears J, Markworth C (2001) Volatile antimicrobials from *Muscodora albus*, a novel endophytic fungus. *Microbiology (Reading)* 147(Pt 11): 2943-2950.
  73. Brady SF, Wagenaar MM, Singh MP, Janso JE, Clardy J (2000) The Cytosporones, New Octaketide Antibiotics Isolated from an Endophytic Fungus. *Org Lett* 2(25): 4043-4046.
  74. Pongcharoen W, Rukachaisirikul V, Phongpaichit S, Kühn T, Pelzing M, et al. (2008) Metabolites from the endophytic fungus *Xylaria* sp. PSU-D14. *Phytochemistry* 69(9): 1900-1902.
  75. Pittayakhajonwut P, Suvannakad R, Thienhirun S, Prabpai S, Kongsaree P, et al. (2005) An anti-herpes simplex virus-type 1 agent from *Xylaria mellisii* (BCC 1005). *Tetrahedron Letters* 46(8): 1341-1344.
  76. Li E, Jiang L, Guo L, Zhang H, Che Y (2008) Pestalochlorides A-C, antifungal metabolites from the plant endophytic fungus *Pestalotiopsis adusta*. *Bioorg Med Chem* 16(17): 7894-7899.
  77. Buatong J, Phongpaichit S, Rukachaisirikul V, Sakayaroj J (2011) Antimicrobial activity of crude extracts from mangrove fungal endophytes. *World J Microbiol Biotechnol* 27: 3005-3008.
  78. Xu L, Zhou L, Zhao J, Li J, Li X, et al. (2008) Fungal endophytes from *Dioscorea zingiberensis* rhizomes and their antibacterial activity. *Lett Appl Microbiol* 46(1): 68-72.
  79. Mao Z, Zhang W, Wu C, Feng H, Peng Y, et al. (2021) Diversity and antibacterial activity of fungal endophytes from *Eucalyptus exserta*. *BMC Microbiol* 21(1): 155.
  80. Cragg GM, Grothaus PG, Newman DJ (2009) Impact of Natural Products on Developing New Anti-Cancer Agents. *Chem Rev* 109(7): 3012-3043.
  81. Joseph B, Priya RM (2011) Bioactive Compounds from Endophytes and their Potential in Pharmaceutical Effect: A Review. *American J of Biochemistry and Molecular Biology* 1(3): 291-309.
  82. Alvin A, Miller KI, Neilan BA (2014) Exploring the potential of endophytes from medicinal plants as sources of antimycobacterial compounds. *Microbiol Res* 169(7-8): 483-495.
  83. Xie S, Ogden A, Aneja R, Zhou J (2016) Microtubule-Binding Proteins as Promising Biomarkers of Paclitaxel Sensitivity in Cancer Chemotherapy. *Med Res Rev* 36(2): 300-312.
  84. Kharwar RN, Mishra A, Gond SK, Stierle A, Stierle D (2011) Anticancer compounds derived from fungal endophytes: their importance and future challenges. *Nat Prod Rep* 28(7): 1208-1228.
  85. Ran X, Zhang G, Li S, Wang J (2017) Characterization and antitumor activity of camptothecin from endophytic fungus *Fusarium solani* isolated from *Camptotheca acuminata*. *Afr Health Sci* 17(2): 566-574.
  86. Giri A, Narasu ML (2000) Production of podophyllotoxin from *Podophyllum hexandrum*: a potential natural product for clinically useful anticancer drugs. *Cytotechnology* 34(1-2): 17-26.
  87. Jiao RH, Xu S, Liu JY, Ge HM, Ding H, et al. (2006) Chaetominine, a Cytotoxic Alkaloid Produced by Endophytic *Chaetomium* sp. IFB-E015. *Org Lett* 8(25): 5709-5712.
  88. Sappapan R, Sommit D, Ngamrojanavanich N, Pengpreecha S, Wiyakrutta S, et al. (2008) 11-Hydroxymonocerin from the Plant Endophytic Fungus *Exserohilum rostratum*. *J Nat Prod* 71(9): 1657-1659.
  89. Zuccaro A, Lahrman U, Langen G (2014) Broad compatibility in fungal root symbioses. *Curr Opin Plant Biol* 20: 135-145.
  90. Tan RX, Zou WX (2001) Endophytes: a rich source of functional metabolites. *Nat Prod Rep* 18(4): 448-459.
  91. Kusari S, Lamshöft M, Zühlke S, Spiteller M (2008) An Endophytic Fungus from *Hypericum perforatum* that Produces Hypericin. *J Nat Prod* 71(2): 159-162.
  92. Huang Z, Guo Z, Yang R, Yin X, Li X, et al. (2009) Chemistry and cytotoxic activities of polyketides produced by the mangrove endophytic fungus *Phomopsis* SP. ZSU-H76. *Chem Nat Compd* 45: 625-628.
  93. Isaka M, Jaturapat A, Rukseree K, Danwisetkanjana K, Tanticharoen M, et al. (2001) Phomoxanones A and B, Novel Xanthone Dimers from the Endophytic Fungus *Phomopsis* Species. *J Nat Prod* 64(8): 1015-1018.
  94. Guo B, Wang Y, Sun X, Tang K (2008) Bioactive natural products from endophytes: a review. *Prikl Biokhim Mikrobiol* 44(2): 153-158.

95. Kralj A, Kehraus S, Krick A, Eguereva E, Kelter G, et al. (2006) Arugosins G and H: Prenylated Polyketides from the Marine-Derived Fungus *Emericella nidulans* var. *a cristata*. *J Nat Prod* 69(7): 995-1000.
96. Wani MC, Taylor HL, Wall ME, Coggon P, McPhail AT (1971) Plant antitumor agents. VI. Isolation and structure of taxol, a novel antileukemic and antitumor agent from *Taxus brevifolia*. *J Am Chem Soc* 93(9): 2325-2327.
97. Logan DC (2007) The mitochondrial compartment. *J Exp Bot* 58(1): 1225-1243.
98. Gill SS, Tuteja N (2010) Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol Biochem* 48(12): 909-930.
99. Foyer CH, Noctor G (2000) Tansley Review No. 112: Oxygen processing in photosynthesis: regulation and signalling. *New Phytologist* 146(3): 359-388.
100. Apel K, Hirt H (2004) Reactive Oxygen Species: Metabolism, Oxidative Stress, and Signal Transduction. *Annu Rev Plant Biol* 55: 373-399.
101. Jaspers P, Kangasjärvi J (2010) Reactive oxygen species in abiotic stress signaling. *Physiol Plant* 138(4): 405-413.
102. Overmyer K, Brosché M, Kangasjärvi J (2003) Reactive oxygen species and hormonal control of cell death. *Trends Plant Sci* 8(7): 335-342.
103. Gara LD, Locato V, Dipierro S, Pinto MCD (2010) Redox homeostasis in plants. The challenge of living with endogenous oxygen production. *Respir Physiol Neurobiol* 173: S13-S19.
104. Foyer CH, Noctor G (2011) Ascorbate and Glutathione: The Heart of the Redox Hub. *Plant Physiol* 155(1): 2-18.
105. Paul S, Bhagobaty RK, Nihalani MC, Joshi SR (2020) Characterization of oleaginous endophytic fungi of biodiesel plants as potential biofuel minifactories. *Biomass and Bioenergy* 142: 105750.
106. Shen B (2015) A New Golden Age of Natural Products Drug Discovery. *Cell* 163(6): 1297-1300.
107. Newsham KK (2011) A meta-analysis of plant responses to dark septate root endophytes. *New Phytol* 190(3): 783-793.
108. Calixto JB (2019) The role of natural products in modern drug discovery. *An Acad Bras Ciênc* 91(Suppl 3): e20190105.
109. Wilson BAP, Thornburg CC, Henrich CJ, Grkovic T, O'Keefe BR (2020) Creating and screening natural product libraries. *Nat Prod Rep* 37(7): 893-918.
110. Strobel G (2018) The Emergence of Endophytic Microbes and Their Biological Promise. *J Fungi (Basel)* 4(2): 57.
111. Abdel-Razek AS, El-Naggar ME, Allam A, Morsy OM, Othman SI (2020) Microbial Natural Products in Drug Discovery. *Processes* 8(4): 470.
112. Wu H, Yang HY, You XL, Li YH (2013) Diversity of endophytic fungi from roots of *Panax ginseng* and their saponin yield capacities. *SpringerPlus* 2(1): 107.
113. Yang K, Liang J, Li Q, Kong X, Chen R, et al. (2013) *Cladosporium cladosporioides* XJ-AC03, an aconitine-producing endophytic fungus isolated from *Aconitum leucostomum*. *World J Microbiol Biotechnol* 29(5): 933-938.
114. Devari S, Jaglan S, Kumar M, Deshidi R, Guru S, et al. (2014) Capsaicin production by *Alternaria alternata*, an endophytic fungus from *Capsicum annum*; LC-ESI-MS/MS analysis. *Phytochemistry* 98: 183-189.
115. Kumara PM, Soujanya KN, Ravikanth G, Vasudeva R, Ganeshiah KN, et al. (2014) Rohitukine, a chromone alkaloid and a precursor of flavopiridol, is produced by endophytic fungi isolated from *Dysoxylum binectariferum* Hook.f and *Amoora rohituka* (Roxb). *Wight & Arn. Phytomedicine* 21(4): 541-546.
116. Zaher AM, Moharram AM, Davis R, Panizzi P, Makboul MA, et al. (2015) Characterisation of the metabolites of an antibacterial endophyte *Botryodiplodia theobromae* Pat. of *Dracaena draco* L. by LC-MS/MS. *Nat Prod Res* 29(24): 2275-2281.
117. Yamazaki T, Tokiwa T (2010) Isofraxidin, a Coumarin Component from *Acanthopanax senticosus*, Inhibits Matrix Metalloproteinase-7 Expression and Cell Invasion of Human Hepatoma Cells. *Bio Pharm Bull* 33(10): 1716-1722.
118. Cheng C, Li W, Lin M, Yang ST (2019) Metabolic engineering of *Clostridium carboxidivorans* for enhanced ethanol and butanol production from syngas and glucose. *Bioresour Technol* 284: 415-423.
119. Oliveira CM, Regasini LO, Silva GH, Pfenning LH, Young MCM, et al. (2011) Dihydroisocoumarins produced by *Xylaria* sp. and *Penicillium* sp., endophytic fungi associated with *Piper aduncum* and *Alibertia*



- macrophylla. *Phytochemistry Letters* 4(2): 93-96.
120. Shukla S, Gupta S (2010) Apigenin: A Promising Molecule for Cancer Prevention. *Pharm Res* 27(6): 962-978.
  121. Zhao Y, Zhao T, Jia H, Li X, Zhu Z, et al. (2017) Optimization of the composition of mixed entrainer for economic extractive distillation process in view of the separation of tetrahydrofuran/ethanol/water ternary azeotrope. Optimization of the composition of mixed entrainer. *J Chem Technol Biotechnol* 92(9): 2433-2444.
  122. Wang X (2020) Groundless Research Published on the International Journal of Plant Sciences. *VP* 6(4): 167-169.
  123. Chapela IH, Petrini O, Hagemann L (1991) Monolignol glucosides as specific recognition messengers in fungus-plant symbioses. *Physiological and Molecular Plant Pathology* 39(4): 289-298.
  124. Kong P, Zhang L, Guo Y, Lu Y, Lin D (2014) Phillyrin, a Natural Lignan, Attenuates Tumor Necrosis Factor  $\alpha$ -Mediated Insulin Resistance and Lipolytic Acceleration in 3T3-L1 Adipocytes. *Planta Med* 80(11): 880-886.
  125. Parthasarathy R, Sathiyabama M (2014) Gymnemagenin-producing Endophytic Fungus Isolated from a Medicinal Plant *Gymnema sylvestris* R.Br. *Appl Biochem Biotechnol* 172(6): 3141-3152.
  126. Suwannarach N, Kumla J, Bussaban B, Nuangmek W, Matsui K, et al. (2013) Biofumigation with the endophytic fungus *Nodulisporium* spp. CMU-UPE34 to control postharvest decay of citrus fruit. *Crop Protection* 45: 63-70.
  127. Sarang H, Rajani P, Vasanthakumari MM, Kumara PM, Siva R, et al. (2017) An endophytic fungus, *Gibberella moniliformis* from *Lawsonia inermis* L. produces lawsone, an orange-red pigment. *Antonie van Leeuwenhoek* 110(7): 853-862.
  128. Kamat S, Kumari M, Taritla S, Jayabaskaran C (2020) Endophytic Fungi of Marine Alga From Konkan Coast, India—A Rich Source of Bioactive Material. *Front Mar Sci* 7: 31.
  129. Ruangrunsi N, Wongpanich V, Tantivatana P, Cowe HJ, Cox PJ, et al. (1985) Traditional Medicinal Plants of Thailand, V. Ancistrocladine, a New Naphthalene-Isoquinoline Alkaloid from *Ancistrocladus tectorius*. *J Nat Prod* 48(4): 529-535.
  130. Liu H, Chen S, Liu W, Liu Y, Huang X, et al. (2016) Polyketides with Immunosuppressive Activities from Mangrove Endophytic Fungus *Penicillium* sp. *ZJ-SY2*. *Mar Drugs* 14(12): 217.
  131. Wang LW, Wang JL, Chen J, Chen JJ, Shen JW, et al. (2017) A Novel Derivative of (-)-mycosamine Produced by the Endophytic Fungus *Mycosphaerella nawae*, Exhibits High and Selective Immunosuppressive Activity on T Cells. *Front Microbiol* 8: 1251.
  132. Chen HP, Zhao ZZ, Cheng GG, Zhao K, Han KY, et al. (2020) Immunosuppressive Nor-isopimarane Diterpenes from Cultures of the Fungicolous Fungus *Xylaria longipes* HFG1018. *J Nat Prod* 83(2): 401-412.
  133. Kajula M, Tejesvi MV, Kolehmainen S, Mäkinen A, Hokkanen J, et al. (2010) The siderophore ferricrocin produced by specific foliar endophytic fungi in vitro. *Fungal Biol* 114(2-3): 248-254.