



Visual Light Leak in High-Frequency Electromagnetic Radiation Hole: A Photon-Electron Theory for Living Tissue for Explaining Myopia

Lee A*

Individual Researcher, China

*Corresponding author: Alex Lee, Xianning West Road 28#, Xi'an city, Shanxi province, China, 710,049; Email: alexlee2019@outlook.com

Hypothesis

Volume 7 Issue 1

Received Date: May 17, 2022

Published Date: June 20, 2022

DOI: 10.23880/oajo-16000243

Abstract

If the frequency of the photon exceeds the threshold, the photon has enough energy to strike the electron away and forms a hole in living tissue include the eye. Longer time of high-frequency electromagnetic radiation will cause more holes and a larger radius of holes. Some visual light leak through the hole and cannot be refracted on the macula, this is myopia symptom of blurred vision, shortening the distance of the eye and the object will include more visual light into cornea and lens, this is myopia another symptom of nearsightedness. The longer eye axis is caused by strike of high energy photon. Blue light and X-ray from screen are two kinds of high-frequency electromagnetic radiation which can shot hole in eye and cause myopia. Healing the shot hole can cure myopia.

Keywords: Myopia; Causes and Processes; Photon-Electron Theory; Screen Radiation; Shot Hole; X-ray; Blue Light; Eye Axis Elongation; Healing Hole; Curing Myopia

Introduction

Myopia is a worldwide visual impairment [1]. Myopia has increased from 25% to 44% between 1972 and 2004 in the United States [1]. In the cities of Asia, the prevalence is over 80% 2021/10/14. There will be 2.62 billion people in myopia by the year 2020, and about 4.76 billion people (49.8% of the world population) will become myopia by the end of 2050 [2]. From 350 BC to now, literature has showed hypotheses about etiology of myopia such as lack of enough outdoor activities and too long time of near work [1], literature suggested that blue light from screen may correlate with myopia [3,4], but the identical effect of blue light remains unclear [3-5]. The specific causes of myopia remain undefined [1,6,7].

A Hypothesis of Particle Property of the High-Frequency Electromagnetic Radiation in Living Tissue

Albert Einstein proposed that light is not a wave passing through space, but a collection of independent wave packets (photons), which has an energy $h\nu$ [8]. In 1923 the "Compton scattering" showed the particle concept of electromagnetic radiation [9]. In 2017, an experiment tested the records of light-by-light scattering at the Large Hadron Collider [10]. The electromagnetic radiation effect on cornea has been investigated on the wave property but not the particle property [11,12]. The effects of ionizing radiation in living tissue are controversial [13,14].

The First Hypothesis

The photon has the particle-like property in living tissue.

A Hypothesis of Photon-Electron Scattering in Living Tissue

According to photon-electron theory, when the photon's frequency reaches or exceeds a threshold, electrons are kicked out only by the hit of photons; this differs from the classical electromagnetic theory [8].

The Second Hypothesis

- a. The photon of visual light's frequency is lower than the threshold, which cannot hit the electron away from its original position; the direction of visual light is changed by the scattering of electron (Figures 1a & 1b).
- b. The photon of high-frequency electromagnetic radiation has enough frequency to hit the electrons away and shot in a straight line in living tissue (Figures 1c & 1d).

A Hypothesis of Massive Photons of High-Frequency Electromagnetic Radiation Producing Holes in the Eye

It has reported the holes produced by heavy-ion radiation. In 1983, literature showed a membrane lesion in the heavy ion irradiated corneas [15], later in 2004, the Brookhaven National Laboratory in the US made an experiment on cells irradiation by heavy ions, which showed that holes may form as the result of heavy ion irradiation in liposomes and that these holes are large enough to allow leakage of cell internal contents [16].

There is literature reported the hole results of electromagnetic radiation. In 1946, literature reported an experiment result on the animal that the Cathode Ray has a penetration without underlying reaction [17]. In 1983, an experiment showed the hole is the physical damage to the lens after microwave radiation on the murine ocular lens [18]. In 2016, literature reported a corneal hole during radiation therapy for cancers of the head and neck [19].

The Third Hypothesis

- a. The visual light can be focused on the macula by the scattering of electron without hole left (Figures 1e & 1f), because the frequency of the visual light is lower than the threshold, when the photon of the visual light strikes the electron in eye, the direction of the light is changed without kicking out the electron from its original position.

- b. The high-frequency electromagnetic radiation can produce holes in living tissue include the eye. Because the photon of the high-frequency electromagnetic radiation has enough frequency to scatter the electron away from the original position in living tissue, massive high-frequency photons can scatter massive electrons away from their original positions in the living tissue, which leads to holes in living tissue (Figures 1g & 1h). The diameter of the hole irradiated by electromagnetic radiation is in the photon's diameter scale.
- c. If the visual light gets into the radiation hole, there are two situations, the first situation is the visual light touches the cornea and lens, this photon is refracted on the macula by the cornea and lens, because the crystalline can scatter the photon of visual light independently [20], most light is refracted in the cornea, the exact focusing is made by the soft lens with changing into different shapes [11,20] (Figure 1i); Another situation is the visual light passes through the hole without touching the cornea and lens, the focusing does not happen, the visual light goes through the radiated hole in a straight line, the macula cannot get the visual light, the image of this visual light is lost in the macula (Figure1j).

Myopia

- a. To illustrate the principle of this design, the writer ignores the pupil, vitreous body, etc. The visual light refracted to retina forms a reversed image, this is omitted. The writer omits eye's other characteristics in this article.
- b. To show the assumption, the scale of photon and photon radiated hole in the figures is not the real scale.
- c. To compare the visual acuity in a healthy eye and in an eye with radiated hole, it is proposed that in a time, three visual light photons are in the incidence angles of α , β , γ , in the following discussions, these three visual light photons are in the fixed incidence angles of α , β , γ as the example visual light photons (Figure 1k).
- d. Taking the distance of S for example, the S is the furthest vision distance for the three chosen visual light photons for the health eye. We can see from (Figure 1k), the visual light photon in the incidence angle of γ will not include into the eye when the distance is further than S.
- e. The three visual light photons fly into the eye in the distance of S, the healthy eye can focus all the three visual light photons to the macula lutea [21] (Figure 1k).
- f. The three visual light photons fly into the eye in the distance of S, the eye with radiated hole loses the photon of the incidence angle of β , the macula gets two photons but not three photons, this means the visual acuity of the example three visual light photons decreased (Figure1l). The vision in the eye with radiated hole is not distinct as in the healthy eye in distance of S, because the macula gets three photons in (Figure 1k), the macula gets two

photons in (Figure 1l). This is one of myopia symptoms: blurred vision in distance.

- g. When the distance of the eye and the object is in Y which is closer than the S , the photon of the incidence angle of β will get into the cornea and lens, the refraction happens

on the photon of the visual light of the incidence angle of β , the macula gets three photons (Figure 1m), the vision is distinct as in (Figure 1k), this is another myopia symptoms: nearsightedness or distinct vision in close distance.




Figure 1: Principle of myopia and the modification of myopia.

- a) Incident photon of visual light with the incidence angle of δ .
- b) The photon of visual light is scattered by the electron, the refraction angle is ϵ .
- c) Incident photon of high-frequency electromagnetic radiation with the incidence Angle of δ .
- d) The photon of high-frequency electromagnetic radiation strikes out the electron from the original position, the refraction does not happen, the photon goes in straight line.
- e) The incident visual light is refracted by cornea and lens on the macula.
- f) No hole left in eye after refraction of visual light.
- g) The high-frequency electromagnetic radiation strikes in a straight line in eye without refraction.
- h) The high-frequency electromagnetic radiation strikes out a hole in eye.
- i) The visual light is refracted on the macula in the radiated hole while it touches the cornea and lens.
- j) The visual light is not refracted on the macula in the radiated hole while it does not touch the cornea and lens.
- k) Three visual light photons in incidence angles of: α , β , γ are all refracted on the macula in healthy eye in the distance of S .
- l) Two photons of visual light (incidence angles of α , γ) are refracted on the macula, one photon of visual light (incidence angle of β) goes straightly through the radiated hole in the distance of S .
- m) All the three photons of visual light are refracted on the macula when shortening the distance ($Y < S$) in the eye with radiated hole.
- n) New high-frequency electromagnetic radiation holes appear in the eye with radiated hole after radiated by more high-frequency electromagnetic radiation.
- o) All the three photons of visual light are not refracted on the macula in the eye with the new high-frequency electromagnetic radiation holes when the distance is S .
- p) All the three photons of visual light are refracted on

the macula in the eye with the new high-frequency electromagnetic radiation holes when shortening the distance ($Z < Y < S$).

- q) In a distance of S , two photons of visual light are scattered by the concave lens on the cornea and lens.

Development of Myopia

1. From Figures 1h to 1n, the eye is radiated by more high-frequency electromagnetic radiation, more radiated holes appear in the eye, they are called the new radiated holes in the following discussions (Figure 1n), which is different with the eye with radiated hole in (Figure 1h).
2. In the distance of further than S , from Figure 1o we can see, the eye with the new radiated holes still cannot focus all the three visual light photons on the macula, because the incidence angles are different which results in leakage of visual light photon in cornea and lens. That means in a further distance than S , the eye with the new radiated holes cannot have a clear vision as a healthy eye shown in Figure 1k. This is myopia symptom: if the distance was further than S , the eye with the new radiated holes still cannot see clearly as a healthy eye.
3. In the distance of S , in the eye with the new radiated holes, the three visual light photons pass through the new radiated holes without touching the cornea and lens, the refraction does not happen, the visual lights go in the straight lines. The macula cannot get the visual light photon, the eye loses the image of these three visual light photons (Figure 1o). This is one of myopia symptoms: blurred vision in distance.
4. In the distance of between Z ($Z < Y < S$) and S , the different incidence angles will cause the visual light photons excluded in cornea and lens in the eye with the new radiated holes. The macula will lose visual light photons because they pass through the new radiated holes without refraction (Figure 1p). This is one of myopia symptoms: blurred vision in distance.
5. In the distance of Z , the distance of Z is the furthest distance for the eye with the new radiated holes to include the three visual light photons to cornea and lens in the example, three visual light photons are focused on the macula in the eye with the new radiated holes by cornea and lens in the distance of Z (Figure 1p), this is another myopia symptom: nearsightedness or distinct vision in close distance.

Concluded from 1-5 above, in the distance of further than Z , the eye with the new radiated holes cannot focus all the three visual light photons on the macula, in the distance of Z and in the distance of the shorter than Z , the eye with the new radiation holes can focus all the three visual light photons on the macula, these are myopia symptom: blurred in distance, only having distinct vision in close distance.

Open Access Journal of Ophthalmology

Comparing Figures 1l to 1o, we can find out that the eye with the new radiated holes cannot focus all the three visual light photons on the macula in the distance of S (Figure 1o), but the eye with radiated hole can focus two visual light photons on the macula in the distance of S (Figure 1l), this is one symptom of the development of myopia: fuzziness is more serious from Figure 1l to Figure 1o in the distance of S .

Comparing Figures 1m and 1p, we can find out that the furthest distance for including all the three visual light photons on the cornea and lens for the eye with the new radiated holes is Z (Figure 1p), which is shorter than the furthest distance (Y) of including all the three visual light photons on the cornea and lens for the eye with the radiated hole (Figure 1m), this is another symptom of the development of myopia: nearsightedness is more serious from (Figure 1m to 1p).

Conclusion from Figures 1a to 1p more radiation of high-frequency radiation, more holes in eye, higher myopia.

The correction of myopia with wearing the concave lens

Myopia glass is a concave lens which can scatter the photon, using a concave lens can help cornea and lens collecting more photons comparing with a non-concave lens in the far distance of S for an eye with holes irradiated by high-frequency electromagnetic radiation, the principle is shown in (Figure 1q). This is why myopic people can see clearly in far distance by wearing myopia glass.

Eye Elongation

Myopic eyes are longer compares with healthy eyes [22], which caused by reduced collagen synthesis and increased collagen degradation [4]. But according to this manuscript, massive high-frequency photons radiate into the eye and hit electrons of the eye, if the radiation incidence angle has a horizontal component which produces a propulsive force in the axial length direction, the eye elongated by the shooting of high-frequency photons.

The Radiated Hole and the Macular Hole

- a) The most common macular change in highly myopic eyes is macular holes [23-28]. In this article, larger radius of the radiated hole and more radiated holes correlate with the higher myopia, because the high-frequency electromagnetic radiation can pass through the eye ball, the high-frequency electromagnetic radiation penetrates not only the cornea and lens but also the macula, according to the photon-electron theory in living tissue in this article, there will appear some radiated holes in

the macula, which is in accordance with the investigation results.

- b) The macular hole is a result of high myopia [4], but according to this article, the hole in the eye causes myopia, longer time of striking of photon of high-frequency electromagnetic radiation causes larger radius of holes and causes higher degree of myopia.
- c) The macular hole is in micron scale [23-28], the author proposes the radiated hole is in scale of photon scale.

Blue Light

The blue light from computer, TV, phone or tablet is think to be no harmful to eye [29], but according to this model, the author proposes the blue light from computer, TV, phone or tablet is one of high-frequency electromagnetic radiation, which can strike the lens and cornea to form a hole. The blue light is one cause of myopia.

X-Ray

Although the Flat-Panel Display have replaced the Cathode Ray Tube Television Sets, Video Displays and Computer Monitors for many years, but literature reports that there were about 232 million Cathode Ray Tube Monitors and Television sets in the United States by 2013, which may be reused [30]. There will be about 0.27 to 5.86 million Cathode Ray Tube Television in Peru by 2025 [30]. In Color Cathode Ray Tube Computer Monitors and Color Cathode Ray Tube Television Set, three guns are used to shoot the electrons to phosphor spots for producing red, green and blue [31], during the shooting the electrons colliding with the shadow mask, the phosphor layer, and the face panel glass produce photons in the X-ray region of the electromagnetic spectrum called Bremsstrahlung X-rays [13,14,32], The mean energy of photons reaching the operator working in front of the monitor was above 17 keV [13]. An experiment showed that the annual equivalent dose of X-ray radiation of Cathode Ray Tube Computer Monitors and Cathode Ray Tube Television Set is about 0.1 mSv/y higher than the background radiation [14], the author supposes the X-ray is one of the high-frequency electromagnetic radiation which causes myopia.

Verification of this Model

- a) From experiment, we can find out that the blue light and X-ray can strike the lens and cornea to form a hole in the radius of photon diameter scale.
- b) In this kind of myopic eyeball, we will find out many holes which have a radius in the photon's radius scale.
- c) Larger radius of the hole and more holes correlate with the higher myopia, according to this article, the hole in the eye causes myopia, longer time of striking of photon of high-frequency electromagnetic radiation

Open Access Journal of Ophthalmology

causes larger radius of holes and causes higher degree of myopia.

- d) Testing myopic person's living or working environment, we will find out high-frequency electromagnetic radiation instrument.

Prevention of Myopia

Decreasing near work time and increasing the outdoor activity treated as the way of prevention of myopia in children [4], but according to this article, prevention of myopia is in two ways:

- a. Using special glass which can avoid the eye from the high-frequency electromagnetic radiation but can let visual light pass through, for example, the lead glass can decrease X-ray beam radiation but let visual light pass [13,33,34].
- b. Using special material which can block the machine from emitting high-frequency electromagnetic radiation.

Curing Myopia

When a person was shot by a bullet, we always need to cure the shot hole, the same as the high-frequency electromagnetic radiation we need to cure the hole in the eye to cure myopia.

Acknowledgements

The author made this research alone and did not use funding source in this research.

Consent Statement/Ethical Approval

Not required.

Conflict of Interest Statement

No

References

1. Cooper J, Tkatchenko AV (2018) A Review of Current Concepts of the Etiology and Treatment of Myopia. *Eye & Contact Lens* 44(4): 231-247.
2. Liu SX, Ye S, Wang QF, Cao YJ, Zhang X (2018) Breastfeeding and myopia: A cross-sectional study of children aged 6-12 years in Tianjin, China. *Scientific Reports* 8(1): 10025.
3. Zhao ZC, Zhou Y, Tan G, Li J (2018) Research progress about the effect and prevention of blue light on eyes. *International Journal of Ophthalmology* 11(12): 1999-

- 2003.
4. Wu PC, Huang HM, Yu HJ, Fang PC, Chen CT (2016) Epidemiology of Myopia. *Asia-Pacific Journal of Ophthalmology* 5(6): 386-393.
 5. Zhang T, Zuo Y, Wei Y, Huang W, Zhou X, et al. (2018) The Prevalence and Associations of Peripheral Retinopathy: Baseline Study of Guangzhou Office Computer Workers. *J Ophthalmol*.
 6. de Jong PTVM (2018) Myopia: its historical contexts. *British Journal of Ophthalmology* 102(8): 1021-1027.
 7. Warner N (2016) Update on myopia. *Current Opinion in Ophthalmology* 27(5): 402-406.
 8. Einstein A (1917) Quantum theory of radiation. *Physikalische Zeitschrift* 18: 121-128.
 9. Compton AHJPR (1923) A quantum theory of the scattering of x-ray by light elements. *Phys Rev* 21: 483-502.
 10. Collaboration A (2017) Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC. *Nat Phys* 13: 852-858.
 11. Meek KM, Knupp C (2015) Corneal structure and transparency. *Prog Retin Eye Res* 49: 1-16.
 12. Meek KM, Quantock AJ (2001) The Use of X-ray Scattering Techniques to Determine Corneal Ultrastructure. *Prog Retin Eye Res* 20(1): 95-137.
 13. Khaledi N, Arbabi A, Dabaghi M (2015) X-Ray Dose Estimation from Cathode Ray Tube Monitors by Monte Carlo Calculation. *Health Phys* 108(4): 401-406.
 14. Perez-Vega C, Zamanillo JM, Ipina JS (2000) Assessment of ionizing radiation from PC monitors and TV receivers. *IEEE Trans Consum Electron* 46(4): 1048-1051.
 15. Nelson AC, Tobias CA (1983) Rapid development of corneal lesions in rats produced by heavy ions. *Adv Space Res* 3(8): 195-209.
 16. Koniarek JP, Thomas JL, Vazquez M (2004) In Space Life Sciences: Radiation Risk Assessment and Radiation Measurement in Low Earth Orbit. *Advances in Space Research-Series* 34: 1373-1377.
 17. Robbins LL, AUB JC (1946) Superficial Burns of Skin and Eyes from Scattered Cathode Rays. *Radiology* 46: 1-23.
 18. Stewart-DeHaan PJ, Creighton MO, Larsen LE, Jacobi JH, Ross WM, et al. (1983) In vitro studies of microwave-induced cataract: Separation of field and heating effects. *Exp Eye Res* 36(1): 75-90.
 19. Elghazi T, Omor Y, Hafidi Z, Eljai A, Elmoize Z, et al. (2016) Unusual complication of radiation therapy, corneal perforation: about a case. *Pan Afr Med J* 25: 64.
 20. Bloemendal H, de Jong W, Jaenicke R, Lubsen NH, Slingsby C, et al. (2004) Ageing and vision: structure, stability and function of lens crystallins. *Prog Biophys Mol Biol* 86(3): 407-485.
 21. Strauss O (2005) The retinal pigment epithelium in visual function. *Physiol Rev* 85(3): 845-881.
 22. Atchison DA, Jones CE, Schmid KL, Pritchard N, Pope JM, et al. (2004) Eye Shape in Emmetropia and Myopia. *Invest Ophthalmol Vis Sci* 45(10): 3380-3386.
 23. Kadonosono K, Yazama F, Itoh N, Uchio E, Nakamura S, et al. (2001) Treatment of retinal detachment resulting from myopic macular hole with internal limiting membrane removal. *Am J Ophthalmol* 131(2): 203-207.
 24. Lai TT, Yang CM (2018) Lamellar Hole-Associated Epiretinal Proliferation in Lamellar Macular Hole and Full-Thickness Macular Hole in High Myopia. *Retina* 38(7): 1316-1323.
 25. Zheng FH, Wu ZH, Leung CKS (2018) Detection of Bruch's Membrane Opening in Healthy Individuals and Glaucoma Patients with and without High Myopia. *Ophthalmology* 125(10): 1537-1546.
 26. You QS, Peng XY, Xu L, Chen CX, Wang YX, et al. (2014) Myopic Maculopathy Imaged by Optical Coherence Tomography The Beijing Eye Study. *Ophthalmology* 121(1): 220-224.
 27. Gu YH, Ke GJ, Wang L, Gu QH, Zhou EL, et al. (2016) Risk factors of rhegmatogenous retinal detachment associated with choroidal detachment in Chinese patients. *Int J Ophthalmol* 9(7): 989-993.
 28. Conart JB, Selton J, Hubert I, Trechot F, El Adssi H, et al. (2014) Outcomes of Macular Hole Surgery with Short-Duration Positioning in Highly Myopic Eyes A Case-Control Study. *Ophthalmology* 121(6): 1263-1268.
 29. Vimont C (2017) Should You Be Worried About Blue Light.
 30. Gusukuma M, Kahhat R (2018) Electronic waste after a digital TV transition: Material flows and stocks. *Resour Conserv Recycl* 138: 142-150.
 31. Holms A, De Cuir CA (2005) Technical Research Report: The Cathode Ray Tube.

32. Coggins AB, Green WH, Marks MM, Manousos RM, Horenstein MG (2005) Basal cell carcinoma and World War II-era cathode ray oscilloscope exposure. *J Am Acad Dermatol* 52(S1): 1-7.
33. Hu PP, Kong Y, Chen B, Liu Q, Zhuo W, et al. (2017) Shielding Effect of Lead Glasses on Radiologists' Eye Lens Exposure in Interventional Procedures. *Radiat Prot Dosimetry* 174(1): 136-140.
34. Distler F, Hartmann J, Pahernik S, Wucherer M (2018) Response to Bieko re: Risk of Radiation-Induced Cataracts: Investigation of Radiation Exposure to the Eye Lens During Endourological Procedures. *J Endourol* 32(10): 905-906.

