



# Everything about Nutritional Value of the Meat Ingredients and How we can Reduce its Microbial Hazards

## Shaltout FAE\*

Department of food Hygiene and control, Benha University, Egypt

\*Corresponding author: Fahim Aziz Eldin Shaltout, Professor of Meat Hygiene, Department of food Hygiene and control, Benha University, Egypt, Email: fahim.shaltout@fvtn.bu.edu.eg

### Review Article

Volume 10 Issue 1

Received Date: November 20, 2024

Published Date: January 06, 2025

DOI: 10.23880/oajvsr-16000283

## Abstract

The Meat ingredients are a valuable part of the human diet as the meat contains essential elements such as meat protein, meat vitamins, and meat minerals. The meat ingredients products types are vulnerable to the microbial meat ingredients pathogens and the meat ingredients spoilage, posing significant risks to the public health and the social meat ingredients quality and safety. The Ionizing radiation method is used in the irradiated meat ingredients to maintain the meat ingredients safety and meat ingredients quality of the consumed meat ingredients products types specifically the beef meat ingredients. During recent decades, the irradiation method of the meat ingredients products reduces the microbial meat contamination and extends the storage period of the meat ingredients. The procedure entails exposing the meat ingredients products types to a regulated amount of the ionizing radiation method, mostly accomplished by applying the gamma rays method, the electron beams method, or the X rays method.. The radiation causes disrupts the DNA and other cellular components of the microbes contaminating the meat ingredients, making the microbes unable to reproduce and causing death of the microbial contamination. The procedure causes breaks down some of the molecules of the meat ingredients which can affect the meat nutritional quality and meat ingredients sensory characters.

**Keywords:** Beef Meat ingredients; DNA; The Gamma Rays Method; The Meat Ingredients Irradiation Method; The Human Health Hazard

## Abbreviations

FDA: Food and Drug Administration; TVBN: Total Volatile Base Nitrogen; TBA: Two Thiobarbituric Acid; WHC: Water Holding Capacity.

## Introduction

Despite the irradiation potential benefits, the irradiated meat ingredients products type's which remains controversially, with concerns to irradiation meat ingredients safety and quality, efficacy, and irradiation

effect on the meat ingredients nutritional quality and the meat ingredients sensory characters of meat ingredients products types. The irradiated meat ingredients products types could create the harmful compounds or destroy the essential nutrients. In contrast, others questioned the need for the irradiation method, considering other meat ingredients safety measures, such as the good manufacturing practices and the meat ingredients testing methods. The Consumer acceptance of the irradiated meat ingredients products types needs to be addressed, with some people expressing concerns about the meat ingredients safety and meat ingredients acceptability [1-6]. Our review article

aims to important evaluate the irradiated meat ingredients products types and its repercussions on the meat ingredients quality and the meat ingredients safety of the beef meat ingredients products types. To proof the meat ingredients irradiation method effectiveness at lowering the microbial meat ingredients contamination and improving the keeping quality of the beef meat ingredients is explored along with its potential effect on the physical and the chemical characteristics, the meat ingredients nutrient content, and the meat ingredients sensory characters. This review article address the regulatory framework for the irradiated meat ingredients types, including the consumed meat ingredients labeling requirements and the consumed meat ingredients government oversight, as well as identify the areas for more meat ingredients researches and meat ingredients policy development [7-12].

### **The Sources and Principles of the Irradiated Meat Ingredients Products Types**

The meat ingredients Ionizing radiation method such as gamma rays method, the X rays method, or the high energy electrons method, is used to irradiate the meat ingredients types. The irradiated meat ingredients products types is determined by the absorbed dose expressed in Gray (Gy) or kilo Gray (kGy), with one Gray being equal to one J/kg of product. The technique is considered a safe and effective way to decrease or eliminate the hazardous microbes, improving the keeping quality, as well as enhance the meat ingredients quality and safety of the meat ingredients types [13-18]. The principles of the irradiated meat ingredients products types are determined by the ability to disrupt the genetic material of microorganisms, preventing them from reproducing or causing illness. The irradiation method affects the microorganisms' genetic material (the DNA or the RNA) directly and indirectly. The Direct irradiation method can break the bonds between base pairs in the genetic material, killing the cell's reproduction ability. The damage to the water molecules creates free radicals and reactive oxygen, which damage genetic material indirectly. The Irradiation method helps to break down certain enzymes and meat ingredients proteins that can contribute to spoilage, thereby improving the of the meat ingredients [19-24]. The United State, Canada, as well as several European and Asian nations, allow the irradiated meat ingredients products types using the Cobalt 60 method, cesium 137 methods and the electron beam accelerators method. The Cobalt 60 method, the most prevalent source of the ionizing radiation method for the irradiated meat ingredients products types, is a radioactive isotope that emits the gamma rays capable of penetrating deep into the meat ingredients products types to destroy the harmful microorganisms. The Cesium 137 method is another source of the ionizing radiation method; it is less commonly used than cobalt 60. The electron beam accelerators are

used for the irradiated meat ingredients products types. The devices generate high energy electrons method that can penetrate the meat ingredients products types to eliminate the harmful microorganisms and improve the beef meat ingredients keeping quality [25-30]. The Irradiating meat ingredients products types have several benefits to the meat ingredients including multifunctional applications as well as guaranteed meat ingredients safety and meat security. The spectrum produced is effective against the bacterial spores across a broad range of concentrations. The processing does not involve heat, it is safe for the meat ingredients products types, does not significantly reduce the meat ingredients nutrient value, so leaves no chemical residues, and is simple to control during use to effectively lengthen the lifespan of the irradiated meat products types. The Radurization method uses low doses of 0.1-1 kGy [31-36]. This dose inhibits respiration, delays the ripening, disinfects the meat ingredients pests, and inactivates the Trichinella parasite. The Radicidation method is referred to as a moderate dose. The meat ingredients radiation uses a quantity of approximately 1-10 kGy, which has the effect of reducing the microbial meat ingredients spoilage and the microbial meat ingredients pathogens including the Salmonella species bacteria and the Listeria monocytogenes bacteria which contaminate the meat ingredients. In regarding to this dosage is typically found in the frozen meat ingredients products types and its application is identical to that of the pasteurization method, except irradiation method does not rely on the thermal energy method [37-42]. The Radapertization method uses extremely high doses which are above or equal to 10 kGy, ranging between 30 and 50 kGy. The irradiation dose is typically used in the sterilization process because its effect can kill all the contaminating microorganisms in the meat ingredients products types up to the level of the spores of the microorganisms. The irradiated meat ingredients products type's origin and the principles are based on the ability of the ionizing radiation method to disrupt the genetic material of the microorganisms, the enzymes, and the proteins in the meat ingredients products types, culminating in improved meat ingredients safety and meat ingredients quality. The use of irradiation method is regulated by the national and the international authorities to ensure its safety and effectiveness in the meat ingredients preservation [43-48].

### **The Action of Irradiation Method on the Irradiated Beef Meat Ingredients Products**

#### **The Microbial Safety of the Irradiated Beef Meat Ingredients Products**

The Microbial meat ingredients safety is important aspect of the beef meat ingredients production and consumption, as these meat ingredients products can be a source of the various harmful microorganisms that can cause the meat borne

illness. The beef meat ingredients products are potentially contaminated with microbial pathogens, such as Salmonella species bacteria, Escherichia coli bacteria, Campylobacter bacteria, and Listeria monocytogenes bacteria, leading to severe illness or death in vulnerable populations [49-54]. The meat ingredients contamination might occur at the production, processing, or distribution stage, including on the farm, during transportation, in slaughterhouses or processing facilities, and in meat ingredients retail outlets or at home. The Improper beef meat ingredients handling and the storage of the beef meat ingredients products can increase the risk of beef meat ingredients contamination. The meat borne illness outbreaks related to the beef meat ingredients have been reported in the world with various types of the meat ingredients products being implicated, including the ground beef meat ingredients products, the chicken meat ingredients products, the pork meat ingredients products, and the processed beef meat ingredients products.

The outbreaks have led to the significant public health hazards, the social meat ingredients quality and the economic consequences, the highlighting the importance of the effective interventions to reduce the risk of the meat ingredients contamination [55-60]. The Irradiation method has been studied extensively for its efficacy in reducing the microbial contamination of the beef meat ingredients. The exposing of the meat ingredients products types to the ionizing radiation method, the latter reduces or eliminates the harmful meat microorganisms that can cause meat borne illness and affect the social meat ingredients quality. The irradiation method could effectively reduce the levels of the meat ingredients microbial pathogens such as Salmonella species bacteria and Escherichia coli bacteria as well as levels of the meat ingredients spoilage bacteria leading to improved microbial meat ingredients safety and a reduced the risk of the meat borne illness and improve the social meat ingredients quality [61-66]. The effectiveness of the different types of the ionizing radiation method on the beef meat ingredients including the gamma rays method and the e beams method, has been used; the gamma ray irradiation method is more effective than the e beam meat irradiation method is at inhibiting the microbial growth in the beef meat ingredients. The UV light method effectively eliminates the Salmonella species bacteria, the Pseudomonas species bacteria, the Micrococcus species bacteria, and the Staphylococcus species bacteria on the beef meat ingredients. The keeping quality of the beef meat ingredients products types is improved by eliminating the microbial meat ingredients contaminant. The Gamma irradiation method at low doses can improve the microbiological meat ingredients safety and quality, ensure meat ingredients safety and quality, and extend the chicken meat's ingredients products ingredients keeping quality without affecting the meat ingredients quality .The 3 kGy gamma irradiated beef meat ingredients reduced the

growth of the mesophilic bacteria, coliforms bacteria, and the Staphylococcus aureus bacteria. The Food and Drug Administration (FDA) determined that a 3.5 kGy gamma ray irradiation method dose effectively eliminates the pathogenic microbes from the fresh beef meat ingredients and improve the social meat ingredients quality. The meat ingredients Irradiation method slows the growth of the bacterial cells and deactivating the bacterial metabolism. The Bacteria are inherently resistant to the action of the irradiation method and, in the lag phase or inactive state, will be more resistant. In contrast, the bacteria in the growth phase will be more effective [67-72].

### **The Chemical Characters of the Irradiated Beef Meat Ingredients Products Types**

The chemical characters of the irradiated beef meat ingredients refer to the changes that occur to the chemical constituents and the compositions of the meat ingredients products types due to exposure to the ionizing radiation method and affect the social meat ingredients quality. The Irradiation method can cause both the desirable and the undesirable action on the chemical characteristics of the beef meat ingredients, depending on the dose and the specific compounds in the meat ingredients products types [73-78]. The most significant changes often observed in the irradiated beef meat products is the formation of the free radicals. They become reactive molecules that damage cellular components and cause oxidative stress. Thus lipid oxidation, which causes off flavors and odors, as well as a decline in the nutritional meat ingredients quality due to the loss of essential fatty acids and other nutrients. The irradiation method at lower doses aids lipid oxidation by reducing the levels of peroxides and other reactive species. This procedure affects the meat ingredients protein content of the beef meat ingredients, leading to alterations in the composition of the amino acids, meat ingredients protein structure, and meat digestibility. The changes have potentially positive and negative action, mostly on the meat ingredients nutritional value and affect the social meat ingredients quality, that are contingent upon the particular meat ingredients proteins involved and the dose of radiation used. The positive action of the irradiation method include the fact that the irradiation method can cause the formation of reactive species, such as the free radicals, which can cause the formation of the covalent bonds between the amino acids in the meat ingredients protein molecules. The cross linking can change the structure of a meat ingredients protein molecule and make it resistant to enzymatic meat ingredients digestion, which causes a decrease in the meat ingredients protein digestibility [79-84]. The Irradiation method can cause the denaturation of the meat ingredients protein molecules. The Denaturation involves opening the meat ingredients protein structure, which can facilitate the interactions between the amino

acids and increase the accessibility of the digestive enzymes to meat ingredients protein molecules, and it can improve the meat ingredients protein digestibility. The irradiation method can cause adverse action; namely, the excessive irradiation method can cause a breakdown of or changes in the amino acid compounds in the meat ingredients protein molecules which cause a decrease in the overall amino acid content and consequently decrease the meat ingredients protein digestibility. The electron beam irradiation method at less than 3 kGy did not affect changes in the meat ingredients quality of the smoked duck flesh (the amino acids, the fatty acids, and the volatiles) during the storage [85-90]. The chemical changes, the irradiation method affects the vitamin content of the beef meat ingredients products types, with some vitamins being more sensitive than others. For example, the irradiation method leads to a loss of the vitamin C, while other vitamins, such as the vitamin A and E, are relatively stable.

The Irradiation method alter the beef meat ingredients oxidation–reduction ability, accelerating the lipid oxidation, the meat ingredients protein breakdown, and the flavor and the odor changes [91-96]. When combined with certain antioxidants, such as the flavonoids, the irradiation method can help prolong the induction period of the lipid oxidation., storing the irradiated beef meat ingredients at 5–10 C for one week almost did not change the pH, the texture, the total volatile base nitrogen (TVBN), or the microbe number. A higher dose of the UV irradiation method increased two thiobarbituric acid (TBA) content, decreased the water holding capacity (WHC), and the decreased the beef meat ingredients color intensity and the tenderness. The 2.5 and 5 kGy gamma irradiation method reduced the nitrite content in the chicken sausages and prevented the oxidation when combined with the antioxidants. The titratable acidity and the acid value in the beef meat ingredients can be reduced by the irradiation method. The beef meat ingredients contamination may occur at the production, the processing, or the distribution stage, including on the farm, during the transport, in the slaughterhouses or the processing facilities, and in the retail outlets or at the home [97-99].

## Conclusion

The Improper handling and the storage of the beef meat ingredients products can increase the risk of the beef meat ingredients contamination. The meat borne diseases outbreaks related to the beef have been reported globally and its effect on the social meat ingredients quality, with the various types of the meat ingredients products types being implicated, including the ground beef meat ingredients products types, the chicken meat ingredients products types, the pork meat ingredients products types, and the processed beef meat ingredients products types.

## References

1. Yemmireddy V, Adhikari A, Moreira J (2022) Effect of Ultraviolet Light Treatment on Microbiological Safety and Quality of Fresh Produce: An Overview. *Front Nutr* 9: 871243.
2. Saucier L (2016) Microbial Spoilage, Quality and Safety within the Context of Meat Sustainability. *Meat Sci* 120: 78-84.
3. Shaltout FA, Riad EM, Abou-Elhassan A (2017) Prevalence Of Mycobacterium Spp . In Cattle Meat And Offal's Slaughtered In And Out Abattoir. *Egyptian Veterinary medical Association* 77(2): 407-420.
4. Elaziz AO, Hassanin, Shaltout FA, Mohamed OA (2021) Prevalence of Some Foodborne Parasitic Affection in Slaughtered Animals in Local Egyptian Abattoir. *Journal of Nutrition Food Science and Technology* 2(3): 1-5.
5. Pereira, PMDCC, Vicente, AFDRB (2013) Meat Nutritional Composition and Nutritive Role in the Human Diet. *Meat Sci* 93: 586-592.
6. Elaziz AO, Hassanin FS, Shaltout FA, Mohamed OA (2021) Prevalence of some zoonotic parasitic affections in sheep carcasses in a local abattoir in Cairo, Egypt. *Advances in Nutrition & Food Science* 6(2): 25-31.
7. Saleh E, Shaltout F, Elaal EA (2021) Effect of some organic acids on microbial quality of dressed cattle carcasses in Damietta abattoirs, Egypt. *Damanhour Journal of Veterinary Sciences* 5(2): 17-20.
8. Edris A, Hassanin FS, Shaltout FA, Elbaba AH, Adel NM (2017) Microbiological Evaluation of Some Heat Treated Fish Products in Egyptian Markets. *EC Nutrition* 12(3): 124-132.
9. Borrego-Soto G, Ortiz-López R, Rojas-Martínez A (2015) Ionizing Radiation-Induced DNA Injury and Damage Detection in Patients with Breast Cancer. *Genet Mol Biol* 38: 420-432.
10. Edris AA, Hassanin FS, Shaltout FA, Elbaba AH, Adel NM (2017) Microbiological Evaluation of Some Heat Treated Fish Products in Egyptian Markets. *EC Nutrition* 12(3): 134-142.
11. Ragab A, Edris AM, Shaltout FAE, Salem AM (2022) Effect of titanium dioxide nanoparticles and thyme essential oil on the quality of the chicken fillet. *Benha Veterinary Medical Journal* 41(2): 38-40.
12. Mkhungo MC, Oyedeji AB, Ijabadeniyi OA (2018) Food Safety Knowledge and Microbiological Hygiene of

- Households in Selected Areas of Kwa-Zulu Natal, South Africa. *Ital J Food Saf* 7: 126-130.
13. Lima F, Vieira K, Santos M, Souza PM (2018) Effects of Radiation Technologies on Food Nutritional Quality. *IntechOpen: London, UK*, pp: 137-146.
  14. Shaltout FA, Amin R, Marionet Z, Nassif, AbdelW (2014) Detection of aflatoxins in some meat products. *Benha Veterinary Med J* 27(2): 368-374.
  15. Marin C, Cerda C, Bodi S, Rebenaque L, Vega S (2022) Research Note: Persistent Salmonella Problem in Slaughterhouses Related to Clones Linked to Poultry Companies. *Poult Sci* 101: 101968.
  16. Shaltout FA, Afify, Jehan EM, Abo E, Asmaa A (2012) Improvement of microbiological status of oriental sausage. *Journal of Egyptian Veterinary Medical Association* 72(2): 157-167.
  17. Castell P, Moreira RG (2021) Irradiation and Consumers Acceptance. *Innov. Food Process Technol A Compr Rev* 2: 122-135.
  18. Shaltout FA, Eldiasty E, Mohamed MS (2014) Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. *Animal Health Research Institute: First International Conference on Food Safety and Technology*, pp: 79-89.
  19. Erkmen O, Bozoglu TF (2016) Food Preservation by Irradiation. In *Food Microbiology: Principles into Practice*. John Wiley & Sons, Ltd. Hoboken, NJ, USA, pp. 106-126.
  20. Klurfeld DM (2018) What Is the Role of Meat in a Healthy Diet? *Anim Front* 8: 5-10.
  21. Hassanien FS, Shaltout FA, Fahmey MZ, Elsukkary HF (2020) Bacteriological quality guides in local and imported beef and their relation to public health. *Benha Veterinary Medical Journal* 39: 125-129.
  22. Bantawa K, Rai K, Limbu SD, Khanal H (2018) Food-Borne Bacterial Pathogens in Marketed Raw Meat of Dharan, Eastern Nepal. *BMC Res Notes* 11: 618.
  23. Khattab E, Shaltout F, Sabik I (2021) Hepatitis A virus related to foods. *Benha Veterinary Medical Journal* 40(1): 174-179.
  24. Saad SM, Shaltout FA, Farag AAA, Mohammed HF (2022) Organophosphorus Residues in Fish in Rural Areas. *Journal of Progress in Engineering and Physical Science* 1(1): 27-31.
  25. Saif M, Saad SM, Hassanin FS, Shaltout FA, Zaghoul M (2019) Molecular detection of enterotoxigenic *Staphylococcus aureus* in ready-to-eat beef products. *Benha Veterinary Medical Journal* 37 (2019): 7-11.
  26. Farag AA, Saad SM, Shaltout FA, Mohammed HF (2023 a) Studies on Pesticides Residues in Fish in Menofia Governorate. *Benha Journal of Applied Sciences* 8(5): 323-330.
  27. Farag AA, Saad SM, Shaltout FA, Mohammed HF (2023 b) Organochlorine Residues in Fish in Rural Areas. *Benha Journal of Applied Sciences* 8(5): 331-336.
  28. Shaltout FA, Hussein MN, Elsayed NK (2023) Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. *Journal of Advanced Veterinary Research* 13(2): 157-160.
  29. Shaltout FA, Heikal GI, Ghanem AM (2022) Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to *Aspergillus flavus* virulent factors. *benha veteriv medical journal veterinary* 42(1): 12-16.
  30. Shaltout FA, Salem RM, Eldiasty EM, Diab FA (2022) Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. *Journal of Advanced Veterinary Research* 12(5): 641-644.
  31. Shaltout FA, Eldiasty E, Salem R, Asmaa (2016) Mycological quality of chicken carcasses and extending shelf - life by using preservatives at refrigerated storage. *Vet M J* 62(3): 1-7.
  32. Shaltout FA, Salem R, Eldiasty E, Diab F (2016) Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Vet M J-Giza* 62(3): 9-14.
  33. Shaltout FA, Eldiasty E, Wafaa A (2015) Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate . *Benha Vet Med J* 28(2): 58-66.
  34. Shaltout FA, Gerges MT, Shewail AA (2018) Impact of Organic Acids and Their Salts on Microbial Quality and Shelf Life of Beef. *Assiut Veterinary Med J* 64(159): 164-177.
  35. Shaltout FA, Hashim M, Elnahas S (2015) Levels of some heavy metals in fish (*tilapia nilotica* and *Claris lazera*) at Menufia Governorate. *Benha Vet Med J* 29(1): 56-64.
  36. Shaltout FA, Zakaria IM, Nabil ME (2018) Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology* 2(5): 429-

- 438.
37. Shaltout FA, Diasty EM, Salem RM, Asmaa MA (2016) Mycological quality of chicken carcasses and extending shelf -life by using preservatives at refrigerated storage. *Vet Med J Giza* 62(3): 1-10.
  38. Shaltout FA, Salem R, Diasty E, Hassan W (2019) Effect of Lemon Fruits and Turmeric Extracts on Fungal Pathogens in Refrigerated Chicken Fillet Meat. *Global Veterinaria* 21(3): 156-160.
  39. Shaltout FA, Salem RM, Diasty E, Fatema H (2016) Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Vet Med J Giza* 62(3): 9-14.
  40. Shaltout FA, Ahmed AA, Maarouf E, Ahmed M (2018) Heavy Metal Residues in chicken cuts up and processed chicken meat products. *Benha Vet Med J* 34(1): 473-483.
  41. Shaltout FA, Hanan M, Ehsan AM, Edris (2020) Bacteriological examination of some ready to eat meat and chicken meals. *Biomed J Sci & Tech Res* 27(1): 20461-20465.
  42. Sobhy A, Shaltout F (2020) Prevalence of some food poisoning bacteria in semi cooked chicken meat products at Qaliubiya governorate by recent Vitek 2 compact and PCR techniques. *Benha Vet Med J* 38(2020): 88-92.
  43. Shaltout FA, Barr AAH, Abdelaziz ME (2022) Pathogenic Microorganisms in Meat Products. *Biomedical Journal of Scientific & Technical Research* 41(4): 32836-32843.
  44. Shaltout FA, Thabet MG, Koura HA (2017) Impact of Some Essential Oils on the Quality Aspect and Shelf Life of Meat. *J Nutr Food Sci* 7: 647.
  45. Shaltout FA, Mohammed IZ, Afify ESA (2020) Bacteriological profile of some raw chicken meat cuts in Ismailia city, Egypt. *Benha Veterinary Medical Journal* 39(2020): 11-15.
  46. Schevey CT, Toshkov S, Brewer MS (2013) Effect of Natural Antioxidants, Irradiation, and Cooking on Lipid Oxidation in Refrigerated, Salted Ground Beef Patties. *J Food Sci* 78: S1793-S1799.
  47. Shaltout FA, Mohammed IZ, Afify ESA (2020) Detection of *E. coli* O157 and *Salmonella* species in some raw chicken meat cuts in Ismailia province, Egypt. *Benha Veterinary Medical Journal* 39 (2020): 101-104.
  48. Shaltout FA, El-diasty EM, Asmaa- Hassan MA (2020) Hygienic Quality of Ready to Eat Cooked Meat in Restaurants at Cairo. *Journal of Global Biosciences* 8(12): 6627-6641.
  49. Shaltout FA, Nasief MZ, Lotfy LM, Gamil BT (2019) Microbiological status of chicken cuts and its products. *Benha Veterinary Medical Journal* 37: 57-63.
  50. Munir MT, Federighi M (2020) Control of Foodborne Biological Hazards by Ionizing Radiations. *Foods* 9: 878.
  51. Hassanin FS, Shaltout FA, Homouda SN, Arakeeb SM (2019) Natural preservatives in raw chicken meat. *Benha Veterinary Medical Journal* 37: 41-45.
  52. Hazaa W, Shaltout FA, Mohamed El-Shate (2019) Prevalence of some chemical hazards in some meat products. *Benha Veterinary Medical Journal* 37(2): 32-36.
  53. Ahn DU, Kim IS, Lee EJ (2013) Irradiation and Additive Combinations on the Pathogen Reduction and Quality of Poultry Meat. *Poult Sci* 92: 534-545.
  54. Hazaa W, Shaltout FA, Mohamed El-Shater (2019) Identification of Some Biological Hazards in Some Meat Products. *Benha Veterinary Medical Journal* 37(2): 27-31.
  55. Gaafar R, Hassanin FS, Shaltout FA, Zaghloul M (2019) Molecular detection of enterotoxigenic *Staphylococcus aureus* in some ready to eat meat-based sandwiches. *Benha Veterinary Medical Journal* 37(2): 22-26.
  56. Gaafar R, Hassanin FS, Shaltout FA, Zaghloul M (2019) Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubiya Governorate, Egypt. *Benha Veterinary Medical Journal* 37(2): 16-21.
  57. Ehlermann DAE (2014) Safety of Food and Beverages: Safety of Irradiated Foods. In: Motarjemi YBT, et al. (Eds.), *Encyclopedia of Food Safety*, Academic Press: Waltham, MA, USA, 3: 447-452.
  58. Saad SM, Shaltout FA, Elroos NAA, El-nahas SB (2019) Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. *J Food Sci Nutr Res* 2(1): 12-20.
  59. Nam KC, Jo C, Ahn DU (2016) Irradiation of Meat and Meat Products. In *Emerging Technologies in Meat Processing: Production, Processing and Technology*; John Wiley & Sons, Ltd. Hoboken, USA, pp: 7-36.
  60. Saad SM, Hassanin FS, Shaltout FA, Nassif MZ, Seif MZ (2019) Prevalence of Methicillin-Resistant *Staphylococcus Aureus* in Some Ready-to-Eat Meat Products. *American Journal of Biomedical Science & Research* 4(6): 460-464.

61. Oh H, Yoon Y, Yoon JW, Oh SW, Lee S, et al. (2023) Salmonella Risk Assessment in Poultry Meat from Farm to Consumer in Korea. *Foods* 12: 649.
62. Shaltout FA, EL-diasty EM, Mohamed MSM (2018) Effects of chitosan on quality attributes fresh meat slices stored at 4 C. *Benha Veterinary Medical Journal* 35(2): 157-168.
63. Ham YK, Kim HW, Hwang KE, Song DH, Kim YJ, et al. (2017) Effects of Irradiation Source and Dose Level on Quality Characteristics of Processed Meat Products. *Radiat Phys Chem* 130: 259-264.
64. Hassanzadeh P, Tajik H, Rohani SMR, Moradi M, Hashemi M, et al. (2017) Effect of Functional Chitosan Coating and Gamma Irradiation on the Shelf-Life of Chicken Meat during Refrigerated Storage. *Radiat Phys Chem* 141: 103-109.
65. Indiarito R, Pratama AW, Sari TI, Theodora HC (2020) Food Irradiation Technology: A Review of the Uses and Their Capabilities. *SSRG Int J Eng Trends Technol* 68: 91-98.
66. Shaltout FA, El Shatter MA, Fahim HM (2019) Studies on Antibiotic Residues in Beef and Effect of Cooking and Freezing on Antibiotic Residues Beef Samples. *Scholarly Journal of Food and Nutrition* 2(1): 1-4.
67. Indiarito R, Qonit MAH (2020) A Review of Irradiation Technologies on Food and Agricultural Products. *Int J Sci Technol Res* 9: 4411-4414.
68. Shaltout FA, Zakaria IM, Nabil ME (2018) Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology* 33(2): 429-438.
69. Arvanitoyannis, I.S. Consumer Behavior toward Irradiated Food. In *Irradiation of Food Commodities: Techniques, Applications, Detection, Legislation, Safety and Consumer Opinion*; Arvanitoyannis, I.S.B.T.-I., Ed.; Academic Press: Boston, MA, USA, 2010; pp. 673-698.
70. Shaltout FA, Maarouf AAA, Elkhoully MES (2017) Bacteriological Evaluation of Frozen Sausage. *Nutrition and Food Toxicology* 1(5): 174-185.
71. Fajardo-Guerrero M, Rojas-Quintero C, Chamorro-Tobar I, Zambrano C, Sampedro F, et al. (2020) Exposure Assessment of Salmonella Spp. in Fresh Pork Meat from Two Abattoirs in Colombia. *Food Sci Technol Int* 26(1): 21-27.
72. Shaltout FA, El-Toukhy EI, Abd El-Hai MM (2019) Molecular Diagnosis of Salmonellae in Frozen Meat and Some Meat Products. *Nutrition and Food Technology Open Access* 5(1): 1-6.
73. Shaltout FA, AM Ali, SM Rashad (2016) Bacterial Contamination of Fast Foods. *Benha Journal of Applied Sciences (BJAS)* 1(2): 45-51.
74. Singh R, Singh A (2019) Food Irradiation: An Established Food Processing Technology for Food Safety and Security. *Defence Life Science Journal* 4(4): 206-213.
75. Shaltout FA, Zakaria IM, Eltanani J, Elmelegy AS (2015) Microbiological status of meat and chicken received to University student hostel. *Benha Veterinary Medical Journal* 29(2): 187-192.
76. Yeh Y, Moura FHD, Broek VD, Mello ASD (2018) Effect of Ultraviolet Light, Organic Acids, and Bacteriophage on Salmonella Populations in Ground Beef. *Meat Science* 139: 44-48.
77. Saad SM, Edris AM, Shaltout FA, Edris, Shimaa (2012) Isolation and identification of salmonellae and *E.coli* from meat and poultry cuts by using A. multiplex PCR. *Benha Vet Med J special* 16-26.
78. Rastogi RP, Richa, Kumar A, Tyagi MB, Sinha RP (2010) Molecular Mechanisms of Ultraviolet Radiation-Induced DNA Damage and Repair. *Journal of Nucleic Acids*.
79. Saad SM, Shaltout FA, Elroos NAA, El-Nahas SB (2019) Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. *Journal of Food Science and Nutrition Research* 2(1): 12-20.
80. Gomez I, Janardhanan R, Ibañez FC, Beriain MJ (2020) The Effects of Processing and Preservation Technologies on Meat Quality: Sensory and Nutritional Aspects. *Foods* 9(10): 1416.
81. Saad SM, Shaltout FA, Elroos NAA, El-nahas SB (2019) Incidence of Staphylococci and *E. coli* in Meat and Some Meat Products. *EC Nutrition* 14(6).
82. Shaltout FA, Riad EM, TES Ahmed, AbouElhassan AA (2017) Studying the Effect of Gamma Irradiation on Bovine Offal's Infected with *Mycobacterium tuberculosis* Bovine Type. *Journal of Food Biotechnology Research* 1(6): 1-5.
83. Shahi S, Khorvash R, Goli M, Ranjbaran SM, Najarian A, et al. (2021) Review of Proposed Different Irradiation Methods to Inactivate Food-Processing Viruses and Microorganisms. *Food Science and Nutrition* 9(10): 5883-5896.
84. Shaltout FA, Zakaria IM, Nabil ME (2018) Incidence

- of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology* 2(5): 429-438.
85. Food and Drug Administration, HHS (2012) Irradiation in the Production, Processing and Handling of Food. Final Rule. Fed Regist 73(164): 49593-49603.
  86. Shaltout A, El-diasty M, El-mesalamy M, El-shaer M (2014) Study on fungal contamination of some chicken meat products with special reference to the use of PCR for its identification. Conference, *Veterinary Medical Journal* 60(2): 1-22.
  87. Bintsis T (2017) Foodborne Pathogens. *AIMS Microbiol* 3(3): 529-563.
  88. Shaltout FA, Thabet MG, Koura HA (2017) Impact of some essential oils on the quality aspect and shelf life of meat. *Benha Veterinary Medical Journal* 33(2): 351-364.
  89. Park JG, Yoon Y, Park JN, Han IJ, Song BS, et al. (2010) Effects of Gamma Irradiation and Electron Beam Irradiation on Quality, Sensory, and Bacterial Populations in Beef Sausage Patties. *Meat Sci* 85: 368-372.
  90. Shaltout FA, Mohammed F, Hosam A, Ibrahim A, Mostafa EM (2017) Afifi4.2017: Incidence of Coliform and *Staphylococcus aureus* in ready to eat fast foods. *Benha Veterinary Medical Journal* 32(1): 13-17.
  91. Shaltout FA, Zakaria IM, Nabil ME (2017) Detection and typing of *Clostridium perfringens* in some retail chicken meat products. *Benha Veterinary Medical Journal* 33(2): 283-291.
  92. Maherani B, Hossain F, Criado P, Fadhel Y, Salmieri S, et al. (2016) World Market Development and Consumer Acceptance of Irradiation Technology. *Foods* 5: 79.
  93. Amiri A, Zandi H, Khosravi HM (2019) Effect of Electron Beam Irradiation on Survival of *Escherichia Coli* O157:H7 and *Salmonella Enterica* Serovar *Thyphimurium* in Minced Camel Meat during Refrigerated Storage. *J Food Qual Hazards Control* 6: 174-178.
  94. Song BS, Lee Y, Park JH, Kim JK, Park HY, et al. (2018) Toxicological and Radiological Safety of Chicken Meat Irradiated with 7.5 MeV X-rays. *Radiat. Phys Chem* 144: 211-217.
  95. Souza C, Apaolaza V, Hartman, P, Brouwer AR, Nguyen N (2021) Consumer Acceptance of Irradiated Food and Information Disclosure-A Retail Imperative. *J Retail Consum Serv* 63: 102699.
  96. Lianou A, Panagou EZ, Nychas GJE (2017) Meat Safety-I Foodborne Pathogens and Other Biological Issues. In Lawrie et al. *Meat Science: 8<sup>th</sup> (Edn.)*, Woodhead Publishing: Cambridge, UK, pp: 521-552.
  97. Sobhy, Asmaa, Shaltout, Fahim (2020) Detection of food poisoning bacteria in some semi-cooked chicken meat products marketed at Qaliubiya governorate. *Benha Vet Med J* 38(2020): 93-96.
  98. Shaltout FA (2024) Abattoir and Bovine Tuberculosis as a Remerging Foodborne Disease. *Clinical Medical Reviews and Report* 6(1): 1-7.
  99. Shaltout FA (2023) Viruses in Beef, Mutton, Chevron, Venison, Fish and Poultry Meat Products. *Food Science & Nutrition Technol* 8(4): 1-10.