



Study of Most Affected Age Group of Anemia among Pregnant Women Attending at El-Nuhud Teaching Hospital, West Kordufan State, 2019

Alnour UBAA^{1*} and Hamad GGA^{†2}

¹Lecture, West Kordofan University, Sudan

²Assistant Professor, West Kordofan University, Sudan

*Corresponding author: Gurashi Gabr Alla Hamad, Faculty of Public & Environmental Health, West Kordofan university, Sudan, Email: khobaib.gor@gmail.com

†Equally contributed towards this article.

Research Article

Volume 8 Issue 1

Received Date: April 08, 2024

Published Date: May 15, 2024

DOI: 10.23880/phoa-16000285

Abstract

Background: Anemia is a common problem in Africa, with prevalence ranging from 21.1% to 64.4%. According to WHO estimates, 56 million pregnant women were anemic in 2005, more than half of them (57.6%) would be in Africa.

Objectives: The main objective of this study to determine the most affected age group of anemia among pregnant women attending at El-Nuhud Teaching Hospital, West Kordufan State, 2019.

Methodology: A cross-sectional descriptive hospital-based study was carried out from August to November 2019. The sample size was total coverage of all pregnant women attending at El-Nuhud Teaching Hospital, West Kordufan State. A number of 420 samples of data of structured administrated questionnaire and blood samples were collected from pregnant women. The blood samples were collected and investigated by laboratory technician by using Colorimeter to measure hemoglobin level. The data were analyzed by using Statistical Package for Social Science (SPSS), version 20 and Chi-Square test (X²) used to check difference between variables, p value less than 0.05 was considered significant.

Results: The results indicated the high affected in illiteracy (77.6%) followed by primary (57.1%), Khalwa (50%), secondary (42.8%), university (26.7%) and post university (0%). This significant relation between educational level for pregnant woman and anemia among pregnant women, (p value = 0.000)..

Conclusion: The study concluded that the prevalence of anemia among pregnant women was more than half. The study recommended to implement health education programmes at all levels, use appropriate vaccinations, supplementations and vitamins during pregnancy to prevent anemia, family planning, optimum maternal and child health, and environmental Sanitation.

Keywords: Pregnant Women; Anemia; El-Nuhud Teaching Hospital

Abbreviation: WHO: World Health Organization.

Introduction

Anemia is common in both developing and industrialized nations. It affects about two billion people worldwide, i.e 30% of the inhabitants of the planet, according to data from

the World Health Organization (WHO). Hence it is considered a public health problem. It greatly affects a person's life, especially in its more severe forms. Anemia cause tiredness, weakness, malaise, and making routine tasks more difficult and painful. Anemia is a common problem in Africa, with prevalence ranging from 21.1% to 64.4%. It has significant impact on morbidity and mortality. The affected person may

be drowsy, with less resistance to physical activities and deterioration in cognitive function and immunity [1].

Anemia defined as low blood hemoglobin concentration has shown to be public health problem that affects low-, middle- and high-income countries and has significant adverse impacts on social and economic development [2].

Globally anemia affects 32.4 million (38.4%) of pregnant women it is major public health problem in South Asia (48.7%) and Africa (46.3%) [3].

Pregnant woman's anemia is a critical issue of public health due to its frequency, multifactorial nature and impact on pregnancy evolution. According to WHO estimates, 56 million pregnant women were anemic in 2005, more than half of them (57.1%) would be in Africa. It contributes to the 20% of maternal deaths related to indirect causes [4]. Anemia of pregnancy an important risk factor for fetal and maternal morbidity is considered a global health problem affecting almost 50% of pregnant women Achebe MM, et al. [5] Anemia is one of nutritional insufficiency disorder and affecting in Ethiopia the prevalence of anemia using a cut off level hemoglobin less than 11 g/dl (33% haematocrit) was 23.5% (72/ 306) [6]. In Africa the prevalence rate of anemia among antenatal women ranges from 35% to 72% while in Asia 37% to 75% [7]. Anemia is a global public health problem, but is more prevalent in pregnant women and young as stated by WHO [2]. Anemia among pregnant women worldwide was 38% and Sudan among countries considered with a moderate type of anemia during pregnancy [8].

In a cross – sectional study was conducted among pregnant women in Erbil city. The result of the study was showed the prevalence of manemia among pregnant women 46.2% [9]. In India anemia is the second most common cause and accounting for 20% of total maternal, the prevalence of anemia ranges from 33% to 89% among pregnant women and is more than women from 60% among adolescent girls with wide variations in different regions of countries Rajamouli j, et al. In fact over 20% of pregnant women in Europe are anemic during pregnancy [10]. The United Kingdom prevalence of maternal anemia in the antenatal period was estimated as 24 % in a recent cross-sectional study [11].

Justifications and Problem State

Anemia is most common medical disorder in pregnancy and postpartum period [12]. Anemia in pregnancy is an important health issue resulting in high maternal morbidity and mortality [13]. The World Health Organization (WHO) has estimated that prevalence of anemia in pregnant women was found 14% in developed, 50% in developing countries

and 60-75% in India. This study conducted among pregnant women attending at ElNuhud Teaching Hospital -West Kordufan State in Sudan to determine the most affected age group with anemia the daily frequency of pregnant women who attending hospital about (10-15) pregnant women and monthly about (300-450) pregnant women El-Nuhud Teaching Hospital, 2018. Prevalence of anemia in females was higher in most regions and age groups South Asia and Central, West and East Sub-Saharan Africa had the highest burden, while East, Southeast and South Asia saw the greatest reduction [14].

Study of pregnant Sudanese women attended delivery in River Nile State Sudan the study revealed that (64.3%) of study population were anemic [12]. Study carried out in Ethiopia among pregnant women revealed that (34.6%) were anemic [15].

Surveys in different parts of India indicate that about 50-60% of women belonging to low socio-economic group are anemic in last trimester of pregnancy [16].

Objective

To determine the most affected age group of anemia among pregnant women attending at El Nuhud Teaching Hospital, West Kordufan State, Sudan-2019

Material & Method

Study Design: A descriptive cross-sectional hospital-based study. Study area: El-Nuhud Teaching Hospital, West Kordufan State, Sudan.

Location: Study conducted at El-Nuhud Teaching Hospital, West Kordufan State - Sudan, 2019. El-Nuhud locality located between latitude 9.25-16.3o North and latitude 27-32o East it surrounded by many areas in the East Alkhewi locality and Abuzabad locality, in the West Wadbanda locality and in the South South Kordufan State, West Kordufan University, 2018.

Building: El-Nuhud Teaching Hospital building were from stable material such as bricks and stone El-Nuhud Teaching Hospital, 2019.

Health services: Regarding to health services there are 2governmental hospitals, 6 health centers, 12 private dispensaries and 8 private clinic El-Nuhud locality, 2018.

Education services: Education services in El-Nuhud locality include basic; higher secondary schools and universities; 194 basic, 15 higher secondary schools and two universities Education office, 2018.

Study Population: All pregnant women attending at El-Nuhud Teaching Hospital that fulfill the inclusion criteria

during the data collection period was considered as study participants. The number population (327051) persons in El-Nuhud locality, (192961) of them as women, (78819) as age reproductive women and (13115) as pregnant women [17].

Sample Size: The sample size was determined by total coverage of pregnant women attending at El-Nuhud Teaching Hospital –West Kordufan State during three months (August-November 2019) to obtained 420 pregnant.

Methodology: Data collected using following tools:

Questionnaire: In the form of administrated questionnaire filled by pregnant women attending at El-Nuhud Teaching Hospital, West Kordufan State. The questionnaire includes the following variables:

-Socio-demographic factors (age, educational status, her husband education, her occupation, her husband occupation, family size, Gestation period and income).

Laboratory Investigation: Hemoglobin level determined using Colorimeter hemoglobin and all laboratory investigations were done by technician's laboratory. Pregnant women was considered anemic when her hemoglobin less than 11gm/dl according (WHO).

Data Analysis: Data entered and analyzed by using computer-based operation practically (Statistic Package for Social Science) version 20 and Chi - Square test (χ^2) was used to check the differences between variables. The (P.value) less than 0.05 considered significant.

Ethnical consideration: Permission was taken from the pregnant women with her full consent and permission was obtained from the Ministry of Health, West Kordufan State, also permission was taken from the Hospital Medical Manger as well as an ethnical approval from the Department of Epidemiology, Faculty of Public and Environmental Health, University of Khartoum.

Results

Anemia Educational level	Positive		Negative		Total	
	N	%	N	%	N	%
Illiteracy	132	78	38	22	170	41
Khalwa	1	50	1	50	2	0.4
Primary	92	57	69	43	161	38
Secondary	24	43	32	57	56	13
University	8	27	22	73	30	7.2
Post university	0	0	1	100	1	0.2
Total	257	61	163	39	420	100

Table 1: Educational level among pregnant women in relation to anemia at El-Nuhud Teaching Hospital, West Kordufan State, 2019.

(n= 420)

$\chi^2 = 45.163$ p = 0.000

Table 1 shows high affected in illiteracy (77.6%) followed by primary (57.1%), Khalwa (50%), secondary (42.8%), university (26.7%) and post university (0%). This significant

relation between educational level for pregnant woman and anemia among pregnant women, (p value = 0.000).

Anemia	Positive		Negative		Total	
	N	%	N	%	N	%
Her job						
Housewife	241	63	144	37	385	92
Worker	15	60	10	40	25	6
Professional	1	10	9	90	10	2.3
Total	257	61	163	39	420	100

Table 2: Pregnant woman's job among pregnant women in relation to anemia at El-Nuhud Teaching Hospital, West Kordufan State, 2019.

(n = 420)

$\chi^2 = 11.370$ p= 0.003

Table 2 shows high affected with housewife (62.6%) followed by worker (60%) and professional (10%). This

significant relation between pregnant women's job and anemia among pregnant women, (p value = 0.003).

Anemia	Positive		Negative		Total	
	N	%	N	%	N	%
Gestation period (months)						
First trimester (1-3 month)	16	46	19	54	135	8.3
Second trimester (4-6 month)	58	50	58	50	116	28
Third trimester (7 - 9 month)	183	68	86	32	269	64
Total	257	61	163	39	420	100

Table 3: Gestation period among pregnant women in relation to anemia at El-Nuhud Teaching Hospital, West Kordufan State, 2019.

(n=420)

$\chi^2 = 14.945$ p= 0.001

Table 3 shows high affected in third trimester 7-9 months (68%) followed by second trimester 4-6 months (50%) and first trimester 1-3 months (45.7%). This significant relation between gestation period and anemia among pregnant women, (p value = 0.001).

Discussion

This a cross-sectional descriptive hospital-based study was conducted at El-Nuhud Teaching, West Kordufan State with an objective to study the prevalence of anemia among pregnant women in period (August- November) 2019.

This study shows that (77.6%), (57.1%), (50%), (42.8%) and (26.7%) anemic of pregnant women's educational level, (illiteracy, primary, Khalwa, secondary and university) respectively, a significant associated between pregnant women's educational level and anemia (p value = 0.000) and this similar with study of Nwizu EN, et al. [18] he revealed that had statistically association with low education attainment and anemia among pregnant women (p value = 0.006), and also agree with study of Jufar AH, et al. [19] they revealed had significant association between education status and anemia in pregnancy, (p value = 0.037).

The study shows that statistically significant association between her husband's educational level and anemia in pregnancy, (p value = 0.000) and this agree with study of Krupp K, et al. [20] they revealed anemia in pregnant women was associated with lower education among spouses (p value = 0.021).

This study shows that (61.6%), (60%) and (10%) anemic of pregnant women's job, (housewife, worker and professional) respectively, statistically significant between pregnant women's job and anemia in pregnancy, (p value =

0.003), this similar with study of Krupp K, et al. [20] they carried out anemia in pregnancy was associated with lower household income (p value = 0.022).

This study shows that (68%), (50%) and (45.7%) anemic of pregnant women their gestation period (7-9 months), (4-6 months) and (1-3 months) respectively, a significant associated between gestation period and anemia among pregnant women (p value = 0.001) and this similar with study of Selim NAA, et al. [21] they revealed that had statistically significant association with gestation period and in pregnancy, (p value = 0.001), also similar with study of Khan NAM, et al. [22] they revealed significantly associated between gestation period and anemia in pregnancy (p value < 0.05), also agree with the study of Melku M, et al. [23] they revealed that anemia was high at third trimester (18.9%), also similar with the study of Jufar AH, et al. [19] they revealed had significant associated between gestation period (third trimester, p value = 0.041) and anemia in pregnancy and similar with the study of Alene KA, et al. [24] they carried out pregnant women in the third and second trimesters were 3.32 times more likely to be 58 affected by anemia as compared to pregnant women in the first trimester.

Conclusions

This a cross-sectional descriptive hospital-based study was conducted with an objective to study prevalence of anemia among pregnant women attending at El-Nuhud Teaching Hospital, West Kordufan State, 2019. The prevalence among pregnant women (61%). The most affected age group was more than 45years old (100%).

The more causes associated with anemia among pregnant women were age group, family size, latrine inside house, pregnant women's educational level, her husband

educational level, pregnant women's job, gestation period, parity, stillbirth history, abortion history, complication during pregnancy, problem health, meals per day poultry eggs intake, fruits and vegetables intake, follow up during pregnancy and folic acid and iron intake during pregnancy.

Recommendation

- Implementation of health education programmes at all levels, especially nutritional education to change dietary habits and feeding practices.
- Provision adequate of quality and quantity of nutrients and food during pregnancy
- Encourage family planning
- Provision universal, perfect, affordable and accessible antenatal care services.

References

1. Omer R, Ahmed B (2015) Determination of iron deficiency anemia in infants in Ahmed Gasim Hospital, Khartoum North (KN), Sudan in the Period from January to June 2011. *International Journal of Innovative Research in Science, Engineering and Technology* 4(9): 9272-9275.
2. WHO (2015) The global prevalence of anemia in 2011. Geneva, Switzerland.
3. Tadesse SE, Seid O, Mariam YG, Fekado A, Wasihum Y, et al. (2017) Determinants of anemia among pregnant women attending Antenatal Care in Dessie Town health facilities, Northern Central Ethiopia, unmatched case – control study. *PLOS ONE* 12(3): e0173173.
4. Salifou K, Obossou AAA, Sidi IR, Hounkpatin Bib, Hounkponou ANF (2015) Factors associated with anemia in pregnant women in 2014 in Parakou. *Journal of Pregnancy Child Health* 2: 174.
5. Achebe MM, Gvili AG (2017) How I treat anemia in pregnancy: Iron, cobalamin and folate. *The American Society of Hematology* 129(8): 940-949.
6. Zekarias B, Meleko A, Hayder A, Nigatu A, Yetagessu T (2017) Prevalence of Anemia and its associated factors among pregnant women attending Antenatal Care (ANC) In Mizan-Tepi University Teaching Hospital, South West Ethiopia. *Health Science Journal* 11(5): 529.
7. Kumari P, Rani M, Saini N, Aparna (2017) Prevalence of anemia among pregnant mothers admitted for delivery at Distinct Hospital Ambala, Haryana: An Observational Study. *International Journal of Pharmaceutical Science Review and Research* 47(1): 27-30
8. ALamain AA, Taha FAN (2016) Incidence and causes of anemia during pregnancy in Antenatal Wards in Khartoum University Hospitals. *The Journal of Middle East and North Africa Sciences* 2(12): 1-7.
9. Abdulwahid RG, Ahmed HM (2017) Prevalence and severity of anemia among pregnant women in Primary Health Centers / Erbil City. *Kurdistan Journal of Applied Research (KJAR)* 2(2): 155.
10. Colman C, Pavord S (2017) Oxford University Hospital NHS Foundation, OUH.
11. Rukuni R, Knight M, Murphy MF, Roberts D, Stanworth SJ (2015) Screening for Iron deficiency and Iron deficiency in pregnancy: a structured review and gap analysis against UK national screening criteria. *BMC Pregnancy and Childbirth* 15: 269.
12. Elnazir FA (2015) Incidence and factors of anemia among women attending delivery in River Nile State Sudan.
13. Anlaakuu P, Anto F (2017) Anemia in pregnancy and associated factors: a cross sectional study of antenatal attendants at the Sunyani Municipal Hospital, Ghana. *BMC Res Notes* 10(1): 402.
14. Namazi A (2018) Anemia and iron supplement in pregnancy: A Mini review. *JOJ Nursing & Health Care* 7(3): 678-680.
15. Nega D, Dana D, Tera T, Eshetu T (2015) Anemia associated with asymptomatic malaria among pregnant women in rural surroundings of Arba Minch, South Ethiopia. *BMC Research Notes* 8: 110.
16. Park JE (2015) Preventive and social medicine In: 23th (Edn.), M/S Banaridas Bhanot Publishers 1167, prem Nagar Jabalpur, India, pp: 642.
17. Ministry of health (2019) West Kordufan State Records.
18. Nwizu EN, Iliyasu Z, Ibrahim SA, Galadanci HS (2011) Sociodemographic and maternal factors in anemia in pregnancy at booking in Kano, Northern Nigeria. *Afr J Reprod Health* 15(4): 33-41.
19. Jufar AH, Zewde T (2014) Prevalence of anemia among pregnant women attending Antenatal Care at Tikur Anbessa Specialized Hospital, Addis Ababa Ethiopia. *Journal of Hematology & Thromboembolic Disease*.
20. Krupp K, Placek CD, Wilcox M, Ravi K, Srinivas V, et al. (2018) Financial decision-making power is associated with moderate to severe anemia: A prospective cohort study among pregnant women in Rural South India. *Midwifery* 61: 15-21.

21. Selim NAA, Al-Mass M, Al-Kuwari M, Ismail MS (2016) Assessment of anemia, IDA and ID among pregnant in Qatar: Cross sectional survey. *SM Journal of Public Health and Epidemiology* 2(3): 1035.
22. Khan NAM, Sonkar VRK, Dimple VK, Inamdar IA (2017) Study of anemia and its associated risk factors among pregnant women in a Rural Field Practice Area of a Medical College. *National Journal of Community Medicine* 8(7): 396-400.
23. Melku M, Addis Z, Alem M, Enawgaw B (2014) Prevalence and predictors of maternal anemia during pregnancy in Gondar, Northwest Ethiopia: An institutional based cross-sectional study. *Journal of Immunology Research*.
24. Alene KA, Dohe AM (2014) Prevalence of Anemia and associated factors among pregnant women in an Urban Area of Eastern Ethiopia. *Hindawi* pp: 7.